

APPENDIX 11A

CORE RESIDENCE TIMES

APPENDIX 11A – CORE RESIDENCE TIMES

The derivation of the core residence times for circulating crud as shown in Subsection 11.1.3 is as follows:

Circulating Crud

The number of radioactive atoms (N_f) in the crud film on in-core surfaces at any time is:

$$\frac{dN_f}{dt} = \sum_i \phi - \lambda_i N_f \quad (\text{Eq. 11A-1})$$

Solving for N_f yields the following:

$$N_f = \frac{\sum_i \phi}{\lambda_i} (1 - e^{-\lambda_i t_{\text{res}}}), \text{ atoms/g} \quad (\text{Eq. 11A-2})$$

Where:

$\sum_i \phi$ = activation rate for each isotope i, d/g-sec

λ_i = decay constant for each isotope i, sec^{-1}

t_{res} = desired core residence time, seconds

The number of radioactive atoms (N_c) released to the reactor coolant at any time is:

$$\frac{dN_c}{dt} = N_f \{ER\} A_c - (\alpha + \beta + \lambda_i) N_c, \text{ atoms/sec}$$

Solving for N_c yields the following:

$$N_c = \frac{N_f \{ER\} A_c}{(\alpha + \beta) + \lambda_i} (1 - e^{-(\alpha + \beta + \lambda_i)t}) \quad (\text{Eq. 11A-3})$$

Where:

$\{ER\}$ = erosion rate, $\text{g/cm}^2\text{-sec}$

A_c = core surface area, cm^2

α = plateout rate, sec^{-1}

APR1400 DCD TIER 2

β = purification cleanup rate, sec^{-1}

λ_i = decay constant, sec^{-1}

Total amount of crud (M_c) released to the reactor coolant at any time is:

$$\frac{dM_c}{dt} = \{ER\}A_T - (\alpha + \beta)M_c \quad (\text{Eq. 11A-4})$$

Where M_c includes both radioactive and nonradioactive material.

Solving for M_c yields:

$$M_c = \frac{\{ER\}A_T}{\alpha + \beta} (1 - e^{-(\alpha + \beta)t}), \text{ grams} \quad (\text{Eq. 11A-5})$$

Where:

$\{ER\}$ = erosion rate, $\text{g/cm}^2\text{-sec}$

A_T = total system area, cm^2

α = plateout rate, sec^{-1}

β = purification cleanup rate, sec^{-1}

The activity (A_i) of the crud released to the reactor coolant is:

$$A_i = \frac{\lambda_i N_c}{M_c}, \text{ Bq/g-crud} \quad (\text{Eq. 11A-6})$$

Substituting the values of N_c and M_c into the above expression and assuming λ_i is small compared to α and β , the activity of the crud is as follows:

$$A_i = \sum_i \phi (1 - e^{-\lambda_i t_{\text{res}}}) \frac{A_c}{A_T}, \text{ Bq/g-crud} \quad (\text{Eq. 11A-7})$$

This activity (A_i) is also assumed to be the activity of the crud that plates out on out-of-core surfaces.

Solving Equation (11A-7) for t_{res} yields Eq. 11.1-8.

Deposited Crud

The activity (A_j) of the deposited crud is:

$$A_j = \lambda_j N_f = \sum_j \phi (1 - e^{-\lambda_j t_{\text{res}}}) \quad (\text{Eq. 11A-8})$$

Solving Eq. 11A-8 for t_{res} yields Eq. 11.1-9.