

Table 2.6-2 Representative Stratigraphic Section – Marsland Expansion Area

Elevation (ft amsl)	Average Depth (ft bgs)	Group	Formation & Member (Schultz and Stout 1955)			Formation and Member (Revised)		References (Revised)
Varying 4150 -4,380	15 - 135	Arikaree Group	Monroe Creek Formation			Upper Harrison Beds		Swinehart et al. (1985)
						Monroe Creek-Harrison Formation		
			Gering Formation			Gering Formation		
Varying 4,140 - 4.020	135 - 285	White River Group	Brule Formation	Whitney Member		Brule Formation	"Brown Siltstones"	LaGarry (1998)
				Orella Member	Orella D		Whitney Member	
					Orella C		Orella Member	
					Orella B			
					Orella A			
4,020 – 3,890	285 – 650		Chadron Formation	Upper Chadron	Chadron C	Chadron Formation	Big Cottonwood Creek Member	Terry (1998) Terry and LaGarry (1998)
	Upper/Middle Chadron			Chadron B				
3,890 – 3,380	650 -925			Middle Chadron	Chadron A	Chamberlain Pass Formation	Peanut Creek Member	Terry (1998) Terry and LaGarry (1998)
3,380 - 3,180	925 – 1,025			Upper Interior Paleosol			Upper Interior Paleosol	Terry (199)
				basal sandstone of the Chadron Formation			Channel Sandstone	Terry (1998) Terry and LaGarry (1998)
				Montana Group	Pierre Shale	Interior Paleosol		Pierre Shale
3,180 – 3,130	1,025 - ? (Bottom not seen in logs)	Pierre Shale				Pierre Shale	Terry (1998) Terry and LaGarry (1998)	

Notes:

- 1) The Shultz and Stout conventions for Formation & Member are utilized throughout this document, with the exception of the Red Clay Horizon, which is referred to as the Upper Interior Paleosol.
- 2) Topsoil, colluvial and alluvial deposits are not shown, but are Quaternary in age and range in thickness from 0 to 30 ft-bgs.
- 3) ft amsl = feet above mean sea level; ft bgs = feet below ground surface
- 4) Elevations are representative averages for MEA only, and based on Log M-1252.