


## **SUPPLEMENTAL EXPERT OPINION REGARDING THE RENEWAL OF ISL URANIUM MINING (CROW BUTTE RESOURCES) NEAR CRAWFORD, NEBRASKA**

Hannan E. LaGarry, Ph. D.  
1002 ½ East 3rd Street, Chadron NE 69337  
hlagarry@olc.edu

| United States Nuclear Regulatory Commission Official Hearing Exhibit              |            |  |                       |
|---|------------|--|-----------------------|
| In the Matter of:   |            | CROW BUTTE RESOURCES, INC.   |                       |
|   |            | (License Renewal for the In Situ Leach Facility, Crawford, Nebraska) |                       |
|  | ASLBP #:   | 08-867-02-OLA-BD01   | Identified: 8/18/2015 |
|   | Docket #:  | 04008943   | Withdrawn:            |
|   | Exhibit #: | INT-013-00-BD01  | Stricken:             |
|   | Admitted:  | 8/18/2015  |                       |
|   | Rejected:  |  |                       |
|   | Other:     |  |                       |

### **INTRODUCTION**

Beginning in 2008 I have provided expert opinions to the Western Nebraska Resources Council and Owe Aku (among others) regarding Crow Butte Resources' ISL uranium mining near Crawford, Nebraska. Since that time, I have offered expert opinions on Crow Butte Resources' proposed expansions, and initiated or participated in several scientific studies intended to further clarify the issues I raised in those written opinions. I am offering this supplemental expert opinion in support of the Consolidated Interveners' new and revised contentions regarding the license renewal of the Crow Butte Resources ISL uranium mine near Crawford, Nebraska. I am concerned that issues regarding the lack of lixiviant containment raised in earlier opinions have not been addressed and also apply to the Consolidated Petitioners' contentions in this particular case. My goal in offering this expert opinion is not to protest uranium mining, but rather to express my concerns regarding threats to the region's diminishing water supplies and the future inhabitability of northwestern Nebraska and adjacent southwestern South Dakota. In this document I will briefly explain the basis for my concerns.

### **PROFESSIONAL BACKGROUND**

I have 25 years' experience studying the geology of northwestern Nebraska and adjacent South Dakota. Following dissertation work on the regional geology from 1988-1995, from 1996-2006 I led teams of geologists from the Nebraska Geological Survey that mapped in detail the geology of most of northwestern Nebraska (a total of 80 1:24,000 quadrangles). The completion of this work frequently required detailed study of equivalent strata in adjacent Fall River, Shannon, and Todd counties in South Dakota. These maps, including digital versions (ArcInfo) and supporting field notes, are available from the University of Nebraska-Lincoln School of Natural Resources (contact Dr. Matthew Joeckel, Director). As a direct consequence of this mapping, I have published peer-reviewed articles on the Chadron Formation (Terry & LaGarry 1998), the Brule Formation (LaGarry 1998), the mapping of surficial deposits (Wysocki & others 2000, 2005), and local faults (Fielding & others 2007). From 2006-2008 I continued this work as an Adjunct Professor of Geology at Chadron State College (CSC) in Chadron, Nebraska. During this time I worked with and advised students studying the region's groundwater, surface water, and faults (Balmat & others 2008, Butterfield & others 2008). Since 2008 I have worked as an Instructor and Researcher in the Department of Math, Science, & Technology at Oglala Lakota College (OLC MST), serving the department as Co-Chair from 2009-2012 and Chair from 2012-2013. Since 2013 I have resumed my duties as a full-time instructor and researcher.

In addition to my ongoing geological work in Nebraska, I have been working with students and faculty to study the geology, groundwater, surface water, and heavy metal contaminants of southwestern South Dakota and the Pine Ridge Reservation. For the past 6 years our research has been funded by the National Science Foundation's Tribal Colleges and Universities Program

and Experimental Program for Stimulating Competitive Research, and the USDA National Institute for Food and Agriculture Tribal College Equity Program. We have formed and maintained partnerships with Chadron State College, the South Dakota Geological Survey, the South Dakota School of Mines and Technology, South Dakota State University, the University of Illinois Urbana-Champaign, the University of Illinois Center for Advanced Materials Purification of Water Systems, the Department of Health Physics at the University of Michigan School of Nuclear Engineering, the University of Washington Native American Research Center for Health, and the Technological University of Darmstadt, Germany. I have authored or co-authored reports detailing the preliminary results of studies describing toxic heavy metal contamination of drinking water (Salvatore & others 2010, Botzum & others 2011), characteristics of local aquifers (Gaddie & LaGarry 2010, LaGarry & others 2012), potential uranium contamination risk to communities on the Pine Ridge Reservation (LaGarry & Yellow Thunder 2012), and the transmission of uranium-contaminated water along regional faults (Bhattacharyya & others 2012).

## **THE CONCERNS**

My concerns regarding the Crow Butte resources ISL uranium mine are the lack of confinement resulting from secondary porosity in the form of faults and joints, the problem of artesian flow, and the horizontal flow of water beyond the uranium-bearing strata. In my 2008 opinion on Crow Butte Resources' ISL mining I described in detail the stratigraphy of the region and potential contamination pathways, and established the plausibility of contaminants migrating away from the mine site into adjacent areas and aquifers. In this supplemental opinion I will concentrate on work that has been completed after 2008. It is beyond the scope of this opinion to review the entire scientific literature for the region, but I provide the most readily available recent research. Where appropriate, I also refer to Crow Butte Resources' FINAL ENVIRONMENTAL ASSESSMENT FOR THE LICENSE RENEWAL OF U.S. NUCLEAR REGULATORY COMMISSION LICENSE NO. SUA-1534.

### ***The problem of secondary porosity***

In order for ISL mining to be considered safe, the mined uranium-bearing strata must be isolated from rocks above and below by confining layers. Confining layers must be continuous, unfractured, and unperforated in order for containment to exist. There are three principal pathways through which contaminated water could migrate away from the uranium-bearing strata through adjacent confining layers (described in detail below): 1) secondary porosity in the form of joints and faults, 2) thinning or pinching out of confining layers, and 3) perforations made by improperly cased or capped wells. I will restrict my comments to situation (1), as situations (2) and (3) have been addressed in my earlier opinions and there is insufficient new data to warrant additional discussion.

Secondary porosity, in the form of intersecting faults and joints, is common in all of the rocks north, east, and south of the Black Hills Dome and along the Pine Ridge Escarpment (see Swinehart & others 1985). These faults and joints are generally oriented NW-SE and SW-NE, and are a result of the ongoing uplift of the Black Hills of southwestern South Dakota. Although many people consider the Black Hills uplift to have ended by the late Cretaceous Period (~65 Ma), the Black Hills were tectonically active in the late Eocene (Evans & Terry 1994), and continued to fault, fracture, and fold the rocks of northwestern Nebraska and southwestern South Dakota into the middle Miocene (Fielding & others 2007). Based on numerous small

earthquakes along the Sandoz Ranch-Whiteclay Fault, the area is tectonically active to this day (McMillan & others 2006). These earthquakes are relatively mild, and don't significantly damage surface infrastructure. However, even small earthquakes represent shifting and flexing of the earth's crust, and are continuously creating, closing, and redistributing the secondary porosity of the region's rocks and changing the flow pathways of the region's groundwater. This means that joints incapable of transmitting water one day may be able to transmit water at a later date. These faults and fractures transect all major bedrock units of the region. These faults connect the uranium-bearing strata to adjacent aquifers as well as modern river alluvium.

Preliminary studies of the interaction of these faults with surface waters in the region show that creeks that provide municipal water supplies can be entirely consumed and redirected by the region's secondary porosity. Chadron Creek, the stream that supplies water to the city of Chadron, Nebraska, went dry for the first time in the city's history. Subsequent study of the creek's water flow rates by Chadron State College students suggested that normal amounts of water are flowing from springs, but the water is disappearing into deeper alluvium or into fractures in the rock (Balmat & others 2008, Butterfield & others 2008). Following these observations, a Chadron State College graduate student studied the lineaments of northwestern Nebraska and southwestern South Dakota using data collected by high-flying aircraft, satellites, and the space shuttle, and showed that these represent widespread faults (Balmat & Leite 2008). Many of the faults in northwestern Nebraska and southwestern South Dakota persist for tens of miles (Diffendal 1994, Fielding & others 2007). Also, many of the ancient river deposits of the Tertiary strata, along with the alluvium deposited by modern rivers such as the Cheyenne River, the White River, and Hat Creek, follow fault zones because fractured rock erodes more easily.

Crow Butte Resources' license renewal application asserts that fault zones are known in the vicinity of the license area. The most notable of these is the White River Fault, which trends along the axis of the syncline in which the uranium-bearing Chamberlain Pass Formation (being mined for uranium in this case) was deposited (see LaGarry and LaGarry 1997, LaGarry 1998). A review of the scientific literature shows that faults and joints are well known in rocks surrounding the Black Hills, and are known to interconnect major aquifers and the land surface (Swinehart & others 1985, Peters & others 1988, Fielding & others 2007). In earlier expert opinions, I described the extensive, detailed geologic mapping of the region conducted by the Nebraska Geological Survey (Table 1), and predicted that faults and joints would be capable of transmitting uranium-contaminated waters from depth onto the land surface. In 2012 my colleagues and I reported preliminary research showing uranium-contaminated artesian springs along the Sandoz Ranch-Whiteclay Fault in Fall River and Shannon counties (Bhattacharyya & others 2012), supporting those assertions. Based on the numerous studies reported here and the maps cited in Table 1, the absence of joints and faults in the vicinity of the proposed mine is a false perception, because joints and faults are ubiquitous in this region.

Despite being obvious when viewed from Earth's orbit (Balmat and Leite 2008), these joints and faults are difficult to observe when covered by Holocene surficial deposits. Crow Butte Resources' assertions that hydrologic modeling shows no evidence of faults is misleading, as it is possible to conduct such tests in ways that are unlikely to reveal the presence of a fault. In order to demonstrate a lack of containment, faults must be mapped and wells installed along them. Test wells can be configured to determine the hydrologic head along a fault relative to adjacent

strata, and measurements of water transmission capacity along faults can be made. Balmat and Leite (2008) and the “Whistleblower letter” (see my 2008 opinion) both reported faults intersecting the Crow Butte Resources mine license area (at about the location of the processing plant), and initial testing could be made there and preliminary measurements made. Additional faults could be tested in the vicinity of Red Cloud Buttes north of Crawford in the Smiley Canyon 1:24,000 Quadrangle and at in the Trunk Butte 1:24,000 quadrangle (Table 1). Joints are ubiquitous in the region, and many are available for testing.

It’s worth noting that despite being made aware of this geological mapping and the published stratigraphic work that resulted from it, Crow Butte Resources continues to use outdated 1960’s era concepts and nomenclature in their FINAL ENVIRONMENTAL ASSESSMENT FOR THE LICENSE RENEWAL OF U.S. NUCLEAR REGULATORY COMMISSION LICENSE NO. SUA–1534. They continue to refer to the uranium-bearing strata as “Basal Chadron Formation” rather than the currently used Chamberlain Pass Formation. If Crow Butte Resources is unable to follow easy to address simple scientific protocols such as using the currently accepted stratigraphic nomenclature, I am concerned about their diligence in addressing challenging issues such as the mapping of faults and joints. Reliance on outdated data, concepts, and assumptions about the geology of the region (including, but not limited to rocks folding, rather than fracturing) will make any and all modeling of the hydrology of the area meaningless.

**Table 1. Open-file (OFM) maps from the Alliance 1:250,000 Quadrangle available from the Nebraska Geological Survey (University of Nebraska-Lincoln Conservation & Survey Division, School of Natural Resources). All maps are Nebraska quadrangles except as noted.**

| Map Name <sup>1</sup>  | Date <sup>2</sup> | Map No. <sup>3</sup> | Authors                       |
|------------------------|-------------------|----------------------|-------------------------------|
| Sherrill Hills (WY-NE) | 1997              | 53                   | LaGarry & LaGarry             |
| Story                  | 1997              | 54                   | LaGarry & LaGarry             |
| Story NE               | 1997              | 55                   | LaGarry & LaGarry             |
| Lone Tree Ranch        | 1997              | 56                   | LaGarry & LaGarry             |
| Wayside                | 1997              | 57                   | LaGarry & LaGarry             |
| Bohemian Creek         | 1997              | 58                   | LaGarry & LaGarry             |
| Kirtley (WY-NE)        | 1997              | 59                   | LaGarry & LaGarry             |
| Warbonnet Ranch        | 1997              | 60                   | LaGarry & LaGarry             |
| Bodarc                 | 1997              | 61                   | LaGarry & LaGarry             |
| Five Points            | 1997              | 62                   | LaGarry & LaGarry             |
| Whitney                | 1997              | 63                   | LaGarry, LaGarry, & Swinehart |
| Trunk Butte            | 1997              | 64                   | LaGarry, LaGarry, & Swinehart |
| Chadron West           | 1997              | 65                   | LaGarry, LaGarry, & Swinehart |
| Crawford               | 1998              | 66                   | LaGarry, Swinehart, & LaGarry |
| Crow Butte             | 1998              | 67                   | LaGarry, Swinehart, & LaGarry |
| Chimney Butte          | 1998              | 68                   | LaGarry, Swinehart, & LaGarry |
| Coffee Mill Butte      | 1998              | 69                   | LaGarry, Swinehart, & LaGarry |
| Dead Mans Creek        | 1998              | 70                   | LaGarry, Swinehart, & LaGarry |
| Belmont                | 1998              | 71                   | LaGarry, Swinehart, & LaGarry |
| Coffee Mill Butte SW   | 1998              | 72                   | LaGarry, Swinehart, & LaGarry |
| Coffee Mill Butte SE   | 1998              | 73                   | LaGarry, Swinehart, & LaGarry |
| Van Tassell (WY-NE)    | 1999              | 74                   | LaGarry, LaGarry, & Swinehart |
| Harrison West          | 1999              | 75                   | LaGarry, LaGarry, & Swinehart |
| Harrison East          | 1999              | 76                   | LaGarry, LaGarry, & Swinehart |
| Andrews                | 1999              | 77                   | LaGarry, LaGarry, & Swinehart |
| Smiley Canyon          | 1999              | 78                   | LaGarry, LaGarry, & Swinehart |
| Van Tassell SE (WY-NE) | 1999              | 79                   | LaGarry, LaGarry, & Swinehart |
| Harrison SW            | 1999              | 80                   | LaGarry, LaGarry, & Swinehart |
| Harrison SW            | 1999              | 81                   | LaGarry, LaGarry, & Swinehart |
| Kyle Creek             | 1999              | 82                   | LaGarry, LaGarry, & Swinehart |

|                                  |      |     |                               |
|----------------------------------|------|-----|-------------------------------|
| Glen                             | 1999 | 83  | LaGarry, LaGarry, & Swinehart |
| Isinglass Buttes                 | 2000 | 84  | LaGarry, LaGarry, & Swinehart |
| Chadron NE                       | 2000 | 85  | LaGarry, LaGarry, & Swinehart |
| Chadron East                     | 2000 | 86  | LaGarry, Swinehart, & LaGarry |
| Bordeaux                         | 2000 | 87  | LaGarry, Swinehart, & LaGarry |
| Kings Canyon/Chadron 3 NW        | 2000 | 88  | LaGarry, Swinehart, & LaGarry |
| Hay Springs Creek/Chadron 3 NE   | 2000 | 89  | LaGarry, Swinehart, & LaGarry |
| Sand Canyon West/Chadron 3 SW    | 2000 | 90  | LaGarry, Swinehart, & LaGarry |
| Sand Canyon East/Chadron 3 SE    | 2000 | 91  | LaGarry, Swinehart, & LaGarry |
| Beaver Wall                      | 2001 | 92  | LaGarry, Swinehart, & LaGarry |
| Whiteclay                        | 2001 | 93  | LaGarry, Swinehart, & LaGarry |
| Whiteclay SW                     | 2001 | 94  | LaGarry, Swinehart, & LaGarry |
| Whiteclay SE                     | 2001 | 95  | LaGarry, Swinehart, & LaGarry |
| Hay Springs                      | 2001 | 96  | LaGarry, Swinehart, & LaGarry |
| Hay Springs NE                   | 2001 | 97  | LaGarry, Swinehart, & LaGarry |
| Hay Springs SW                   | 2001 | 98  | LaGarry, Swinehart, & LaGarry |
| Spring SE                        | 2001 | 99  | LaGarry, Swinehart, & LaGarry |
| Montrose                         | 2001 | 108 | LaGarry & LaGarry             |
| Orella                           | 2001 | 109 | LaGarry & LaGarry             |
| Wolf Butte                       | 2001 | 110 | LaGarry & LaGarry             |
| Roundtop                         | 2001 | 111 | LaGarry, LaGarry, & Swinehart |
| Horn                             | 2001 | 112 | LaGarry & LaGarry             |
| Clinton NW                       | 2002 | 113 | LaGarry, Swinehart, & LaGarry |
| Clinton NE                       | 2002 | 114 | LaGarry, Swinehart, & LaGarry |
| Clinton SW                       | 2002 | 115 | LaGarry, Swinehart, & LaGarry |
| Clinton                          | 2002 | 116 | LaGarry, Swinehart, & LaGarry |
| Rushville SE                     | 2002 | 117 | LaGarry, Swinehart, & LaGarry |
| Rushville SW                     | 2002 | 118 | LaGarry, Swinehart, & LaGarry |
| Rushville NE                     | 2002 | 119 | LaGarry, Swinehart, & LaGarry |
| Rushville                        | 2002 | 120 | LaGarry, Swinehart, & LaGarry |
| Gordon NW                        | 2003 | 121 | LaGarry, Swinehart, & LaGarry |
| Hog Island                       | 2003 | 122 | LaGarry, Swinehart, & LaGarry |
| Gordon                           | 2003 | 123 | LaGarry, Swinehart, & LaGarry |
| Gordon SE                        | 2003 | 124 | LaGarry, Swinehart, & LaGarry |
| South of Gordon                  | 2003 | 125 | LaGarry, Swinehart, & LaGarry |
| Coburn Canyon                    | 2003 | 126 | LaGarry, Swinehart, & LaGarry |
| Bovee Valley West/Rushville 3 SW | 2003 | 127 | LaGarry, Swinehart, & LaGarry |
| Bovee Valley East/Rushville 3 SE | 2003 | 128 | LaGarry, Swinehart, & LaGarry |
| Whistle Creek NW                 | 2004 | 129 | LaGarry, Swinehart, & LaGarry |
| Whistle Creek NE                 | 2004 | 130 | LaGarry, Swinehart, & LaGarry |
| Marsland NW                      | 2004 | 131 | LaGarry, Swinehart, & LaGarry |
| Marsland                         | 2004 | 132 | LaGarry, Swinehart, & LaGarry |
| Box Butte Reservoir West         | 2004 | 133 | LaGarry, Swinehart, & LaGarry |
| Box Butte Reservoir East         | 2004 | 134 | LaGarry, Swinehart, & LaGarry |
| Box Butte NW                     | 2005 | 137 | LaGarry, Swinehart, & LaGarry |
| Box Butte NE                     | 2005 | 138 | LaGarry, Swinehart, & LaGarry |
| Skunk Lake NW                    | 2005 | 139 | LaGarry, Swinehart, & LaGarry |
| Skunk Lake NE                    | 2005 | 140 | LaGarry, Swinehart, & LaGarry |
| Smith Lake                       | 2005 | 141 | LaGarry, Swinehart, & LaGarry |
| Twin Lakes NE                    | 2005 | 142 | LaGarry, Swinehart, & LaGarry |
| Dolly Warden Lake                | 2005 | 143 | LaGarry, Swinehart, & LaGarry |
| Billys Lake                      | 2005 | 144 | LaGarry, Swinehart, & LaGarry |

---

**Notes for Table 1:**

- 1. Where two names are given, the former name is listed first, and the current name second**
- 2. Date indicates when map went on open file.**
- 3. Numbers were assigned by the Conservation & Survey Division**

### ***The problem of artesian flow***

Artesian flow occurs when there is a hydrologic connection, through faults or highly permeable strata, between groundwater sources high on the landscape and the land surface lower down. The weight of water in overlying strata exerts pressure downward into water within the uranium-bearing strata, which can then be released as artesian water flow (like a fountain) where the topographically lower uranium-bearing strata is exposed at the surface, or where it is punctured by drilling. Artesian flow was observed by Crow Butte Resources and was acknowledged in their FINAL ENVIRONMENTAL ASSESSMENT FOR THE LICENSE RENEWAL OF U.S. NUCLEAR REGULATORY COMMISSION LICENSE NO. SUA-1534. Artesian flow is most likely where the upper confining layer is perforated by secondary porosity (faults and joints), poorly constructed or improperly sealed exploration wells, or thinning or absence of upper confining layers. Artesian flow could transmit lixiviant, the most toxic mineral-laden of waters, onto the land surface (and into White River alluvium) and discharge large amounts of contaminants into connected aquifers or faults in a very short time.

### ***The problem of horizontal flow***

Confining layers above and below uranium-bearing strata limit the unwanted spread of contaminants from an ISL site. However, horizontal flows within the uranium-bearing strata and along intersecting faults and joints are also of concern. Such flow can rapidly redirect lixiviant or mine waste away from the mine site and into unexpected breaches in the confining layers. In my 2008 opinion I cited research on the transmission of water along secondary porosity (faults and joints) in the Brule Formation of up to 1500 feet/day. I also provide detailed, plausible conditions under which this contamination could spread to adjacent aquifers. Crow Butte acknowledges such secondary porosity in the Brule Formation (but not the faults that create it) in their FINAL ENVIRONMENTAL ASSESSMENT FOR THE LICENSE RENEWAL OF U.S. NUCLEAR REGULATORY COMMISSION LICENSE NO. SUA-1534. This acknowledged secondary porosity, if breached by unconfined lixiviant, would transmit contaminants to the major, mapped faults north of the Pine Ridge in Nebraska in only a few years, and from there into adjacent regions.

## **CONCLUDING REMARKS**

Based on the arguments presented above, it is my expert opinion that ISL mining in the area surrounding Crawford, Nebraska cannot be adequately contained. Reports of artesian flow, the acknowledged and prevalent jointing and faulting leading to widespread secondary porosity, along with potentially high horizontal flow through regional faults indicate that during the course of its operation the Crow Butte Resources ISL uranium mine will most likely contaminate the region with unconfined lixiviant. This contamination will pollute and render unusable ground and surface water southwards into Nebraska and surface waters within the White River drainage northeastwards into greater South Dakota. Also, based on my reading of Crow Butte Resources' FINAL ENVIRONMENTAL ASSESSMENT FOR THE LICENSE RENEWAL OF U.S. NUCLEAR REGULATORY COMMISSION LICENSE NO. SUA-1534, no comprehensive review of the geologic literature was conducted despite my criticisms on that issue in the 2008 opinion. In my view, the use of outdated scientific literature, or in this case, a general lack of review of recent study, should not be seen as an opportunity to operate in a knowledge vacuum. Much of the Great Plains region was studied prior to the 1980's and the general acceptance of Plate Tectonics Theory, and therefore generally misrepresents the geologic setting of the region. This was true

of the geologic literature used to justify ISL mining near Crawford, Nebraska, and is also true of the data used to justify proposed mining near Edgemont, South Dakota. Crow Butte Resources' complete and continued reliance of modeling based on outdated and incomplete parameters will never be sufficient to satisfy concerns based on actual mapping and recent published studies. It is incumbent upon potential ISL operators, as it is with any natural resource extractors, to seek out the most recent research and expert opinions on the geological settings in which they propose to operate.

## REFERENCES

- Balmat, J. L. & M. B. Leite. 2008. Identification of geological structures from lineaments on remotely sensed images of the Black Hills-Pine Ridge region. Proceedings of the 118<sup>th</sup> Annual Meeting of the Nebraska Academy of Sciences, p.28.
- Balmat, J. W., J. L. Balmat, M. J. Culver, A. C. Butterfield, C. A. Kaiser, J. Zweibel, H. E. LaGarry, & M. B. Leite. 2008. Preliminary geology of the Chadron Creek watershed, Dawes County, Nebraska. Proceedings of the 118th Annual Meeting of the Nebraska Academy of Sciences, p.64.
- Bhattacharyya, P., K. Converse, J. W. Ejnick, H. E. LaGarry, & A. Riesen. 2012. Studying uranium contamination levels in groundwater from the Pine Ridge Reservation, South Dakota: a community-university partnership. Geological Society of America Abstracts With Programs 44(6):77.
- Botzum, C. J., J. W. Ejnick, K. Converse, H. E. LaGarry, & P. Bhattacharyya. 2011. Uranium contamination in drinking water in the Pine Ridge Reservation, southwestern South Dakota. Geological Society of America Abstracts with Programs 43(5):125.
- Butterfield, A. C., J. L. Balmat, J. W. Balmat, K. Young, M. Culver, C. A. Kaiser, J. Zweibel, M. B. Leite, & H. E. LaGarry. 2008. Discharge of Chadron Creek, Dawes County, Nebraska. Proceedings of the 118th Annual Meeting of the Nebraska Academy of Sciences, pp. 64-65.
- Diffendal, R. F. 1994. Geomorphic and structural features of the Alliance 10 x 20 Quadrangle, western Nebraska, discernible from synthetic-aperture radar imagery and digital shaded-relief maps. Rocky Mountain Geology 30(2):137-147.
- Evans, J. E. & D. O. Terry, Jr. 1994. The significance of incision and fluvial sedimentation in the basal White River Group (Eocene-Oligocene), badlands of South Dakota, U.S.A. Sedimentary Geology 90:137-152.
- Fielding, C. R., H. E. LaGarry, L. A. LaGarry, B. E. Bailey, & J. B. Swinehart. 2007. Sedimentology of the Whiteclay gravel beds in northwestern Nebraska, USA: structurally controlled drainage promoted by early Miocene uplift of the Black Hills Dome. Sedimentary Geology 202:58-71.
- Gaddie, H. & H. E. LaGarry. 2010. Preliminary description of newly recognized Oligocene and early Miocene strata near Wanblee, South Dakota. Geological Society of America Abstracts with Programs 42(3)14.
- LaGarry, H. E. 1998. Lithostratigraphic revision and redescription of the Brule Formation (White River Group), northwestern Nebraska, pp. 63-92 in (D. O. Terry, Jr., H. E. LaGarry, & R. M. Hunt, Jr., eds.) Depositional environments, lithostratigraphy, and biostratigraphy of

- the White River and Arikaree Groups (late Eocene to early Miocene, North America). Geological Society of America Special Paper 325, 216 p.
- LaGarry, H. E. & L. A. LaGarry. 1997. Geology of the Montrose, Orella, Wolf Butte, Roundtop, and Horn 7.5' USGS quadrangles, Sioux and Dawes counties, Nebraska. University of Nebraska-Lincoln Conservation and Survey Division Open-file Report 48, 161 p.
- LaGarry, H. E. & L. A. LaGarry. 2010. Proposed lithostratigraphic revision, redescription, and redefinition of the White River Group (Eocene-Oligocene), South Dakota. Geological Society of America Abstracts with Programs 42(3):14.
- LaGarry, H. E., C. Belile, & H. Gaddie. 2012. Revised lithostratigraphic correlation of the Arikaree Group from northwestern Nebraska to southwestern South Dakota. Proceedings of the 122nd Annual Meeting of the Nebraska Academy of Sciences, pp. 92- 93.
- McMillan, M. E., P. L. Heller, & S. L. Wing. 2006. History and causes of post-Laramide relief in the Rocky Mountain Orogenic Plateau. Geological Society of America Bulletin 118(3/4):393-405.
- Peter, K. D., K. E. Kolm, & T. C. Nichols, Jr. 1988. Lineaments: significance, criteria for determination, and varied effects on ground-water systems – a case history in the use of remote sensing, pp. 46-68 in (A. I. Johnson & C. B. Petersson, eds.) Geotechnical Applications of Remote Sensing and Remote Data Transmission. American Society for Testing and Materials, Philadelphia, Pennsylvania.
- Rothrock, E. P. 1931. The Cascade Anticline. South Dakota Geological Survey Reports of Investigations 8, 25 p.
- Rothrock, E. P. 1931. The Chilson Anticline. South Dakota Geological Survey Reports of Investigations 9, 26 p.
- Rothrock, E. P. 1948. Structures in Fall River County. South Dakota Geological Survey Reports of Investigations 62, 64 p.
- Salvatore, M., H. E. LaGarry, P. Lee, & L. One Feather. 2010. Preliminary report: isolation and detection of filterable contaminants in drinking water, Pine Ridge Reservation, South Dakota. Geological Society of America Abstracts with Programs 42(3):44.
- Swinehart, J. B., V. L. Souders, H. M. DeGraw, & R. F. Diffendal Jr. 1985. Cenozoic paleogeography of western Nebraska, pp. 209– 229 in (R. M. Flores & S. S. Kaplan, eds.), Cenozoic Paleogeography of West-Central United States. Proceedings of Rocky Mountain Paleogeography Symposium 3, Society of Economic Paleontologists and Mineralogists. Denver, Colorado.
- Terry, D. O., Jr. & H. E. LaGarry. 1998. The Big Cottonwood Creek Member: a new member of the Chadron Formation in northwestern Nebraska, pp. 117-142 in (D. O. Terry, Jr., H. E. LaGarry, & R. M. Hunt, Jr., eds.) Depositional environments, lithostratigraphy, and biostratigraphy of the White River and Arikaree Groups (late Eocene to early Miocene, North America). Geological Society of America Special Paper 325, 216 p.
- Terry, D. O., Jr. 1998. Lithostratigraphic revision and correlation of the lower part of the White River Group: South Dakota to Nebraska, pp. 15-38 in (D. O. Terry, Jr., H. E. LaGarry, & R. M. Hunt, Jr., eds.) Depositional environments, lithostratigraphy, and biostratigraphy of the White River and Arikaree Groups (late Eocene to early Miocene, North America).



Geological Society of America Special Paper 325, 216 p.

Wysocki, D. A., P. J. Schoeneberger, & H. E. LaGarry. 2005. Soil Surveys: a window to the subsurface. *Geoderma* 126:167-180.

Wysocki, D. A., Schoeneberger, & H. E. LaGarry. 2000. Geomorphology of soil landscapes, pp. E5-E39 in (M. E. Sumner, editor-in-chief) *CRC Handbook of Soil Science*. Chemical Rubber Company Press.