

AN ECOLOGICAL STUDY
OF THE SUSQUEHANNA RIVER
IN THE VICINITY OF
THE THREE MILE ISLAND NUCLEAR STATION

Annual Report For 1976

by

George A. Nardacci, Project Leader, and Associates
Ichthyological Associates, Inc.
P.O. Box 223, Ellers, Pennsylvania 17319

for

METROPOLITAN EDISON COMPANY

POOR ORIGINAL

ICHTHYOLOGICAL ASSOCIATES, INC.
Edward C. Raney, Ph.D., Director
301 Forest Drive, Ithaca, New York 14850

February 1977

1565 038

7911010

754

AN ECOLOGICAL STUDY OF THE SUSQUEHANNA RIVER
IN THE VICINITY OF THE THREE MILE ISLAND NUCLEAR STATION

Annual Report For 1976

by

George A. Nardacci, Project Leader, and Associates

Ichthyological Associates, Inc.
P. O. Box 223, Etters, Pennsylvania 17319

POOR ORIGINAL

for

METROPOLITAN EDISON COMPANY

ICHTHYOLOGICAL ASSOCIATES, INC.
Edward C. Raney, Ph.D., President
301 Forest Drive, Ithaca, New York 14850

February 1977

1565 039

TABLE OF CONTENTS

Section		Page
1.0	INTRODUCTION	1
1.1	SUMMARY	2
1.1.1	Fish	2
1.1.2	Macroinvertebrates	2
1.1.3	Ambient Water Quality	4
1.1.3.1	Physicochemical Parameters	4
1.1.3.2	Fecal Coliform	4
1.1.4	Thermal Plume Mapping	4
1.1.5	Effects Of Cooling Tower Salt Drift On Agricultural Crops And Natural Vegetation	5
1.1.5.1	Plant Pathogens	5
1.1.5.2	Quantitative Vegetation Studies	5
2.0	FISH	
	Robert W. Malick, Jr.	
2.1	Methods	6
2.2	Results	9
2.2.1	Trapnet	9
2.2.2	Seine	11
2.2.3	Reproductive Status And Condition Factor	13
2.2.4	Parasites	14
2.3	Discussion	15
3.0	MACROINVERTEBRATES	
	John H. Epler, III, Janet L. Polk	
3.1	Methods	86
3.1.1	Taxonomic Treatment	87
3.1.2	Data Analysis	90
3.2	Results	92
3.2.1	Density	93
3.2.2	Biomass	94
3.2.3	Diversity (D)	94
3.2.4	Percent Similarity Composition	94
3.2.5	Analysis Of Variance	94
3.3	Discussion	97
4.0	AMBIENT WATER QUALITY	
4.1	Methods	156
4.2	Results	156
4.2.1	PHYSICOCHEMICAL PARAMETERS	156
	Eric S. McClellan, James H. Smith	
4.2.2	FECAL COLIFORM	158
	Linda M. Wike	
5.0	THERMAL PLUME MAPPING	
	George A. Nardacci	
5.1	Methods	167
5.2	Results	168
6.0	EFFECTS OF COOLING TOWER SALT DRIFT ON AGRICULTURAL CROPS AND NATURAL VEGETATION	
6.1	PLANT PATHOGENS	
	James D. Montgomery, Robert F. Eppley, Jr., Lynn D. Wike	
6.1.1	Methods	194
6.1.2	Results	195
6.1.3	Other Vegetation Disturbances	196
6.2	QUANTITATIVE VEGETATION STUDIES	
	James D. Montgomery, Lynn D. Wike	
6.2.1	Methods	207
6.2.2	Results	208
6.2.3	Discussion	212

POOR ORIGINAL

Table		Page
2.1-1	Location of trapnet and seine stations.	18
2.1-2	List of common and scientific names of fishes taken by trapnet and seine during 1974, 1975, and 1976 in the Susquehanna River in the vicinity of TMINS.	19
2.2-1	Fishes taken by trapnet on 3-5 March 1976 in the vicinity of TMINS.	20
2.2-2	Fishes taken by trapnet on 15-17 March 1976	20
2.2-3	Fishes taken by trapnet on 29-31 March 1976	21
2.2-4	Fishes taken by trapnet on 13-15 April 1976	21
2.2-5	Fishes taken by trapnet on 26-28 April 1976	22
2.2-6	Fishes taken by trapnet on 10-12 May 1976	22
2.2-7	Fishes taken by trapnet on 26-28 May 1976	23
2.2-8	Fishes taken by trapnet on 7-9 June 1976.	23
2.2-9	Fishes taken by trapnet on 29 June - 1 July 1976.	24
2.2-10	Fishes taken by trapnet on 6-8 July 1976.	25
2.2-11	Fishes taken by trapnet on 19-21 July 1976.	26
2.2-12	Fishes taken by trapnet on 2-4 August 1976.	27
2.2-13	Fishes taken by trapnet on 16-18 August 1976.	28
2.2-14	Fishes taken by trapnet on 1-3 September 1976	28
2.2-15	Fishes taken by trapnet on 13-15 September 1976	29
2.2-16	Fishes taken by trapnet on 27-29 September 1976	30
2.2-17	Fishes taken by trapnet on 18-20 October 1976	31
2.2-18	Fishes taken by trapnet on 25-27 October 1976	31
2.2-19	Summary of fishes taken at trapnet station TM-AQF-11A3 during 1976	32
2.2-20	Summary station TM-AQF-11A2.	32
2.2-21	Summary station TM-AQF-11A3.	33
2.2-22	Summary station TM-AQF-9B2	33
2.2-23	Summary of fishes taken at trapnet stations during March through October 1976	34
2.2-24	Monthly summary of fishes taken at trapnet stations upstream (U) and downstream (D) from the Discharge during March through October 1976	35
2.2-25	Indices of percent similarity of species composition between trapnet stations during 1974, 1975, and 1976.	36
2.2-26	Indices of percent similarity of species composition between seine stations during 1974, 1975, and 1976.	36
2.2-27	Fishes taken by seine on 2 March 1976 in the vicinity of TMINS.	37
2.2-28	Fishes taken by seine on 15 March 1976.	37
2.2-29	Fishes taken by seine on 29 March 1976.	38
2.2-30	Fishes taken by seine on 13 April 1976.	38

2.2-31	Fishes taken by seine on 30 April 1976.	39
2.2-32	Fishes taken by seine on 10 May 1976.	39
2.2-33	Fishes taken by seine on 24 May 1976.	39
2.2-34	Fishes taken by seine on 7 June 1976.	40
2.2-35	Fishes taken by seine on 21 June 1976.	40
2.2-36	Fishes taken by seine on 7 July 1976.	41
2.2-37	Fishes taken by seine on 19 July 1976.	41
2.2-38	Fishes taken by seine on 2 August 1976.	42
2.2-39	Fishes taken by seine on 16 August 1976.	42
2.2-40	Fishes taken by seine on 1 September 1976.	43
2.2-41	Fishes taken by seine on 13 September 1976.	43
2.2-42	Fishes taken by seine on 29 September 1976.	44
2.2-43	Fishes taken by seine on 19 October 1976.	44
2.2-44	Fishes taken by seine on 25 October 1976.	45
2.2-45	Summary of fishes taken at seine station TM-AQF-1A2 during 1976.	46
2.2-46	Summary . . . station TM-AQF-16A1.	46
2.2-47	Summary . . . station TM-AQF-10A2.	47
2.2-48	Summary . . . station TM-AQF-9A1.	47
2.2-49	Summary . . . station TM-AQF-9B3.	48
2.2-50	Summary of fishes taken at seine stations during March through October 1976.	48
2.2-51	Monthly summary of fishes taken at seine stations upstream (U) and downstream (D) from the Discharge during March through October 1976.	49
2.2-52	Condition factors and reproductive status of channel catfish taken by trapnet during March through October 1976 downstream (D) and upstream (U) from the TMINS Discharge.	50
2.2-53	Condition factors and reproductive status of rock bass.	51
2.2-54	Condition factors and reproductive status of pumpkinseed.	52
2.2-55	Condition factors and reproductive status of white crappie.	53
2.2-56	Condition factors and reproductive status of black crappie.	54
2.2-57	Condition factors and reproductive status of spottail shiner.	55
2.2-58	Condition factors and reproductive status of spotfin shiner.	56
2.2-59	Condition factors and reproductive status of white sucker.	56
2.2-60	Length frequency and mean weights per 5 m group of fishes taken by trapnet on 3-5 March 1976 downstream and upstream from the TMINS Discharge.	57
2.2-61	Length frequency . . . fishes taken by trapnet on 15-17 March 1976.	57
2.2-62	Length frequency . . . fishes taken by trapnet on 29-31 March 1976.	58
2.2-63	Length frequency . . . fishes taken by trapnet on 13-15 April 1976.	58
2.2-64	Length frequency . . . fishes taken by trapnet on 26-28 April 1976.	59
2.2-65	Length frequency . . . fishes taken by trapnet on 10-12 May 1976.	60

2.2-66	Length frequency fishes taken by trapnet on 26-28 May 1976	61
2.2-67	Length frequency fishes taken by trapnet on 7-9 June 1976.	61
2.2-68	Length frequency fishes taken by trapnet on 29 June - 1 July 1976.	62
2.2-69	Length frequency fishes taken by trapnet on 6-8 July 1976.	63
2.2-70	Length frequency fishes taken by trapnet on 19-21 July 1976.	64
2.2-71	Length frequency fishes taken by trapnet on 2-4 August 1976.	65
2.2-72	Length frequency fishes taken by trapnet on 16-18 August 1976.	66
2.2-73	Length frequency fishes taken by trapnet on 1-3 September 1976	66
2.2-74	Length frequency fishes taken by trapnet on 13-15 September 1976	67
2.2-75	Length frequency fishes taken by trapnet on 27-29 September 1976	68
2.2-76	Length frequency fishes taken by trapnet on 18-20 October 1976	69
2.2-77	Length frequency fishes taken by trapnet on 25-27 October 1976	69
2.2-78	Length frequency and mean weights per 5 mm group of fishes taken by seine on 2 March 1976 downstream and upstream from the THINS Discharge.	70
2.2-79	Length frequency fishes taken by seine on 15 March 1976.	70
2.2-80	Length frequency fishes taken by seine on 29 March 1976.	71
2.2-81	Length frequency fishes taken by seine on 13 April 1976.	71
2.2-82	Length frequency fishes taken by seine on 30 April 1976.	72
2.2-83	Length frequency fishes taken by seine on 10 May 1976.	72
2.2-84	Length frequency fishes taken by seine on 24 May 1976.	73
2.2-85	Length frequency fishes taken by seine on 7 June 1976.	73
2.2-86	Length frequency fishes taken by seine on 21 June 1976	74
2.2-87	Length frequency fishes taken by seine on 7 July 1976.	74
2.2-88	Length frequency fishes taken by seine on 19 July 1976	75
2.2-89	Length frequency fishes taken by seine on 2 August 1976.	76
2.2-90	Length frequency fishes taken by seine on 16 August 1976	77
2.2-91	Length frequency fishes taken by seine on 1 September 1976	78
2.2-92	Length frequency fishes taken by seine on 13 September 1976.	79
2.2-93	Length frequency fishes taken by seine on 29 September 1976.	80
2.2-94	Length frequency fishes taken by seine on 19 October 1976.	81
2.2-95	Length frequency fishes taken by seine on 25 October 1976.	82
2.3-1	Summary of number of specimens, number of species, number of collection, and number per collection of fishes taken by trapnet and seine upstream and downstream from the THINS Discharge during 1974 through 1976.	83
2.3-2	Kendall-tau correlation coefficients for species composition at trapnet and seine stations during 1974-1975, 1974-1976, and 1975-1976	84

FIGURE

Figure		
2.1-1	Location of trapnet and seine stations in the vicinity of THINS	85

SECTION 3.0 MACROINVERTEBRATES

TABLE

Table

3.1-1	Location and description of macroinvertebrate stations.	1-7
3.1-2	Macroinvertebrates taken from the Susquehanna River in the vicinity of Three Mile Island.	118
3.2-1	Numbers and milligrams of biomass, # eggs, of benthic organisms per Ponar grab (529 cm ²) at Station TM-AQI-1A1, April 1976	111
3.2-2	Numbers and milligrams of biomass at station TM-AQI-1A1, May 1976.	111
3.2-3	Numbers and milligrams of biomass at station TM-AQI-1A1, June 1976	111
3.2-4	Numbers and milligrams of biomass at station TM-AQI-1A1, July 1976	111
3.2-5	Numbers and milligrams of biomass at station TM-AQI-1A1, August 1976	111
3.2-6	Numbers and milligrams of biomass at station TM-AQI-1A1, September 1976.	111
3.2-7	Numbers and milligrams of biomass at station TM-AQI-1A1, October 1976.	111
3.2-8	Numbers and milligrams of biomass at station TM-AQI-1A2, April 1976.	111
3.2-9	Numbers and milligrams of biomass at station TM-AQI-1A2, May 1976.	111
3.2-10	Numbers and milligrams of biomass at station TM-AQI-1A2, June 1976	111
3.2-11	Numbers and milligrams of biomass at station TM-AQI-1A2, July 1976	111
3.2-12	Numbers and milligrams of biomass at station TM-AQI-1A2, August 1976	111
3.2-13	Numbers and milligrams of biomass at station TM-AQI-1A2, September 1976.	111
3.2-14	Numbers and milligrams of biomass at station TM-AQI-1A2, October 1976.	111
3.2-15	Numbers and milligrams of biomass at station TM-AQI-11A1, April 1976	111
3.2-16	Numbers and milligrams of biomass at station TM-AQI-11A1, May 1976	111
3.2-17	Numbers and milligrams of biomass at station TM-AQI-11A1, June 1976.	111
3.2-18	Numbers and milligrams of biomass at station TM-AQI-11A1, July 1976.	111
3.2-19	Numbers and milligrams of biomass at station TM-AQI-11A1, August 1976.	111
3.2-20	Numbers and milligrams of biomass at station TM-AQI-11A1, September 1976	111
3.2-21	Numbers and milligrams of biomass at station TM-AQI-11A1, October 1976	111
3.2-22	Numbers and milligrams of biomass at station TM-AQI-11A2, April 1976	111
3.2-23	Numbers and milligrams of biomass at station TM-AQI-11A2, May 1976	111
3.2-24	Numbers and milligrams of biomass at station TM-AQI-11A2, June 1976.	111
3.2-25	Numbers and milligrams of biomass at station TM-AQI-11A2, July 1976.	111
3.2-26	Numbers and milligrams of biomass at station TM-AQI-11A2, August 1976.	111
3.2-27	Numbers and milligrams of biomass at station TM-AQI-11A2, September 1976	111
3.2-28	Numbers and milligrams of biomass at station TM-AQI-11A2, October 1976	111
3.2-29	Numbers and milligrams of biomass at station TM-AQI-9B1, April 1976.	111
3.2-30	Numbers and milligrams of biomass at station TM-AQI-9B1, May 1976.	111
3.2-31	Numbers and milligrams of biomass at station TM-AQI-9B1, June 1976	111
3.2-32	Numbers and milligrams of biomass at station TM-AQI-9B1, July 1976	111

v

POOR ORIGINAL

1565 044

3.2-33	Numbers and milligrams of biomass at Station TM-AQ1-981, August 1976	143
3.2-34	Numbers and milligrams of biomass at Station TM-AQ1-981, September 1976.	144
3.2-35	Numbers and milligrams of biomass at Station TM-AQ1-981, October 1976.	145
3.2-36	Numbers and percent composition of benthic organisms collected at the macroinvertebrate stations, April through October 1976.	146
3.2-37	Estimates of density (number/m ²) of the dominant taxa at the macroinvertebrate stations, April through October 1976.	147
3.2-38	Estimates of biomass (mg/m ²) of the dominant taxa at the macroinvertebrate stations, April through October 1976.	148
3.2-39	Diversity (D) values at the macroinvertebrate stations, April through October 1976.	149
3.2-40	Indices of percent similarity of species composition (PNC) between the macroinvertebrate stations sampled April through October 1974, 1975, and 1976	149
3.2-41	Analysis of variance, randomized block design for number of taxa, April through October 1976.	149
3.2-42	Comparison of mean number of taxa at the macroinvertebrate stations using Student-Newman-Keuls multirange test, April through October 1976	149
3.2-43	Comparison of mean number of taxa by sampling date (April through October 1976) using Student-Newman-Keuls multirange test.	149
3.2-44	Three-factor analysis of variance for densities of <u>Limnodrilus hoffmeisteri</u> at the macroinvertebrate stations, May through October 1974 through 1976	150
3.2-45	Comparison of logarithmic mean density of <u>Limnodrilus hoffmeisteri</u> by year (May through October 1974-1976) using Student-Newman-Keuls multirange test	150
3.2-46	Comparison of logarithmic mean density of <u>Limnodrilus hoffmeisteri</u> by sampling date (May through October 1974-1976) using Student-Newman-Keuls multirange test	150
3.2-47	Comparison of logarithmic mean density of <u>Limnodrilus hoffmeisteri</u> at the macroinvertebrate stations using Student-Newman-Keuls multirange test, May through October 1974-1976.	150
3.3-1	Summary of data collected at the macroinvertebrate stations for 1974, 1975, and 1976.	151

FIGURES

Figure		
3.1-1	Location of benthic macroinvertebrate stations in the vicinity of TMINS	153
3.2-1	Percent composition of macroinvertebrate taxa, based on the total number of individuals collected at the macroinvertebrate stations, April through October 1976	153
3.3-1	Numbers of taxa taken at the macroinvertebrate stations, April through October 1976	154
3.3-2	Diversity (D) values at macroinvertebrate stations, April through October 1974, 1975, and 1976.	155

SECTION 4.0 AMBIENT WATER QUALITY

TABLES

Table		
4.1-1	Parameter, method used, and reference for analysis of water taken in the vicinity of TMINS in 1976	160
4.2-1	Monthly mean, minimum, and maximum concentrations of selected water quality parameters taken in the vicinity of TMINS in 1976.	161
4.2-2	Summary of selected physicochemical parameters taken on 6 and 20 April 1976 in the vicinity of TMINS	162
4.2-3	Summary on 4 and 18 May 1976	162

POOR ORIGINAL

1565 045

4.2-4	Summary on 1 and 15 June 1976.	163
4.2-5	Summary on 5 and 20 July 1976.	164
4.2-6	Summary on 3 and 17 August 1976.	164
4.2-7	Summary on 7 and 21 September 1976	164
4.2-8	Summary on 5 and 19 October 1976	165
4.2-9	Combinations of five consecutive samples for determination of the geometric mean of total coliform density at ambient water quality stations, April through October 1976.	166

SECTION 5.0 THERMAL PLUME MAPPING

TABLES

POOR ORIGINAL

Table

5.2-1	Thermal plume data for TMINS on 16 February 1976.	170
5.2-2	Thermal plume data for TMINS on 20 February 1976.	171
5.2-3	Thermal plume data for TMINS on 21 February 1976.	172
5.2-4	Thermal plume data for TMINS on 21 February 1976.	173
5.2-5	Thermal plume data for TMINS on 21 February 1976.	174
5.2-6	Thermal plume data for TMINS on 21 February 1976.	175
5.2-7	Thermal plume data for TMINS on 21 February 1976.	176
5.2-8	Thermal plume data for TMINS on 21 February 1976.	177
5.2-9	Thermal plume data for TMINS on 21 February 1976.	178
5.2-10	Thermal plume data for TMINS on 21 February 1976.	179
5.2-11	Thermal plume data for TMINS on 21 February 1976.	180
5.2-12	Thermal plume data for TMINS on 23 February 1976.	181
5.2-13	Thermal plume data for TMINS on 23 February 1976.	182
5.2-14	Thermal plume data for TMINS on 5 March 1976.	183
5.2-15	Thermal plume data for TMINS on 19 March 1976	183
5.2-16	Thermal plume data for TMINS on 1 April 1976.	184
5.2-17	Thermal plume data for TMINS on 15 April 1976	184
5.2-18	Thermal plume data for TMINS on 29 April 1976	185
5.2-19	Thermal plume data for TMINS on 13 May 1976	185
5.2-20	Thermal plume data for TMINS on 27 May 1976	186
5.2-21	Thermal plume data for TMINS on 2 June 1976	186
5.2-22	Thermal plume data for TMINS on 10 June 1976.	187
5.2-23	Thermal plume data for TMINS on 24 June 1976.	187
5.2-24	Thermal plume data for TMINS on 15 July 1976.	188
5.2-25	Thermal plume data for TMINS on 30 July 1976.	188
5.2-26	Thermal plume data for TMINS on 12 August 1976.	189
5.2-27	Thermal plume data for TMINS on 26 August 1976.	189

5.2-28	Thermal plume data for TMINS on 9 September 1976.	190
5.2-29	Thermal plume data for TMINS on 22 September 1976	190
5.2-30	Thermal plume data for TMINS on 7 October 1976.	191
5.2-31	Thermal plume data for TMINS on 21 October 1976	
5.2-32	Thermal plume data for TMINS on 4 November 1976	
5.2-33	Thermal plume data for TMINS on 18 November 1976.	192
5.2-34	Summary of surface water temperatures at selected plume map stations, discharge and ambient river temperature differences, air temperatures, wind conditions, discharge rate, river flow, and station operation level February through November 1976.	193

SECTION 6.0 EFFECTS OF COOLING TOWER SALT DRIFT ON AGRICULTURAL CROPS AND NATURAL VEGETATION

6.1 PLANT PATHOGENS

TABLES

Table

6.1-1	Plants observed in flower in salt drift transects in the vicinity of TMINS in 1976. . . .	197
6.1-2	Parasitic plant diseases observed on agricultural crops in plant pathology transects, April through October 1976.	203
6.1-3	Parasitic plant diseases observed on natural vegetation in plant pathology transects, April through October 1976.	204
6.1-4	Vegetation exhibiting significant insect damage in plant pathology transects, April through October 1976.	205

FIGURE

Figure

6.1-1	Location of salt drift transects and areas of quantitative vegetation study, in the vicinity of TMINS	206
-------	---	-----

SECTION 6.2 QUANTITATIVE VEGETATION STUDIES

TABLES

Table

6.2-1	Comparison of trees and saplings on 24 selected plots in the bottomland hardwood forest on Shelley Island, 1973-1976.	214
6.2-2	Comparison of shrubs and vines on 24 selected plots in the bottomland hardwood forest on Shelley Island, 1973-1976	215
6.2-3	Ground cover occurring in the bottomland hardwood forest on Shelley Island, 1976. . . .	216
6.2-4	Comparison of dominant ground cover occurring in the bottomland hardwood forest on Shelley Island, 1973-1976	217
6.2-5	Tree seedlings in bottomland hardwood forest on Shelley Island, 1974, 1975, and 1976. . .	217
6.2-6	Comparison of trees and saplings on 17 selected plots in the south forest on Three Mile Island, 1973-1976	218
6.2-7	Comparison of shrubs and vines on 17 selected plots in the south forest on Three Mile Island, 1973-1976	218
6.2-8	Ground cover occurring in the "vine" portion of the black locust forest on Three Mile Island, 1976.	219
6.2-9	Comparison of the dominant ground cover occurring in the "vine" portion of the south woodlot on Three Mile Island, 1973-1976	220

6.2-10	Tree seedlings in the "vine" portion of the black locust forest on Three Mile Island, 1973-1976	220
6.2-11	Ground cover occurring in the "non-vine" portion of the black locust forest on Three Mile Island, 1976.	221
6.2-12	Comparison of the dominant ground cover occurring in the "non-vine" portion of the south woodlot on Three Mile Island, 1973-1976	222
6.2-13	Tree seedlings in the "non-vine" portion of the black locust forest on Three Mile Island, 1973, 1975, and 1976.	222
6.2-14	Ground cover occurring in the south field on Three Mile Island, 1976.	223
6.2-15	Comparison of dominant ground cover occurring in the south field on Three Mile Island, 1973-1976	224
6.2-16	Tree seedlings in the south field on Three Mile Island, 1973-1976	224
6.2-17	Ground cover occurring in the field near the south ponds on Three Mile Island, 1976	225
6.2-18	Comparison of dominant ground cover occurring in the field near the south ponds on Three Mile Island, 1973-1976.	226
6.2-19	Tree seedlings in the field near the south ponds on Three Mile Island, 1976	226
6.2-20	Ground cover occurring in the north field on Three Mile Island, 1976.	227
6.2-21	Comparison of the dominant ground cover occurring in the north field on Three Mile Island, 1973-1976	229
6.2-22	Tree seedlings in the north field on Three Mile Island, 1973-1976	229
6.2-23	Ground cover occurring in the field near the 500 kv substation, 1976.	230
6.2-24	Comparison of the dominant ground cover occurring in the field near the 500 kv substation on Route 441, 1974-1976	231
6.2-25	Tree seedlings in the field near the 500 kv substation, 1974-1976	231

POOR ORIGINAL

1565 048

POOR ORIGINAL

ACKNOWLEDGMENTS

We wish to thank Barbara Beck, Michael Buring, Thomas Jenckes, Bud Koch, James Mudge, John Romanski, and James Stacey of Metropolitan Edison Company for their assistance in the past year. We also appreciate the help of Three Mile Island Nuclear Station Security without whose assistance our work would have been impossible.

The following biologists participated in the study in 1976:
John Epler, Robert Eppley, Richard Evans, Harold Hagerty, Barbara Lathrop, Robert Malick, Eric McClellan, James Montgomery, George Nardacci, Janet Polk, Wayne Potter, Philip Ritson, James Smith, Walter Soya, Linda Wike, and Lynn Wike.

Linda Wike provided secretarial services and assistance.

We would also like to acknowledge the following specialists for their advice and assistance in confirming the performing determinations:
Dr. Marvin L. Bobb (Hemiptera), Mr. Jarl K. Hiltunen (Tubificidae), Mr. Michael S. Loden (Naididae), Mr. Thomas P. Poe (Hirudinea), Dr. John W. Reynolds (Megadriles), Dr. Selwyn S. Roback (Chironomidae), Dr. Ole A. Saether (Chironomidae), Mr. Clark N. Shiffer (Odonata). Others who assisted in providing help include: Dr. Robert Denoncourt, the Pennsylvania Fish Commission and a special thanks to Dr. Guy L. Steucke (Millersville State College) and Jeffery J. Hondo (Ichthyological Associates, Inc., Absecon, N.J.) for their statistical advice.

1565 049

1.0 INTRODUCTION

This document is the third annual postoperational report on the ecology of York Haven Pond (Lake Frederic), Susquehanna River, Pennsylvania in the vicinity of the Three Mile Island Nuclear Station (TMINS). Ichthyological Associates, Inc. initiated the study in February 1974. This report covers the period from January through December 1976. Sections are presented to meet the Environmental Technical Specifications (ETS) for TMINS, Unit 1, Appendix B. Parameters analyzed are the same as reported in Potter and Associates (1976) with the exception of impingement of fishes, entrainment of ichthyoplankton, entrainment of plankton (phytoplankton and zooplankton), and bird impaction on cooling towers. The ETS requirements for surveillance of these parameters were fulfilled and the programs were terminated.

The TMINS Unit 1 achieved criticality on 5 June 1974. Unit 1 has been at various levels of operation ranging from complete shutdown to 100%, 830 megawatts. In 1976, the TMINS Unit 1 was shutdown for refueling from 21 February to 27 May.

Potter, W.A. and Associates. 1976. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Annual Report for 1975. Ichthyological Associates, Inc. 395 pp.

POOR ORIGINAL

1565 050

POOR ORIGINAL

1.1 SUMMARY

1.1.1 Fish

Fishes were sampled at four trapnet and five seine stations, March through October 1976.

A total of 144 trapnet collections yielded 822 fish of 25 species. The pumpkinseed and channel catfish were most abundant. Common fishes were taken in equal or greater numbers downstream from the Discharge than upstream. No significant change in species rank at individual stations was observed from 1974 through 1976.

A total of 90 seine collections yielded 10,478 fish of 35 species. The spottail shiner ranked first in overall abundance and was the most common species at all stations. Most common species showed nearly equal distribution among upstream and downstream stations. Significant differences in species rank at one upstream and one downstream station were observed between 1975 and 1976.

No appreciable differences were noted in condition factor or weights downstream and upstream from the Discharge where sufficient numbers were present; most values were within the ranges observed in previous years.

The fish leech, Myxobdella lugubris, and anchor worm (Lernaea spp.) were common ectoparasites.

Variations in fish abundance may be attributable to year class fluctuations. The impact of TMINS Unit 1 on fishes was negligible.

1565 051

1.1.2 Macroinvertebrates

Macroinvertebrate collections were taken semimonthly April through October 1976. A total of 90,567 specimens representing 112 taxa was taken. Limnodrilus hoffmeisteri, Chironomus tentans, and Nais elinguis were the most abundant taxa. Limnodrilus hoffmeisteri was usually most abundant at all stations each month. Coniobasis virginica, L. hoffmeisteri, and C. decorus had the greatest biomass of all specimens weighed.

Monthly estimates of diversity ranged from 0.66 to 2.76. Index of percent similarity values between stations were in the intermediate to high affinity range (>50% similarity). High similarities in species composition were found between Station 1A2 and 9B1 (91%), and 11A1 and 11A2 (89%).

Comparison of number of taxa between macroinvertebrate stations and sample dates for 1976 was accomplished by analysis of variance, randomized block design. The Student-Newman-Keuls multirange test revealed that Stations 1A2 and 9B1 and 11A1, 11A2, and 9B1 were similar in number of taxa.

Three-factor analysis of variance performed on densities of Limnodrilus hoffmeisteri May through October 1974 through 1976 revealed that years, sample dates, and stations were significantly different.

The macroinvertebrate communities appeared to be more affected by ice and high river flow than by the operation of TAMS.

1565 052

POOR ORIGINAL

1.1.3 Ambient Water Quality

1.1.3.1 Physicochemical Parameters

Except for dissolved oxygen and total suspended solids the monthly distribution of parameters was similar. Mean values were high in September and low in April and May. The monthly mean concentrations of most parameters at Station 11A1 (TMINS Discharge) were generally higher than at the other stations.

1.1.3.2 Fecal Coliform

The highest overall mean density of fecal coliform was at Station 1A1; the lowest was at Station 11A1 (TMINS Discharge). Monthly mean fecal coliform densities were lowest in April, increased throughout the summer, and peaked in September. For all stations the geometric mean of fecal coliform densities exceeded the limit established for the Commonwealth of Pennsylvania.

1.1.4 Thermal Plume Mapping

Thirty-three plume surveys were conducted at various river flow conditions and station operation levels. The discharge temperatures were within the limits established in the ETS. A return of the discharge temperature to within 2.7 C of ambient was used to define the plume. In 28 of 33 surveys the plume was limited to 5 m offshore and 25 m downstream from the Discharge.

1565 053

the analytical plume model for normal cooldown conditions was compared with the cooldown for refueling on 21 February. The plume was recorded no further than 20 m offshore and 1000 m downstream from the Discharge. This varies from the predicted model presented in the environmental statement for the operation of TMINS. High river flow during the cooldown operations may account for this variance.

1.1.5 Effects of Cooling Tower Salt Drift on Agricultural Crops and Natural Vegetation

1.1.5.1 Plant Pathogens

Plant pathology transects were examined from April through October 1976. No differences were noted in flowering time or appearance of the 219 taxa observed with respect to the location of possible salt drift. Plant parasitic diseases were found on four agricultural crops and 21 taxa of natural vegetation, and insect damage was noted on nine taxa. None of the damage caused significant defoliation and no pattern was observed relative to the operation of the cooling towers at TMINS.

1.1.5.2 Quantitative Vegetation Studies

Two forests and four fields were surveyed late August through mid-October 1976; results were statistically compared with those obtained in 1973 and 1975. There were few changes in the overstory and understory in the forests. Some of the statistically significant changes in ground cover in forests and fields were related to natural or human disturbance; others were normal in the course of secondary succession. No pattern of change was found that was attributed to the operation of the cooling towers at TMINS.

2.0 FISH

The ETS, Appendix B, Section 4.1.1D requires replicate fish samples be taken both inside and outside of the thermal plume every two weeks, March through October.

POOR ORIGINAL

2.1 METHODS

Fish were sampled every two weeks at four trapnet and five seine stations, March⁷ through October 1976 (Table 2.1-1 and Figure 2.1-1). High river flows during June and October necessitated collections be taken on succeeding weeks. Trapnet collections taken on 29 June through 1 July were analyzed as June data.

Habitat differences in water velocity, depth, substrate, and available cover were observed at the stations. Riffles were noted at the upstream seine stations during low river flow; no riffles were encountered downstream. Mud and silt were common substrates downstream while mud, rubble, and boulders predominated upstream. Rubble, boulders, and vegetation provided limited cover. Water depths ranged to 2 m.

Fishes taken by trapnet and seine since 1974 are listed in Table 2.1-2. Common and scientific names and taxonomic order of presentation followed Bailey et al. (1970).

References used for fish identification included Denoncourt (1975), Gibbs (1957), Hubbs and Lagler (1964), Pflieger (1975), Scott and Crossman (1973), Snelson (1968), and Trautman (1957).

1565 055

Identification of young of the white sucker from those of the shorthead redhorse was difficult. Young were distinguishable only upon internal examination of gas bladder chambers. Therefore, specimens were listed as white sucker/shorthead redhorse until they were large enough to be identified by external characteristics.

Condition factor (K) for fishes comprising more than 10% of the trapnet or seine catch during 1974, 1975, or 1976 was calculated using the formula:

$$K = 100W/L^3$$

where W = mean weight (g) per 5 mm group, and L = upper limit of 5 mm fork length interval expressed in cm.

Species diversity indices (D) were calculated for each trapnet and seine station using the Shannon-Weaver function presented by Lloyd et al. (1968):

$$D = C/N(N \log_{10} N - \sum n_i \log_{10} n_i)$$

where C = 3.321928 (converts base 10 log to base 2), N = total number of individuals, and n_i = total number of individuals in the i th species.

An index was computed to identify the percent similarity between stations with respect to composition of fishes (Whittaker and Fairbanks 1958). It was expressed as:

$$PSc = \sum \min(a,b)$$

where PSc = the percent similarity and a and b = the percentages of a species in samples A and B. PSc values range from 0.0 (no similarity) to 100.0 (complete similarity). This index measures relative similarity in terms of species populations and often leads to grouping of communities by dominants or major species.

POOR ORIGINAL

1565 056

Kendall's coefficient of rank correlation (Sokal and Rohlf 1973) was applied to the catch data to analyze the yearly variation in rankings of species at a station. This did not test whether numbers collected in different years were significantly different, only whether their respective rankings were correlated (Summerfelt and Minckley 1969).

Physicochemical data included time, weather, secchi disc, and air and water temperatures. Dissolved oxygen concentration and pH were determined from water samples taken at each station. River stage was obtained from the Harrisburg River Gauge Station for 0700 hr. Data for trapnet catches were presented as two, separate 24-hr collections. Minimum values for parameters were listed first regardless of collection date. Single recordings for a parameter indicate identical observations throughout the period.

Each trapnet consisted of a 0.91 m by 15.24 m lead net and a 0.91 m by 1.83 m metal frame connected to two traps (4, 0.76 m diameter hoops). The lead net and trapnet were of 1.27 cm mesh. Nets were set for 24-hr, were checked, and reset for a second 24-hr period. Effort was made to set nets perpendicular to shore; however, high flows often caused nets to be set at angles of 45 degrees or less to shore to prevent rolling. Fishes were identified, measured, weighed, and released in the field.

A 3.05 m by 1.22 m seine with 0.32 cm mesh was used; at least three hauls were made for each collection. Additional hauls were taken when further effort might alter the relative abundance of fishes or yield other species. Fishes were preserved in 10% formalin for one week, rinsed in water and let stand for two days, and stored in 40% isopropanol.

POOR ORIGINAL

1565 057

Specimens were measured to within a 5 mm fork length interval. All fish of a species within the same interval in each collection were weighed to the nearest 0.1 g. A subsample was measured and weighed when a large number (100+) of one species was taken in a collection.

Reproductive status for fishes was defined as follows: young were spawned during the current calendar year; juveniles were incapable of reproduction, or minnows and darters less than 26 mm collected prior to the current spawning season; and adults were capable of reproduction. Classifications were based on field observations and information in the literature (Carlander 1953, 1969; Miller and Buss 1963; Scott and Crossman 1973; Trautman 1957).

Fishes were examined for ectoparasites.

2.2 RESULTS

2.2.1 TRAPNET

Results of March through October trapnet collections are reported in Tables 2.2-1 through 2.2-13 and are summarized in Tables 2.2-19 through 2.2-24. A total of 144 trapnet collections (36 upstream from the Discharge and 108 downstream) yielded 822 fish of 25 species (Tables 2.2-23 and 2.2-24). Most fish (295) and most species (20) were taken at Stations 1A3 and 11A2, respectively. The number of specimens per collection (n/Coll.) was 8.19 upstream and 4.88 downstream (Table 2.2-24). Except for the white sucker and northern hog sucker, all species were taken in equal or greater numbers downstream. Catch per month was greatest at three of the four stations (1A3, 11A2, 9B2) in September (Tables 2.2-19 through 2.2-22).

POOR ORIGINAL

1565 058

Sunfishes and catfishes dominated the catch. The five most abundant fishes (74.5% of the total catch) were the pumpkinseed, channel catfish, rock bass, black crappie, and redbreast sunfish (Table 2.2-23).

Nine sunfishes comprised 62.5% of the trapnet catch. The pumpkinseed ranked first in overall abundance (26.0%) and was the most common species at Stations 1A3 and 9B2 (Table 2.2-23). Other sunfishes that comprised more than 5.0% of the catch were the rock bass (11.7%), black crappie (10.1%), and redbreast sunfish (8.6%).

Five catfishes accounted for 23.1% of the catch; slightly greater numbers were taken downstream (Table 2.2-24). The channel catfish (18.1%) and brown bullhead (5.0%) were most common.

Two juvenile alewife were captured at Station 922 on 13-14 September (Table 2.2-15). This species had not been recorded from the lower Susquehanna drainage in Pennsylvania since the 1930's (Fowler 1940). Introductions were made in 1969 into impoundments in the Susquehanna watershed in Centre, Columbia, and Luzerne counties (Robert B. Hesser, Pennsylvania Fish Commission, personal communication). The specimens may have migrated downstream from one of these impoundments.

Other fishes taken for the first time by trapnet included the muskellunge, northern hog sucker, yellow perch, and walleye. Four species previously taken by trapnet, the goldfish, bluntnose minnow, fallfish, and tessellated darter were not collected in 1976.

POOR ORIGINAL

1565 059

POOR ORIGINAL

Diversity indices calculated from summary data for each station ranged from 2.95 (1A3) to 3.35 (11A2) Table 2.2-23. Monthly indices at most stations peaked in July or August and ranged from 0.00 at 11A3 in October to 3.24 at 11A2 in August (Tables 2.2-19 through 2.2-22). Lower diversity indices reflected the dominance of a single species.

Percent similarity indices between stations were moderate (Table 2.2-25). The greatest similarity (71.2) was between Stations 1A3 and 11A2; it was caused by the dominance of the channel catfish and rock bass. The similarity between Stations 1A3 and 9B2 (66.1) was characterized by like habitats and dominance by the pumpkinseed. The least similarity of 49.4, between Stations 11A3 and 9B2, was caused by dissimilar habitats and the absence of common dominant species.

2.2.2 SEINE

Results of March through October seine collections for each date are presented in Tables 2.2-27 through 2.2-44 and are summarized in Tables 2.2-45 through 2.2-51. A total of 90 collections (36 upstream from the Discharge and 54 downstream) yielded 10,478 fish of 35 species (Tables 2.2-50 and 2.2-51). Most fish (2,645) and most species (29) were taken at Stations 1A2 and 16A1, respectively. The numbers of specimens per collection upstream and downstream were comparable (136.50 and 103.04, respectively). Most dominant species showed nearly equal distribution among upstream and downstream stations.

1565 060

Seine catches were small and erratic from March through May; only 48 specimens were collected in May (Table 2.2-51). The catch reached a peak in June (3,611 fish) and corresponded with a large spawn of the spottail shiner. Catches at all stations except 1A2 decreased during July and August. A secondary peak of 2,050 fish was recorded in September when young of the spotfin shiner were abundant. The number of specimens per collection was similar at all stations except 9A1 (Table 2.2-50). Fewest fish (1,006) and least species (19) were taken at Station 9A1. Seines were difficult to fish at Station 9A1 because of boulders and deep mud.

The spottail shiner comprised 52.7% of the catch and was the most abundant species at all stations (Table 2.2-50). The spotfin shiner ranked second in overall abundance (19.8%); nearly half of the specimens were taken at Station 1A2. Young of the white sucker/shorthead redhorse accounted for 6.7% of the catch and were most common at Station 16A1. The tessellated darter ranked fourth in overall abundance (6.5%) and was most common in September. Young of the channel catfish ranked fifth in overall abundance (4.7%); more than 99% were taken downstream from the Discharge (Table 2.2-51).

Fishes taken for the first time from the study area by seine included the fathead minnow, yellow bullhead, and shield darter. The yellow perch was the only species taken in previous years not collected in 1976.

Overall station diversity indices were low due to dominance by the spottail shiner and spotfin shiner and ranged from 1.75 (1A2) to 2.45 (10A2) Table 2.2-50. The highest monthly index (2.71) occurred at Station 1A2 in August (Table 2.2-45).

POOR ORIGINAL

1565 061

Percent similarity indices between stations were high (Table 2.2-26). The greatest similarity (82.0) occurred between downstream Stations 10A2 and 9A1. The lowest similarity (64.4) was found between the two upstream stations, 1A2 and 16A1, and resulted from differences in abundance of the spotfin shiner and suckers. Similarities between upstream and downstream stations were reduced by the unequal distribution of the channel catfish. Similarities seemed less dependent on the presence of like habitats and more dependent on affinities of dominant species.

2.2.3 REPRODUCTIVE STATUS AND CONDITION FACTOR

Reproductive status and condition factor (K) per 5 mm fork length interval for the channel catfish, rock bass, pumpkinseed, white crappie, and black crappie taken by trapnet and the spottail shiner, spotfin shiner, and white sucker taken by seine are presented in Tables 2.2-52 through 2.2-59. Mean weights per 5 mm fork length intervals of fishes taken by trapnet and seine are given in Tables 2.2-60 through 2.2-95.

Juvenile channel catfish were common in trapnet catches during March, April, September, and October; most adults were taken in October. Young channel catfish first appeared in seine catches on 7 July; this may indicate that spawning began in June. Most rock bass taken by trapnet were adults. Ripe male and female pumpkinseed were common in June and July trapnet catches. Young of the white crappie and black crappie were taken by trapnet in September.

POOR ORIGINAL

1565 062

Most spottail shiners taken by seine were young or juvenile; young were first collected in June. Spottails spawned in May and June grew to 50-60 mm FL by October. The spotfin shiner appeared to have an extended spawning season with peaks in June and August. Winter survival of young spotfins of the 1975 year class was good as indicated by the abundance of juveniles in spring 1976 seine catches. Most white suckers spawned in June; young were common in July seine catches.

No appreciable differences were noted in condition factors or weights downstream and upstream from the Discharge where sufficient numbers were present; most values were within the ranges observed in previous years.

2.2.4 PARASITES

The fish leech, Myzobdella lugubris, was a common parasite of the tessellated darter. Infestation on darters was first observed in August and reached a peak in October. Some darters were parasitized by up to five leeches. The site of attachment was always on or near a fin. No parasitized darters smaller than 31 mm were observed. One adult channel catfish was also parasitized. M. lugubris shows the greatest distribution of any piscicolid reported in North America (Hugghins 1972) and is probably the most abundant fish leech of catfishes in the Midwest (Klemm 1972).

Anchor worms (Lernaea spp.) were found on a few specimens of the comely shiner, spottail shiner, spotfin shiner, bluntnose minnow, bluegill, and smallmouth bass. Most of these infected fishes were in poor condition.

A slight infection of black spot, caused by metacercariae of digenetic trematodes, was observed on one juvenile creek chub.

1565 063

POOR ORIGINAL

2.3 DISCUSSION

Trapnet catches have declined since the inception of the program in 1974 (Table 2.3-1). The smallest yearly catch (822) occurred in 1976 and was roughly half of the 1975 catch (1,610). The decline in the 1976 catch resulted from decreases in the number of fish taken at the downstream stations. The upstream catches in 1975 and 1976 were comparable (324 and 295, respectively). The number of specimens per upstream collection increased from 6.35 in 1975 to 8.19 in 1976. The total number of specimens captured in 1974 and 1975 was inflated by large collections of juvenile channel catfish downstream from the Discharge. No large collections of channel catfish occurred in 1976. Variation in the relative abundance of some species may have resulted from fluctuations in year class strength. Results of Kendall's coefficient of rank correlation tests showed no significant changes ($P \geq .05$) in species ranks at individual stations during 1974 through 1976 (Table 2.3-2).

Fluctuations in yearly seine catches resulted from large differences in the numbers of specimens taken upstream from the Discharge (Table 2.3-1). Catch data from downstream stations were similar throughout the study period. Catches were most influenced by the availability of young of the spottail shiner, spotfin shiner, and white sucker/shorthead redhorse. Results of Kendall's coefficient of rank correlation tests showed significant differences ($P \geq .05$) in species ranks at Stations 1A2 (upstream from the Discharge) and 9A1 (downstream from the Discharge) in 1975 and 1976 (Table 2.3-2).

1565 064

It is concluded that the impact from the operation of Unit 1 on the populations of fishes vulnerable to trapnet and seine was negligible. The ETS requirements for Unit 1 are fulfilled and programs will continue as specified in the ETS for Unit 2.

Bailey, B.M., J.E. Fitch, E.S. Herald, E.A. Lachner, C.C. Lindsey, C.R. Robins, and W.B. Scott. 1970. A list of common and scientific names of fishes from the United States and Canada. Amer. Fish. Soc. Special Publ. No. 6. 150 pp.

Carlander, K.D. 1953. Handbook of freshwater fishery biology with the first supplement. Wm. C. Brown Co., Dubuque, Iowa. 430 pp.

_____. 1969. Handbook of freshwater fishery biology. Vol. 1. Life history data on freshwater fishes of the United States and Canada, exclusive of the Perciformes. Iowa State Univ. Press. Ames, Iowa. 752 pp.

Denoncourt, R.F. 1975. Key to the families and genera of Pennsylvania freshwater fishes and the species of freshwater fishes of the Susquehanna River drainage above Conowingo Dam. Proc. Pa. Acad. Sci. 49:82-88.

Fowler, H.W. 1940. A list of the fishes recorded from Pennsylvania. Comm. of Pa., Bd. Fish Comm. Bull. 7. 25 pp.

Gibbs, R.H., Jr. 1957. Cyprinid fishes of the subgenus Cyprinella of Notropis. II. Distribution and variation of Notropis spilopterus, with the description of a new subspecies. Lloydia 20(3):186-211.

Hubbs, C.L. and K.F. Lagler. 1964. Fishes of the Great Lakes region. Univ. Mich. Press, Ann Arbor, Mich. 213 pp.

Hughins, E.J. 1972. Parasites of fishes in South Dakota. South Dakota Dept. Game, Fish, and Parks. Bull. 484. 73 pp.

Klemm, D.J. 1972. The leeches (Annelida:Hirudinea) of Michigan. Mich. Academician IV(4):405-444.

Lloyd, M., J.H. Zar, and J.R. Karr. 1968. On the calculation of information-theoretical measures of diversity. Amer. Midl. Nat. 79(2): 257-272.

POOR ORIGINAL

1565 065

- Miller, J. and K. Buss. [1963?] . The age and growth of the fishes in Pennsylvania. Pa. Fish Comm. 26 pp.
- Pfleiger, W.L. 1975. The fishes of Missouri. Missouri Dept. of Cons. 343 pp.
- Scott, W.B. and E.J. Crossman. 1973. Freshwater fishes of Canada. Fish. Res. Bd. Can. Bull. 184. 966 pp.
- Snelson, F.F., Jr. 1968. Systematics of the Cyprinid fish Notropis argemus, with comments on the subgenus Notropis. Copeia 1968 (4): 776-802.
- Sokal, R.R. and F.J. Rohlf. 1973. Introduction to biostatistics. W.H. Freeman and Co. San Francisco, Ca. 368 pp.
- Summerfelt, R.C. and C.O. Minckley. 1969. Aspects of the life history of the sand shiner, Notropis stramineus (Cope), in the Smoky Hill River, Kansas. Trans. Am. Fish. Soc. 98(3):444-453.
- Trautman, M.B. 1957. The fishes of Ohio with illustrated keys. Ohio State Univ. Press, Columbus, Ohio. 683 pp.
- Whittaker, R.H. and C.W. Fairbanks. 1958. A study of plankton copepod communities in the Columbia Basin, Southeastern Washington. Ecology 39:46-65.

POOR ORIGINAL

1565 066

POOR ORIGINAL

Table 2.1-1

Location of trapnet and seine stations.

Station Number	Location
TRAPNET	
	UPSTREAM
TM-AQF-1A3*	Off southwest shore of St. Johns Island.
	DOWNSTREAM
TM-AQF-11A2	TMINS Discharge.
TM-AQF-11A3	200 m downstream from TMINS Discharge.
TM-AQF-9B2	1900 m downstream from TMINS Discharge.
SEINE	
	UPSTREAM
TM-AQF-1A2	Northwest St. Johns Island.
TM-AQF-16A1	25 m upstream from TMI boat dock.
	DOWNSTREAM
TM-AQF-10A2	150 m downstream from TMINS Discharge.
TM-AQF-9A1	1500 m downstream from TMINS Discharge.
TM-AQF-9B3	2000 m downstream from TMINS Discharge.

* Polar coordinate prefix TM-AQF- deleted from station numbers for discussion in text.

1565 067

POOR ORIGINAL

Table 2.1-2

List of common and scientific names of fishes taken by trapnet and seine during 1974, 1975, and 1976 in the Susquehanna River in the vicinity of EMCS.

Common Name	Scientific Name	Trapnet			Seine		
		1974	1975	1976	1974	1975	1976
Alewife	<u>Alosa pseudoharengus</u> (Wilson)	-	-	x	-	-	-
Muskellunge	<u>Esox masquinongy</u> Mitchill	-	-	x	-	-	-
Goldfish	<u>Carassius auratus</u> (Linnaeus)	x	-	-	-	-	-
Carp	<u>Cyprinus carpio</u> Linnaeus	x	x	x	-	x	x
Cutlips minnow	<u>Foxallium maxillaria</u> (Lesueur)	-	-	-	x	-	x
River chub	<u>Notropis macrodonum</u> (Cope)	-	-	-	x	x	x
Golden shiner	<u>Notemigonus crysoleucas</u> (Mitchill)	x	x	x	x	x	x
Comely shiner	<u>Notropis anogenus</u> (Abbott)	-	-	-	x	x	x
Common shiner	<u>Notropis cornutus</u> (Mitchill)	-	-	-	x	x	x
Spottail shiner	<u>Notropis hudsonius</u> (Clinton)	x	x	x	x	x	x
Swallowtail shiner	<u>Notropis procerus</u> (Cope)	-	-	-	x	x	x
Rosyface shiner	<u>Notropis rubellus</u> (Agassiz)	-	-	-	-	x	x
Spotfin shiner	<u>Notropis spilopterus</u> (Cope)	x	x	x	x	x	x
Bluntnose minnow	<u>Pimephales notatus</u> (Rafinesque)	-	x	-	x	x	x
Fathead minnow	<u>Pimephales promelas</u> Rafinesque	-	-	-	-	-	x
Blacknose dace	<u>Shimichthys atratulus</u> (Hermaun)	-	-	-	x	x	x
Longnose dace	<u>Shimichthys catarractae</u> (Valenciennes)	-	-	-	x	x	x
Creek chub	<u>Semotilus atromaculatus</u> (Mitchill)	-	-	-	x	x	x
Fallfish	<u>Semotilus corporalis</u> (Mitchill)	x	-	-	x	x	x
Quillback	<u>Carpiodes cyprinus</u> (Lesueur)	x	x	x	x	x	x
White sucker	<u>Catostomus commersoni</u> (Lacepede)	x	x	x	x	x	x
Northern hog sucker	<u>Hypoclinemus nigricans</u> (Lesueur)	-	-	x	x	x	x
Shorthead redhorse	<u>Notropis macrolepidotum</u> (Lesueur)	x	x	x	x	x	x
White catfish	<u>Ictalurus uatus</u> (Linnaeus)	x	x	x	x	-	x
Yellow bullhead	<u>Ictalurus natalis</u> (Lesueur)	x	x	x	-	-	x
Brown bullhead	<u>Ictalurus nebulosus</u> (Lesueur)	x	x	x	x	-	x
Channel catfish	<u>Ictalurus punctatus</u> (Rafinesque)	x	x	x	x	x	x
Margined madtom	<u>Noturus insignis</u> (Richardson)	x	x	x	-	-	-
Rock bass	<u>Ambloplites rupestris</u> (Rafinesque)	x	x	x	x	x	x
Redbreast sunfish	<u>Lepomis gibbosus</u> (Linnaeus)	x	x	x	x	x	x
Pumpkinseed	<u>Lepomis gibbosus</u> (Linnaeus)	x	x	x	x	x	x
Bluegill	<u>Lepomis macrochirus</u> Rafinesque	x	x	x	x	x	x
Smallmouth bass	<u>Micropterus dolomieu</u> Lacpede	x	x	x	x	x	x
Largemouth bass	<u>Micropterus salmoides</u> (Lacepede)	x	x	x	x	x	x
White crappie	<u>Pomoxis annularis</u> Rafinesque	x	x	x	x	x	x
Black crappie	<u>Pomoxis nigromaculatus</u> (Lesueur)	x	x	x	x	x	x
Tessellated darter	<u>Etheostoma olmstedti</u> Storer	-	x	-	x	x	x
Sanded darter	<u>Etheostoma caeruleum</u> (Cope)	-	-	-	x	x	x
Yellow perch	<u>Perca flavescens</u> (Mitchill)	-	-	x	-	x	-
Shield darter	<u>Percina nebulosa</u> (Stauffer)	-	-	-	-	-	x
Walleye	<u>Stizostedion vitreum vitreum</u> (Mitchill)	-	-	x	-	-	-

1565 068

Table 2.2-1
Fishes taken by trapnet on 3-5 March 1976 in the vicinity of THNS.

Station	13 MAR-1976					14 MAR-1976					15 MAR-1976				
	3-4	4-5	5-6	6-7	7-8	3-4	4-5	5-6	6-7	7-8	3-4	4-5	5-6	6-7	7-8
Air Temp. (C)	3.0, 7.0	7.0, 9.0	6.0, 9.0	6.0, 9.0	6.0, 9.0	3.0, 6.0	6.0, 11.5	6.0, 11.5	6.0, 11.5	6.0, 11.5	3.0, 6.0	6.0, 11.5	6.0, 11.5	6.0, 11.5	6.0, 11.5
Water Temp. (C)	5.5	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.0, 5.5	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.0, 5.5	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.5, 6.0
Dissolved Oxygen (ppm)	9.4	9.4, 9.9	9.4, 9.9	9.4, 9.9	9.4, 9.9	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2
pH	7.9, 8.2	8.2, 8.3	8.2, 8.3	8.2, 8.3	8.2, 8.3	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4
Secchi disc (cm)	51, 56	82, 83	82, 83	82, 83	82, 83	33, 53	53	53	53	53	66, 66	51, 66	51, 66	51, 66	51, 66
Water stage (ft)	6.55, 6.62	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.55, 6.62	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.55, 6.62	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.67, 6.70
Weather	Light rain,					Light rain,					Light rain,				
No. of fish	12	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Species	12	1	1	1	1	1	1	1	1	1	1	1	1	1	1
White sucker	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Shortnose sturgeon	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Channel catfish	6	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Parakeet fish	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Bluegill	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

Table 2.2-2
Fishes taken by trapnet on 13-17 March 1976 in the vicinity of THNS.

Station	13 MAR-1976					14 MAR-1976					15 MAR-1976				
	3-4	4-5	5-6	6-7	7-8	3-4	4-5	5-6	6-7	7-8	3-4	4-5	5-6	6-7	7-8
Air Temp. (C)	3.0, 7.0	7.0, 9.0	6.0, 9.0	6.0, 9.0	6.0, 9.0	3.0, 6.0	6.0, 11.5	6.0, 11.5	6.0, 11.5	6.0, 11.5	3.0, 6.0	6.0, 11.5	6.0, 11.5	6.0, 11.5	6.0, 11.5
Water Temp. (C)	5.5	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.0, 5.5	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.0, 5.5	5.5, 6.0	5.5, 6.0	5.5, 6.0	5.5, 6.0
Dissolved Oxygen (ppm)	9.4	9.4, 9.9	9.4, 9.9	9.4, 9.9	9.4, 9.9	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2	9.4, 10.2
pH	7.9, 8.2	8.2, 8.3	8.2, 8.3	8.2, 8.3	8.2, 8.3	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4	8.2, 8.4
Secchi disc (cm)	51, 56	82, 83	82, 83	82, 83	82, 83	33, 53	53	53	53	53	66, 66	51, 66	51, 66	51, 66	51, 66
Water stage (ft)	6.55, 6.62	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.55, 6.62	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.55, 6.62	6.67, 6.70	6.67, 6.70	6.67, 6.70	6.67, 6.70
Weather	Light rain,					Light rain,					Light rain,				
No. of fish	12	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Species	12	1	1	1	1	1	1	1	1	1	1	1	1	1	1
White sucker	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Shortnose sturgeon	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Channel catfish	6	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Parakeet fish	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Bluegill	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

POOR ORIGINAL

1565 069

Table 2.2-5

Fishes taken by trapnet on 26-28 April 1976 in the vicinity of TWINS.

Station	IN APR-1976			IN APR-1976			IN APR-1976			IN APR-1976			Total
	Date	Time	26-27	27-28	28-29	29-30	26-27	27-28	28-29	29-30	30-31	31-1	
Air Temp. (C)	0930-0950	0855-0910	3.5, 9.0	5.5, 7.5	8.0, 9.0	8.0, 9.0	6.0, 9.0	6.0, 8.5	6.0, 8.5	6.0, 8.5	6.0, 8.5	6.0, 8.5	57
	0930-0950	0855-0910	11.0, 15.0	10.0, 11.0	11.0, 15.0	11.0, 15.0	11.0, 15.0	11.0, 15.0	11.0, 15.0	11.0, 15.0	11.0, 15.0	11.0, 15.0	
	0930-0950	0855-0910	6.5, 8.1	6.1, 8.3	7.0, 8.9	7.0, 8.9	7.1, 8.9	8.9, 9.3	8.9, 9.3	8.9, 9.3	8.9, 9.3	8.9, 9.3	
	0930-0950	0855-0910	8.1, 8.2	8.1, 8.3	8.0, 8.1	8.0, 8.1	8.1, 8.2	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	
	0930-0950	0855-0910	30, 58	53, 58	38, 53	38, 53	38, 53	51, 61	51, 61	51, 61	51, 61	51, 61	
River Stage (ft)	0930-0950	0855-0910	4.33, 4.40	4.40, 4.48	4.33, 4.40	4.33, 4.40	4.33, 4.40	4.40, 4.48	4.40, 4.48	4.40, 4.48	4.40, 4.48	4.40, 4.48	57
	0930-0950	0855-0910	Overcast, clear	Partly cloudy	Overcast, clear	Overcast, clear	Overcast, clear	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	Partly cloudy	
No. of Spec.	15	11	3	3	3	3	3	3	3	3	3	3	6
No. of Spec.	3	5	3	3	3	3	3	3	3	3	3	3	
Carp	2	-	-	-	-	-	-	-	-	-	-	-	1
Brown bullhead	2	-	-	-	-	-	-	-	-	-	-	-	
Channel catfish	1	2	-	-	-	-	-	-	-	-	-	-	3
Rock bass	2	2	-	-	-	-	-	-	-	-	-	-	
Redbreast sunfish	1	2	-	-	-	-	-	-	-	-	-	-	3
Duck logperch	9	3	-	-	-	-	-	-	-	-	-	-	

Table 2.2-6

Fishes taken by trapnet on 10-12 May 1976 in the vicinity of TWINS.

Station	IN APR-1976			IN APR-1976			IN APR-1976			IN APR-1976			Total
	Date	Time	10-11	11-12	12-13	13-14	10-11	11-12	12-13	13-14	14-15	15-16	
Air Temp. (C)	1330-1350	1305-1325	22.0, 25.0	19.0, 22.0	17.0, 17.5	17.0, 17.5	22.0, 25.0	19.0, 22.0	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	68
	1330-1350	1305-1325	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	17.0, 17.5	
	1330-1350	1305-1325	9.2, 10.6	8.5, 9.2	7.9, 8.2	7.9, 8.2	9.2, 10.6	8.5, 9.2	7.9, 8.2	7.9, 8.2	7.9, 8.2	7.9, 8.2	
	1330-1350	1305-1325	8.2, 8.5	7.9, 8.2	7.9, 8.2	7.9, 8.2	8.2, 8.5	7.9, 8.2	7.9, 8.2	7.9, 8.2	7.9, 8.2	7.9, 8.2	
	1330-1350	1305-1325	91	86, 91	51, 61	51, 61	91	86, 91	51, 61	51, 61	51, 61	51, 61	
River Stage (ft)	1330-1350	1305-1325	4.61, 4.51	4.38, 4.41	4.61, 4.51	4.38, 4.41	4.61, 4.51	4.38, 4.41	4.61, 4.51	4.38, 4.41	4.61, 4.51	4.38, 4.41	68
	1330-1350	1305-1325	Overcast, clear	Overcast, clear	Overcast, clear	Overcast, clear	Overcast, clear	Overcast, clear	Overcast, clear	Overcast, clear	Overcast, clear	Overcast, clear	
No. of Spec.	11	10	3	3	3	3	3	3	3	3	3	3	31
No. of Spec.	3	6	3	3	3	3	3	3	3	3	3	3	
Carp	-	-	-	-	-	-	-	-	-	-	-	-	1
Spot tail shiner	-	1	-	-	-	-	-	-	-	-	-	-	
Yellow perch	-	-	-	-	-	-	-	-	-	-	-	-	1
Yellow perch	-	-	-	-	-	-	-	-	-	-	-	-	
Brown bullhead	-	-	-	-	-	-	-	-	-	-	-	-	2
Channel catfish	-	-	-	-	-	-	-	-	-	-	-	-	
Rock bass	-	1	-	-	-	-	-	-	-	-	-	-	1
Redbreast sunfish	-	1	-	-	-	-	-	-	-	-	-	-	
Pumpkinseed	5	3	-	-	-	-	5	3	-	-	-	-	8
Bluegill	1	3	-	-	-	-	1	3	-	-	-	-	
Smallmouth bass	-	-	-	-	-	-	-	-	-	-	-	-	1
White crappie	-	-	-	-	-	-	-	-	-	-	-	-	
Black crappie	-	-	-	-	-	-	-	-	-	-	-	-	1
Black crappie	-	-	-	-	-	-	-	-	-	-	-	-	

POOR ORIGINAL

1565 071

Table 2.2-7

Ectoparasite taken by pregnant on 26-28 May 1976 in the vicinity of Minc.

Date	26-27 1938-1939	27-28 1939-1940	28-29 1940-1941	29-30 1941-1942	30-31 1942-1943
Time	1638-1645	1646-1649	1650-1653	1654-1657	1658-1661
Air Temp. (°C)	12.0, 25.0	27.0, 25.0	21.5, 22.0	22.0	22.0, 26.0
Water Temp. (°C)	14.5, 16.0	16.0, 18.0	15.5, 16.0	16.0, 18.0	15.5, 17.5
Dissolved oxygen (ppm)	9.6	9.2	9.0	10.4	9.5
pH	7.5, 7.7	7.3, 7.5	7.0, 7.5	7.5, 7.7	7.5, 7.6
Sediment (cm)	63	61, 66	61, 66	61, 71	66, 81
Sludge layer (ft.)	5.00, 5.20	5.00, 5.00	5.00, 5.00	5.00, 5.20	5.00, 5.00
Weather	overcast,	clear,	light rain,	light rain,	clear,
No. of Spec.	15	10	10	10	10
No. of Spd.	3	6	5	2	2
Specimen shiner	-	-	0	1	6
"white sucker"	-	-	2	1	1-9
brown bullhead	-	-	-	-	3
channel catfish	2	1	-	2	5
Margined madtom	-	1	-	-	1-9
Rock bass	5	1	1	-	8
Belted sunfish	-	2	1	1	7
Redbreast blackfish	8	3	2	-	13-2
Invertebrates	-	-	-	-	51-2

Table 2.2-8

Fishes taken by trapnet on 7-8 June 1976 in the vicinity of TMIN.

Station	T-TAQ-1A2		T-TAQ-11A2		T-TAQ-11A3		T-TAQ-9B2		T-TAQ-9B3		T-TAQ-9B4		T-TAQ-9B5		T-TAQ-9B6		T-TAQ-9B7		T-TAQ-9B8		T-TAQ-9B9		T-TAQ-9B10		T-TAQ-9B11		T-TAQ-9B12		T-TAQ-9B13		T-TAQ-9B14		T-TAQ-9B15		T-TAQ-9B16		T-TAQ-9B17		T-TAQ-9B18		T-TAQ-9B19		T-TAQ-9B20		T-TAQ-9B21		T-TAQ-9B22		T-TAQ-9B23		T-TAQ-9B24		T-TAQ-9B25		T-TAQ-9B26		T-TAQ-9B27		T-TAQ-9B28		T-TAQ-9B29		T-TAQ-9B30		T-TAQ-9B31		T-TAQ-9B32		T-TAQ-9B33		T-TAQ-9B34		T-TAQ-9B35		T-TAQ-9B36		T-TAQ-9B37		T-TAQ-9B38		T-TAQ-9B39		T-TAQ-9B40		T-TAQ-9B41		T-TAQ-9B42		T-TAQ-9B43		T-TAQ-9B44		T-TAQ-9B45		T-TAQ-9B46		T-TAQ-9B47		T-TAQ-9B48		T-TAQ-9B49		T-TAQ-9B50		T-TAQ-9B51		T-TAQ-9B52		T-TAQ-9B53		T-TAQ-9B54		T-TAQ-9B55		T-TAQ-9B56		T-TAQ-9B57		T-TAQ-9B58		T-TAQ-9B59		T-TAQ-9B60		T-TAQ-9B61		T-TAQ-9B62		T-TAQ-9B63		T-TAQ-9B64		T-TAQ-9B65		T-TAQ-9B66		T-TAQ-9B67		T-TAQ-9B68		T-TAQ-9B69		T-TAQ-9B70		T-TAQ-9B71		T-TAQ-9B72		T-TAQ-9B73		T-TAQ-9B74		T-TAQ-9B75		T-TAQ-9B76		T-TAQ-9B77		T-TAQ-9B78		T-TAQ-9B79		T-TAQ-9B80		T-TAQ-9B81		T-TAQ-9B82		T-TAQ-9B83		T-TAQ-9B84		T-TAQ-9B85		T-TAQ-9B86		T-TAQ-9B87		T-TAQ-9B88		T-TAQ-9B89		T-TAQ-9B90		T-TAQ-9B91		T-TAQ-9B92		T-TAQ-9B93		T-TAQ-9B94		T-TAQ-9B95		T-TAQ-9B96		T-TAQ-9B97		T-TAQ-9B98		T-TAQ-9B99		T-TAQ-9B100		T-TAQ-9B101		T-TAQ-9B102		T-TAQ-9B103		T-TAQ-9B104		T-TAQ-9B105		T-TAQ-9B106		T-TAQ-9B107		T-TAQ-9B108		T-TAQ-9B109		T-TAQ-9B110		T-TAQ-9B111		T-TAQ-9B112		T-TAQ-9B113		T-TAQ-9B114		T-TAQ-9B115		T-TAQ-9B116		T-TAQ-9B117		T-TAQ-9B118		T-TAQ-9B119		T-TAQ-9B120		T-TAQ-9B121		T-TAQ-9B122		T-TAQ-9B123		T-TAQ-9B124		T-TAQ-9B125		T-TAQ-9B126		T-TAQ-9B127		T-TAQ-9B128		T-TAQ-9B129
---------	-----------	--	------------	--	------------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	-----------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------	--	-------------

POOR ORIGINAL
1565 072

Table 2-2-12

Tides taken by trawler on 2-4-48 at 1976 in the vicinity of 1948.

Station	7-82-11A1				7-82-11A2				7-82-11A3				7-82-11A4				7-82-11A5				7-82-11A6				7-82-11A7				7-82-11A8				7-82-11A9				7-82-11A10				7-82-11A11				7-82-11A12				7-82-11A13				7-82-11A14				7-82-11A15				7-82-11A16				7-82-11A17				7-82-11A18				7-82-11A19				7-82-11A20				7-82-11A21				7-82-11A22				7-82-11A23				7-82-11A24				7-82-11A25				7-82-11A26				7-82-11A27				7-82-11A28				7-82-11A29				7-82-11A30				7-82-11A31				7-82-11A32				7-82-11A33				7-82-11A34				7-82-11A35				7-82-11A36				7-82-11A37				7-82-11A38				7-82-11A39				7-82-11A40				7-82-11A41				7-82-11A42				7-82-11A43				7-82-11A44				7-82-11A45				7-82-11A46				7-82-11A47				7-82-11A48				7-82-11A49				7-82-11A50				7-82-11A51				7-82-11A52				7-82-11A53				7-82-11A54				7-82-11A55				7-82-11A56				7-82-11A57				7-82-11A58				7-82-11A59				7-82-11A60				7-82-11A61				7-82-11A62				7-82-11A63				7-82-11A64				7-82-11A65				7-82-11A66				7-82-11A67				7-82-11A68				7-82-11A69				7-82-11A70				7-82-11A71				7-82-11A72				7-82-11A73				7-82-11A74				7-82-11A75				7-82-11A76				7-82-11A77				7-82-11A78				7-82-11A79				7-82-11A80				7-82-11A81				7-82-11A82				7-82-11A83				7-82-11A84				7-82-11A85				7-82-11A86				7-82-11A87				7-82-11A88				7-82-11A89				7-82-11A90				7-82-11A91				7-82-11A92				7-82-11A93				7-82-11A94				7-82-11A95				7-82-11A96				7-82-11A97				7-82-11A98				7-82-11A99				7-82-11A100				7-82-11A101				7-82-11A102				7-82-11A103				7-82-11A104				7-82-11A105				7-82-11A106				7-82-11A107				7-82-11A108				7-82-11A109				7-82-11A110				7-82-11A111				7-82-11A112				7-82-11A113				7-82-11A114				7-82-11A115				7-82-11A116				7-82-11A117				7-82-11A118				7-82-11A119				7-82-11A120				7-82-11A121				7-82-11A122				7-82-11A123				7-82-11A124				7-82-11A125				7-82-11A126				7-82-11A127				7-82-11A128				7-82-11A129				7-82-11A130				7-82-11A131				7-82-11A132				7-82-11A133				7-82-11A134				7-82-11A135				7-82-11A136				7-82-11A137				7-82-11A138				7-82-11A139				7-82-11A140				7-82-11A141				7-82-11A142				7-82-11A143				7-82-11A144				7-82-11A145				7-82-11A146				7-82-11A147				7-82-11A148				7-82-11A149				7-82-11A150				7-82-11A151				7-82-11A152				7-82-11A153				7-82-11A154				7-82-11A155				7-82-11A156				7-82-11A157				7-82-11A158				7-82-11A159				7-82-11A160				7-82-11A161				7-82-11A162				7-82-11A163				7-82-11A164				7-82-11A165				7-82-11A166				7-82-11A167				7-82-11A168				7-82-11A169				7-82-11A170				7-82-11A171				7-82-11A172				7-82-11A173				7-82-11A174				7-82-11A175				7-82-11A176				7-82-11A177				7-82-11A178				7-82-11A179				7-82-11A180				7-82-11A181				7-82-11A182				7-82-11A183				7-82-11A184				7-82-11A185				7-82-11A186				7-82-11A187				7-82-11A188				7-82-11A189				7-82-11A190				7-82-11A191				7-82-11A192				7-82-11A193				7-82-11A194				7-82-11A195				7-82-11A196				7-82-11A197				7-82-11A198				7-82-11A199				7-82-11A200				7-82-11A201				7-82-11A202				7-82-11A203				7-82-11A204				7-82-11A205				7-82-11A206				7-82-11A207				7-82-11A208				7-82-11A209				7-82-11A210				7-82-11A211				7-82-11A212				7-82-11A213				7-82-11A214				7-82-11A215				7-82-11A216				7-82-11A217				7-82-11A218				7-82-11A219				7-82-11A220				7-82-11A221				7-82-11A222				7-82-11A223				7-82-11A224				7-82-11A225				7-82-11A226				7-82-11A227				7-82-11A228				7-82-11A229				7-82-11A230				7-82-11A231				7-82-11A232				7-82-11A233				7-82-11A234				7-82-11A235				7-82-11A236				7-82-11A237				7-82-11A238				7-82-11A239				7-82-11A240				7-82-11A241				7-82-11A242				7-82-11A243				7-82-11A244				7-82-11A245				7-82-11A246				7-
---------	-----------	--	--	--	-----------	--	--	--	-----------	--	--	--	-----------	--	--	--	-----------	--	--	--	-----------	--	--	--	-----------	--	--	--	-----------	--	--	--	-----------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	-------------	--	--	--	----

Table 2.2-13

Fishes taken by trawler on 10-18 August 1976 in the vicinity of BATES.

DATE		11 AUG - 1962					14 AUG - 1962					17 AUG - 1962					20 AUG - 1962				
Date	Time	16-17	17-18	18-19	19-20	20-21	16-17	17-18	18-19	19-20	20-21	16-17	17-18	18-19	19-20	20-21	16-17	17-18	18-19	19-20	20-21
Air Temp. (C)		22.5, 26.5	24.5, 29.5	20.5, 27.0	20.5, 21.5	22.5, 24.0	22.5, 24.0	24.0, 26.5	20.5, 21.5	20.5, 21.5	22.5, 24.0	22.5, 24.0	24.0, 26.5	20.5, 21.5	20.5, 21.5	22.5, 24.0	22.5, 24.0	24.0, 26.5	20.5, 21.5	20.5, 21.5	22.5, 24.0
Water Temp. (C)		20.5, 21.5	20.5, 22.0	7.5, 8.1	7.5, 7.9	7.5, 7.9	7.5, 7.9	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1	7.5, 8.1
Dissolved Oxygen (ppm)		8.0	8.0, 8.2	8.0	8.0	8.0	8.0	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2
pH		8.0	8.0, 8.2	8.0	8.0	8.0	8.0	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2	8.0, 8.2
Stock blue (cm)		18, 28	16, 31	18, 31	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28	20, 28
Water Stage (ft)		4.58, 4.52	4.58, 4.58	4.58, 4.58	4.58, 4.52	4.58, 4.52	4.58, 4.52	4.58, 4.58	4.58, 4.52	4.58, 4.52	4.58, 4.52	4.58, 4.52	4.58, 4.58	4.58, 4.52	4.58, 4.52	4.58, 4.52	4.58, 4.52	4.58, 4.58	4.58, 4.52	4.58, 4.52	4.58, 4.58
Weather		3	6	6	7	7	7	6	7	7	7	7	6	7	7	7	7	6	7	7	7
No. of Spec.		3	6	6	7	7	7	6	7	7	7	7	6	7	7	7	7	6	7	7	7
No. of Fish		3	6	6	7	7	7	6	7	7	7	7	6	7	7	7	7	6	7	7	7
Temp		3	6	6	7	7	7	6	7	7	7	7	6	7	7	7	7	6	7	7	7
Yellow perch		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
White sucker		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Channel catfish		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Rock bass		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Redbreast sunfish		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Bluegill		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Black crappie		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1
Yellow perch		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1

Table 2.2-16

Fish taken by trawler on 27-29 September 1976 in the vicinity of TONG.

Station	19 AUG 1976			20 AUG 1976			21 AUG 1976			22 AUG 1976			23 AUG 1976			24 AUG 1976			25 AUG 1976			26 AUG 1976			27 AUG 1976			28 AUG 1976			29 AUG 1976			Total			
	27-28	28-29	29-30	30-31	31-01	01-02	02-03	03-04	04-05	05-06	06-07	07-08	08-09	09-10	10-11	11-12	12-13	13-14	14-15	15-16	16-17	17-18	18-19	19-20	20-21	21-22	22-23	23-24	24-25	25-26	26-27	27-28	28-29	29-30	30-31	31-01	01-02
Time	0047-0817	0822-0916	0922-0946	0954-0957	1004-1006	1012-1014	1022-1024	1032-1034	1042-1044	1052-1054	1102-1104	1112-1114	1122-1124	1132-1134	1142-1144	1152-1154	1202-1204	1212-1214	1222-1224	1232-1234	1242-1244	1252-1254	1302-1304	1312-1314	1322-1324	1332-1334	1342-1344	1352-1354	1402-1404	1412-1414	1422-1424	1432-1434	1442-1444	1452-1454	1502-1504	1512-1514	1522-1524
Air Temp. (°C)	14.5, 16.5	9.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	14.5, 14.5	
Water Temp. (°C)	15.0, 17.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	14.0, 15.0	
Dissolved Oxygen (ppm)	7.0, 7.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	7.0, 9.4	
pH	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	8.0, 8.1	
Secchi Disc (cm)	10, 36	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	
Water State (ft)	3.61, 3.72	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85	3.72, 3.85		
Weather	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	Light rain,	
No. of Spec.	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Golden shiner	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Yellow perch	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Rock bass	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
White sucker	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Blackchin shiner	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Crayfish	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Bluegill	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Shiner	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		
Bluegill	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1		

POOR ORIGINAL

1565 079

Table 2.2-17

stages taken by trapnet on 18-20 October 1976 in the vicinity of IMN.

[illegible]

Table 2.2-10

specimens taken by trapnet on 25-27 October 1976 in the vicinity of TMS.

[illegible]

POOR ORIGINAL

1565 080

POOR ORIGINAL

Table 2.2-19

Summary of fishes taken at trapnet station TN-AQF-1A3 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	% Catch
Carp	1	-	1	2	-	1	2	-	7	2.4
Golden shiner	-	-	-	-	1	-	-	-	1	0.3
Spottail shiner	6	-	1	-	-	-	-	-	7	2.4
Quillback	-	-	-	-	3	2	-	-	5	1.7
White sucker	2	-	-	1	-	1	1	-	5	1.7
Northern hog sucker	2	1	-	-	-	-	-	-	3	1.0
Shorthead redhorse	1	-	-	-	-	-	-	-	1	0.3
White catfish	-	1	-	-	-	-	1	-	2	0.7
Yellow bullhead	-	-	-	-	-	-	2	-	2	0.7
Brown bullhead	1	2	3	1	4	2	-	-	13	4.4
Channel catfish	23	3	4	3	1	10	8	10	62	21.0
Margined madtom	1	-	1	-	-	-	-	-	2	0.7
Rock bass	-	7	8	7	-	7	5	6	40	13.6
Redbreast sunfish	-	3	11	3	1	4	2	-	24	8.1
Pumpkinseed	1	15	18	11	11	18	29	-	103	34.9
Bluegill	-	-	1	-	1	-	-	-	2	0.7
White crappie	2	-	-	-	-	-	-	-	2	0.7
Black crappie	-	-	-	-	-	1	11	-	12	4.1
Galley	-	-	-	-	-	-	2	-	2	0.7
No. of Spmn.	10	32	48	28	22	46	63	16	295	
No. of Spp.	10	7	9	7	7	9	10	2	19	
No. of Coll.	6	4	4	4	4	4	6	4	36	
n/Coll.	6.67	8.00	12.00	7.00	5.50	11.50	10.50	4.00	8.19	
Diversity Index	2.18	2.19	2.46	2.34	2.13	2.48	2.44	0.95	2.95	

Table 2.2-20

Summary of fishes taken at trapnet station TN-AQF-11A2 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	% Catch
Blackchin shiner	-	-	-	-	-	-	1	-	1	0.5
Carp	-	-	-	-	-	1	-	-	1	0.5
Golden shiner	-	-	-	-	1	2	1	-	4	2.2
Spottail shiner	-	6	-	-	-	-	-	1	7	3.8
Spotfin shiner	-	-	6	-	-	-	-	-	6	3.2
Quillback	-	-	-	-	-	1	-	-	1	0.5
White sucker	-	-	-	-	-	2	-	-	2	1.1
Shorthead redhorse	1	-	-	1	-	-	-	-	2	1.1
Brown bullhead	-	-	-	3	-	-	-	-	3	1.6
Channel catfish	3	7	-	4	3	7	1	21	46	24.9
Margined madtom	-	-	-	1	-	-	-	-	1	0.5
Rock bass	-	7	2	5	6	1	7	-	28	15.1
Redbreast sunfish	-	2	4	-	-	3	7	-	16	8.6
Pumpkinseed	-	-	4	7	9	4	4	-	28	15.1
Bluegill	-	-	-	-	2	2	7	-	11	5.9
Smallmouth bass	-	-	-	1	-	-	-	-	1	0.5
Largemouth bass	-	-	-	-	1	-	-	-	1	0.5
White crappie	1	-	-	1	-	1	1	-	4	2.2
Black crappie	-	-	-	6	2	5	6	1	20	10.8
Galley	-	-	-	1	-	1	-	-	2	1.1
No. of Spmn.	5	22	36	30	24	30	35	23	185	
No. of Spp.	3	4	4	10	7	12	9	3	20	
No. of Coll.	6	4	4	4	4	4	6	4	36	
n/Coll.	0.83	5.50	4.00	7.50	6.00	7.50	5.83	5.75	5.14	
Diversity Index	1.37	1.88	1.91	2.97	2.39	3.24	2.77	0.51	3.35	

1565 081

POOR ORIGINAL

Table 2.2-21

Summary of fishes taken at trapnet station TM-AQF-11A3 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	Catch
Carp	-	1	-	-	2	1	-	-	4	5.3
Golden shiner	-	-	-	-	1	1	-	-	2	2.7
Spotfin shiner	-	-	-	-	1	-	-	-	1	1.3
Quillback	-	-	-	-	-	-	1	-	1	1.3
White sucker	-	-	1	1	-	-	-	-	2	2.7
Brown bullhead	-	-	-	2	-	-	-	-	2	2.7
Channel catfish	1	2	-	2	-	-	-	1	6	8.0
Margined madtom	2	-	-	1	1	-	-	-	4	5.3
Rock bass	2	1	3	5	1	-	1	-	13	17.3
Redbreast sunfish	1	1	12	4	1	1	2	-	22	29.3
Pumpkinseed	-	-	1	-	3	-	7	-	11	14.7
Bluegill	1	-	-	-	1	-	-	-	2	2.7
Smallmouth bass	-	-	1	-	-	-	-	-	1	1.3
Black crappie	-	-	-	2	2	-	-	-	4	5.3
No. of Spmn.	7	5	18	17	13	3	11	1	75	
No. of Spp.	5	4	5	7	9	3	4	1	14	
No. of Coll.	6	4	4	4	4	4	6	4	36	
n/Coll.	1.17	1.25	4.50	4.25	3.25	0.75	1.83	0.25	2.08	
Diversity Index	2.24	1.92	1.52	2.54	3.03	1.59	1.49	0.00	3.14	

Table 2.2-22

Summary of fishes taken at trapnet station TM-AQF-952 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	Catch
Alewife	-	-	-	-	-	-	2	-	2	0.7
Carp	-	-	-	-	1	-	3	2	6	2.2
Golden shiner	-	-	-	-	2	-	9	-	11	4.1
Spottail shiner	-	-	-	-	1	-	-	-	1	0.4
Quillback	-	-	1	1	1	2	7	-	12	4.5
White catfish	1	-	-	-	1	-	-	-	2	0.7
Yellow bullhead	-	-	2	-	-	-	-	-	2	0.7
Brown bullhead	2	6	1	5	5	1	3	-	23	8.6
Channel catfish	5	7	5	2	9	-	3	4	35	13.1
Margined madtom	-	-	-	-	-	-	-	1	1	0.4
Rock bass	1	2	4	1	1	-	6	-	15	5.6
Redbreast sunfish	-	2	7	-	-	-	-	-	9	3.4
Pumpkinseed	4	12	31	3	10	4	8	-	72	27.0
Bluegill	-	-	2	-	1	3	1	1	8	3.0
Smallmouth bass	1	-	-	-	-	-	-	-	1	0.4
White crappie	-	-	1	-	1	-	12	-	14	5.2
Black crappie	2	-	5	5	17	5	13	-	47	17.6
Yellow perch	-	-	-	-	2	1	-	-	3	1.1
Walleye	1	-	-	-	-	-	1	1	3	1.1
No. of Spmn.	17	29	59	17	52	16	68	9	267	
No. of Spp.	9	5	10	6	13	8	12	5	19	
No. of Coll.	6	4	4	4	4	4	6	4	36	
n/Coll.	2.83	7.25	14.75	4.25	13.00	4.00	11.33	2.25	7.42	
Diversity Index	2.70	2.02	2.25	2.32	2.83	2.35	2.22	2.06	3.32	

1565 082

Table 2.2-23

Summary of fishes taken at trapnet stations during March through October 1976.

Station	TH-AQF-1A3	TH-AQF-11A2	TH-AQF-11A3	TH-AQF-2B2	Total	% Catch
Alewife	-	-	-	2	2	0.2
Muskellunge	-	1	-	-	1	0.1
Carp	7	1	4	6	18	2.2
Golden shiner	1	4	2	11	18	2.2
Spottail shiner	7	7	-	1	15	1.8
Spotfin shiner	-	6	1	-	7	0.9
Quillback	5	1	1	12	19	2.3
White sucker	5	2	2	-	9	1.1
Northern hog sucker	3	-	-	-	3	0.4
Shorthead redhorse	1	2	-	-	3	0.4
White catfish	2	-	-	2	4	0.5
Yellow bullhead	2	-	-	2	4	0.5
Brown bullhead	13	3	2	23	41	5.0
Channel catfish	62	46	6	35	149	18.1
Margined madtom	2	1	4	1	8	1.0
Rock bass	50	28	13	15	96	11.7
Redbreast sunfish	24	16	22	9	71	8.6
Pumpkinseed	103	28	11	72	214	26.0
Bluegill	2	11	2	8	23	2.8
Smallmouth bass	-	1	1	1	3	0.4
Largemouth bass	-	1	-	-	1	0.1
White crappie	2	4	-	14	20	2.4
Black crappie	12	20	4	47	83	10.1
Yellow perch	-	-	-	3	3	0.4
Walleye	2	2	-	3	7	0.9
No. of Spmn.	295	185	75	267	822	
No. of Spp.	19	20	14	19	25	
No. of Coll.	36	36	36	36	144	
Diversity Index	2.95	3.35	3.14	3.32	3.41	

^aAsphyxial loss of fishes taken at trapnet stations upstream (U) and downstream (D) from the discharge during March through October 1976.

Month	March		April		May		June		July		August		September		October		Total catch	Total	CATCH	TOTAL	CATCH	TOTAL	CATCH	TOTAL
	E	P	E	P	E	P	E	P	E	P	E	P	E	P	E	P								
Albacore	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Basking shark	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Carp	1	-	1	1	-	-	2	-	1	1	1	1	2	3	2	2	7	2.5	11	2.1	16	2.2	1	
Golden shiner	-	-	-	-	-	-	-	-	1	4	-	-	-	-	-	-	7	2.5	17	3.2	18	2.2	1	
Silver shiner	6	-	6	1	-	-	-	-	-	-	-	-	-	-	-	-	7	2.5	6	1.5	15	1.8	1	
Spot tail shiner	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
Goldfish	-	-	-	-	-	-	-	-	1	1	2	3	-	-	-	-	5	1.7	7	1.3	7	0.9	1	
White sucker	2	-	-	-	-	-	-	-	1	1	2	1	6	-	-	-	5	1.7	14	2.7	19	2.3	1	
Yellow perch	1	-	1	1	-	-	1	-	-	-	-	-	-	-	-	-	3	1.0	5	0.6	9	1.1	1	
Shortfin mako	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	1.0	-	-	-	-	1	
White catfish	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4	0.3	2	0.4	3	0.4	1	
Yellow catfish	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.2	2	0.4	5	0.5	1	
Brown bullhead	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.2	2	0.4	4	0.5	1	
Channel catfish	2	-	2	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.2	2	0.4	4	0.5	1	
White catfish	21	9	3	16	5	8	1	10	5	5	1	12	10	7	8	5	33	5.5	28	5.1	51	8.0	1	
Striped bass	1	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	11	6.2	21.0	27	16.5	17.4	18.1	
Rock bass	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1	0.2	1	0.4	6	1.1	1	
Smallmouth bass	1	7	10	8	9	7	11	-	8	7	1	5	15	6	-	-	50	15.6	56	16.6	96	11.7	1	
Parkinson	1	3	5	11	23	5	4	1	4	2	5	2	9	-	-	-	25	8.1	57	8.9	71	8.6	1	
Bluegill	1	4	15	12	18	16	11	10	11	27	16	8	29	19	-	-	101	32.9	111	31.1	151	26.0	1	
Black crappie	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.3	21	4.0	23	2.8	1	
White crappie	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	0.6	3	0.6	3	0.3	1	
Black sculpin	2	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.7	18	3.5	20	2.4	1	
Yellow perch	2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	12	4.1	21	11.5	83	16.1	1	
White perch	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3	0.6	3	0.6	3	0.3	1	
Bluegill	1	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	2	0.7	5	0.9	7	0.9	1	
Sum of sp.	40	26																						

Table 2.2-25

Indices of percent similarity of species composition between trapnet stations during 1974, 1975, and 1976.

1974				1975		
		82.5	11A2	38.7		
	70.9	65.1	11A3	64.8	45.5	
57.2	73.0	55.4	9B2	67.1	46.9	63.8
11A3	11A2	1A3		1A3	11A2	11A3
49.4	60.3	66.1	9B2			
	62.6	58.3	11A3			
		71.2	11A2			
1976						

POOR ORIGINAL

Table 2.2-26

Indices of percent similarity of species composition between seine stations during 1974, 1975, and 1976.

1974				1975			
		84.0	16A1	44.7			
		88.9	77.0	10A2	61.0	70.3	
	71.1	71.3	64.1	9A1	71.0	37.8	44.4
84.4	72.9	69.7	60.1	9B3	56.5	73.3	83.2
9A1	10A2	16A1	1A2		1A2	16A1	10A2
79.3	77.9	67.8	71.4	9B3			
	82.0	79.4	68.2	9A1			
		70.5	80.9	10A2			
1976							
		64.4	16A1				

1565 085

Table 2.2-27

Fishes taken by seine on 2 March 1976 in the vicinity of TMNS.

Station	TM-ACF-1A2	TM-ACF-1A1	TM-ACF-1A2	TM-ACF-9A1	TM-ACF-9B3	Total	Catch
Time	1010	1030	1045	1115	1135		
Air Temp. (C)	6.0	6.0	5.5	6.0	5.0		
Water Temp. (C)	6.5	6.5	6.5	6.5	6.5		
Dissolved Oxygen (ppm)	8.8	9.0	9.7	9.3	8.6		
pH	8.0	8.1	8.1	8.0	7.9		
Secchi Disc (cm)	51	51	51	36	51		
River Stage (ft)	6.80	6.80	6.80	6.80	6.80		
Weather	Fog	Fog	Fog	Fog	Fog		
No. of Spmn.	11	16	7	3	44	81	
No. of Sp.	2	6	3	2	6	10	
Comely shiner	-	3	-	-	-	3	3.7
Common shiner	-	-	-	-	1	1	1.2
Spottail shiner	-	2	-	-	1	3	3.7
Swallowtail shiner	-	1	-	1	7	9	11.1
Spotfin shiner	5	8	5	-	33	51	63.0
Bluntnose minnow	3	-	1	-	1	5	6.2
Blacknose dace	-	1	-	-	-	1	1.2
Redbreast sunfish	-	1	-	-	1	2	2.5
Tessellated darter	-	-	1	2	-	3	3.7
Banded darter	3	-	-	-	-	3	3.7

POOR ORIGINAL

Table 2.2-28

Fishes taken by seine on 15 March 1976 in the vicinity of TMNS.

Station	TM-ACF-1A2	TM-ACF-1A1	TM-ACF-1A2	TM-ACF-9A1	TM-ACF-9B3	Total	Catch
Time	1030	1045	1100	1130	1200		
Air Temp. (C)	11.0	12.0	13.0	13.0	11.5		
Water Temp. (C)	6.0	5.5	6.0	6.0	6.0		
Dissolved Oxygen (ppm)	10.8	10.8	10.0	9.8	9.5		
pH	7.8	7.7	7.6	7.8	7.8		
Secchi Disc (cm)	61	61	43	61	64		
River Stage (ft)	5.84	5.84	5.84	5.84	5.84		
Weather	Clear	Clear	Clear	Clear	Partly Cloudy		
No. of Spmn.	24	24	32	31	-	111	
No. of Sp.	3	4	7	4	-	9	
Comely shiner	-	2	-	-	-	2	1.5
Spottail shiner	-	-	4	1	-	5	4.5
Swallowtail shiner	3	3	2	2	NO	10	9.0
Emeryface shiner	-	-	4	-	-	4	3.0
Spotfin shiner	15	18	14	26	-	73	65.8
Bluntnose minnow	6	-	4	2	FISH	12	10.8
Fathead minnow	-	1	-	-	-	1	0.9
Redbreast sunfish	-	-	2	-	-	2	1.8
Tessellated darter	-	-	2	-	TAKEN	2	1.8

1565 086

Table 2.2-29

Fishes taken by seine on 29 March 1976 in the vicinity of TMINS.

Station	TM-AQF-1A2	TM-AQF-1A1	TM-AQF-1A2	TM-AQF-9A1	TM-AQF-9B3	Total	Mean
Time	1020	1040	1055	1135	1140		
Air Temp. (C)	12.0	12.0	12.0	13.0	13.0		
Water Temp. (C)	10.5	10.5	10.0	10.5	10.5		
Dissolved Oxygen (ppm)	8.8	9.2	9.7	9.1	8.8		
pH	7.9	7.9	7.9	7.8	7.8		
Secchi Disc (cm)	33	41	41	36	46		
River Stage (ft)	5.52	5.52	5.52	5.52	5.52		
Weather	Overcast	Overcast	Overcast	Overcast	Overcast		
No. of Spmn.	13	16	30	4	13	76	
No. of Sp.	3	4	4	2	3	16	
Spottail shiner	-	1	11	-	-	12	22.3
Swallowtail shiner	-	-	1	-	6	7	7.3
Spotfin shiner	8	1	13	-	6	28	29.2
Bluntnose minnow	-	4	8	1	-	13	13.8
Blacknose dace	1	-	-	-	-	1	1.0
Creek chub	-	-	1	-	-	1	1.0
Channel catfish	-	-	1	-	-	1	1.0
Tessellated darter	4	10	2	3	1	20	27.6
Banded darter	-	-	3	-	-	3	3.1

POOR ORIGINAL

Table 2.2-30

Fishes taken by seine on 13 April 1976 in the vicinity of TMINS.

Station	TM-AQF-1A2	TM-AQF-1A1	TM-AQF-1A2	TM-AQF-9A1	TM-AQF-9B3	Total	Mean
Time	0922	0935	0945	1020	1030		
Air Temp. (C)	9.5	10.0	10.5	10.0	11.5		
Water Temp. (C)	7.0	8.0	8.0	8.0	8.0		
Dissolved Oxygen (ppm)	10.7	10.8	11.0	10.5	10.4		
pH	7.7	7.6	7.7	7.7	7.7		
Secchi Disc (cm)	NA	122	137	137	137		
River Stage (ft)	4.83	4.83	4.83	4.83	4.83		
Weather	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy		
No. of Spmn.	1	6	35	1	5	48	
No. of Sp.	1	3	6	1	3	14	
Spottail shiner	-	-	13	-	2	15	15.7
Swallowtail shiner	-	1	1	-	1	3	3.3
Rosyface shiner	1	-	-	-	-	1	1.0
Spotfin shiner	-	4	11	-	2	17	17.4
Bluntnose minnow	-	-	2	-	-	2	2.2
Pumpkinseed	-	-	-	1	-	1	1.0
Tessellated darter	-	1	2	-	-	3	3.3
Banded darter	-	-	1	-	-	1	1.0

1565 087

Table 2.2-31

Fishes taken by seine on 30 April 1976 in the vicinity of THINS.

Station	TH-AQF-1A2	TH-AQF-1A3	TH-AQF-1A4	TH-AQF-9A1	TH-AQF-9B3	Total	Catch
Time	1450	1440	1422	1400	1350		
Air Temp. (C)	20.0	20.5	20.5	19.0	19.0		
Water Temp. (C)	15.0	15.0	14.0	14.5	14.5		
Dissolved Oxygen (ppm)	10.0	9.9	9.7	10.6	10.2		
pH	7.5	7.5	7.6	7.9	8.0		
Secchi Disc (cm)	61	38	48	61	64		
River Stage (ft)	5.28	5.28	5.28	5.28	5.28		
Weather	Clear	Clear	Clear	Clear	Clear		
No. of Spmn.	17	2	2	1	3	4	
No. of Spn.	3	2	5	1	1	4	
Spottail shiner	-	-	-	-	-	-	3.2
Swallowtail shiner	7	-	1	-	-	8	16.3
Spotfin shiner	6	1	19	1	3	30	61.3
Bluntnose minnow	-	-	1	-	-	1	2.0
Smallmouth bass	-	-	1	-	-	1	2.0
Tessellated darter	4	1	-	-	-	5	10.3

Table 2.2-32

Fishes taken by seine on 10 May 1976 in the vicinity of THINS.

Station	TH-AQF-1A2	TH-AQF-1A3	TH-AQF-1A4	TH-AQF-9A1	TH-AQF-9B3	Total	Catch
Time	1035	1015	1000	0942	0930		
Air Temp. (C)	22.0	23.0	20.0	19.0	18.0		
Water Temp. (C)	16.5	16.0	16.0	15.5	15.0		
Dissolved Oxygen (ppm)	8.7	9.2	8.4	9.1	8.6		
pH	7.9	7.9	7.9	7.9	8.0		
Secchi Disc (cm)	107	107	107	107	107		
River Stage (ft)	4.51	4.51	4.51	4.51	4.51		
Weather	Clear	Clear	Clear	Clear	Clear		
No. of Spmn.	1	-	2	-	-	3	
No. of Spn.	1	-	2	-	-	3	
Bluntnose minnow	-	-	1	-	-	1	33.3
Tessellated darter	1	NO FISH TAKEN	1	NO FISH TAKEN	-	2	11.4

POOR ORIGINAL

Table 2.2-33

Fishes taken by seine on 24 May 1976 in the vicinity of THINS.

Station	TH-AQF-1A2	TH-AQF-1A3	TH-AQF-1A4	TH-AQF-9A1	TH-AQF-9B3	Total	Catch
Time	1040	1025	1005	0943	0927		
Air Temp. (C)	16.5	16.0	15.0	15.0	15.0		
Water Temp. (C)	16.5	16.0	15.5	15.5	16.0		
Dissolved Oxygen (ppm)	NA	NA	NA	NA	NA		
pH	7.5	7.5	7.5	7.5	7.7		
Secchi Disc (cm)	33	41	41	51	56		
River Stage (ft)	5.83	5.83	5.83	5.83	5.83		
Weather	Clear	Clear	Clear	Clear	Clear		
No. of Spmn.	3	1	4	-	37	45	
No. of Spn.	3	1	2	-	5	7	
Spottail shiner	-	-	3	-	1	4	7.7
Swallowtail shiner	1	-	-	NO	-	1	2.2
Spotfin shiner	1	-	-	-	-	1	2.2
Creek chub	-	-	-	FISH	4	4	8.9
White sucker/Shorthead redhorse	-	-	-	-	30	30	66.7
Smallmouth bass	1	1	1	TAKEN	1	4	8.9
Tessellated darter	-	-	-	-	1	1	2.2

NA = Not Available

1565 088

Table 2.2-34

Fishes taken by seine on 7 June 1976 in the vicinity of TMINS.

Station	TM-AQF-1A2	TM-AQF-16A1	TM-AQF-10A2	TM-AQF-9A1	TM-AQF-9B1	Total	Count
Time	1040	1030	1010	0945	0930		
Air Temp. (C)	22.5	22.0	22.5	21.5	21.5		
Water Temp. (C)	19.5	19.5	19.0	19.0	19.5		
Dissolved Oxygen (ppm)	8.2	7.9	8.7	7.9	8.2		
pH	8.2	8.1	7.9	8.0	8.1		
Secchi Disc (cm)	61	56	51	56	61		
River Stage (ft)	4.69	4.69	4.69	4.69	4.69		
Weather	Clear	Clear	Clear	Clear	Clear		
No. of Spmn.	79	635	219	333	876	2192	
No. of Spp.	7	4	9	6	4	11	
Spottail shiner	36	339	77	240	773	1465	39.2
Shallowtail shiner	2	-	-	-	-	2	0.1
Spotfin shiner	26	8	6	-	-	40	1.3
Bluntnose minnow	-	-	4	-	-	4	0.2
Longnose dace	1	-	2	-	7	10	0.3
Creek chub	-	-	1	4	12	17	0.4
White sucker/Shorthead redhorse	8	324	123	85	84	624	28.5
Redbreast sunfish	-	-	2	-	-	2	0.1
Bluegill	-	-	-	1	-	1	-
Tessellated darter	5	-	3	1	-	9	0.2
Shield darter	1	14	1	2	-	18	0.4

+ = less than 0.05%.

POOR ORIGINAL

Table 2.2-35

Fishes taken by seine on 21 June 1976 in the vicinity of TMINS.

Station	TM-AQF-1A2	TM-AQF-16A1	TM-AQF-10A2	TM-AQF-9A1	TM-AQF-9B1	Total	Count
Time	1030	1015	1000	0932	0920		
Air Temp. (C)	25.5	26.0	25.0	24.5	24.5		
Water Temp. (C)	25.0	24.0	25.0	25.0	25.0		
Dissolved Oxygen (ppm)	7.8	6.4	7.0	6.5	6.1		
pH	8.7	8.1	8.3	8.5	8.2		
Secchi Disc (cm)	30	30	30	30	30		
River Stage (ft)	4.42	4.42	4.42	4.42	4.42		
Weather	Overcast	Overcast	Overcast	Overcast	Overcast		
No. of Spmn.	391	306	403	89	230	1419	
No. of Spp.	6	4	7	6	5	12	
Common shiner	-	3	1	-	-	4	0.3
Spottail shiner	371	261	372	77	193	1274	39.2
Spotfin shiner	2	8	6	5	8	29	0.2
Blacknose dace	-	1	-	-	-	1	0.1
Longnose dace	-	1	-	-	-	1	0.1
Creek chub	-	2	8	2	12	24	1.0
White sucker	-	1	1	-	-	2	0.1
Shorthead redhorse	-	16	-	-	-	16	1.2
White sucker/Shorthead redhorse	4	-	10	3	16	33	1.3
Pumpkinseed	1	-	-	-	-	1	0.1
Smallmouth bass	1	-	-	1	-	2	0.1
Tessellated darter	12	11	5	1	1	30	0.2
Shield darter	-	2	-	-	-	2	0.1

1565 089

Table 2.2-36

Fishes taken by seine on 7 July 1976 in the vicinity of THINS.

Station	TH-AOP-1A2	TH-AOP-1A1	TH-AOP-1A2	TH-AOP-1A1	TH-AOP-1A2	Total	std
Time	0935	1020	1040	1115	1130		
Air Temp. (C)	22.0	21.5	21.5	22.5	24.0		
Water Temp. (C)	24.0	24.5	24.5	24.5	25.0		
Dissolved Oxygen (ppm)	6.8	6.2	6.8	7.0	6.9		
pH	7.3	7.6	7.3	7.3	7.2		
Secchi Disc (cm)	36	30	30	33	30		
River Stage (ft)	4.46	4.46	4.46	4.46	4.46		
Weather	Overcast	Overcast	Overcast	Overcast	Overcast		
No. of Spmn.	636	515	179	19	359	1678	
No. of Spp.	7	10	9	7	8	17	
Comely shiner	-	23	-	-	-	23	1.1
Common shiner	1	-	-	-	-	1	0.1
Spottail shiner	595	355	150	61	200	1361	81.1
Spotfin shiner	31	3	1	1	-	36	2.1
Bluntnose minnow	-	2	1	-	-	3	0.2
Blacknose dace	1	-	-	-	-	1	0.1
Fallfish	-	1	4	8	3	16	1.1
White sucker	-	-	8	6	131	145	8.6
Northern hog sucker	-	1	-	-	-	1	0.1
Shorthead redhorse	-	-	-	-	1	1	0.1
White sucker/Shorthead redhorse	-	10	6	-	-	16	1.0
Brown bullhead	-	-	-	-	4	4	0.2
Channel catfish	-	-	5	-	7	12	0.7
Rock bass	1	-	-	-	-	1	0.1
Smallmouth bass	5	9	2	3	5	24	1.4
White crappie	-	-	-	3	-	3	0.2
Tessellated darter	2	9	2	7	8	28	1.7
Banded darter	-	-	-	-	-	2	0.1

Table 2.2-37

Fishes taken by seine on 19 July 1976 in the vicinity of THINS.

Station	TH-AOP-1A2	TH-AOP-1A1	TH-AOP-1A2	TH-AOP-1A1	TH-AOP-1A2	Total	std
Time	1446	1355	1337	1308	1253		
Air Temp. (C)	24.5	29.0	25.5	27.5	27.0		
Water Temp. (C)	25.0	24.0	24.5	24.5	24.5		
Dissolved Oxygen (ppm)	7.4	7.2	7.4	7.4	7.2		
pH	7.8	7.8	7.8	7.7	7.6		
Secchi Disc (cm)	30	23	28	28	30		
River Stage (ft)	4.51	4.51	4.51	4.51	4.51		
Weather	Clear	Clear	Clear	Clear	Clear		
No. of Spmn.	218	164	124	73	30	679	
No. of Spp.	10	17	10	9	7	21	
Carp	-	-	1	-	-	1	0.1
Comely shiner	4	3	2	-	-	9	1.3
Common shiner	4	2	-	-	-	6	0.9
Spottail shiner	218	128	83	52	12	493	72.6
Spotfin shiner	47	3	2	3	-	55	8.1
Bluntnose minnow	2	-	-	2	1	5	0.7
Blacknose dace	1	-	-	-	-	1	0.1
Creek chub	-	-	1	-	-	1	0.1
Fallfish	-	3	-	1	-	4	0.6
Quillback	-	1	-	-	-	1	0.1
White sucker	4	3	2	6	5	20	2.9
Shorthead redhorse	1	6	23	4	2	36	5.3
Brown bullhead	-	-	-	-	3	3	0.4
Rock bass	-	1	-	-	-	1	0.1
Redbreast sunfish	-	-	-	1	-	1	0.1
Pumpkinseed	1	-	-	-	-	1	0.1
Smallmouth bass	6	3	3	1	1	14	2.1
Largemouth bass	-	1	-	-	-	1	0.1
White crappie	-	-	1	-	-	1	0.1
Tessellated darter	-	9	6	3	6	24	3.5
Banded darter	-	1	-	-	-	1	0.1

POOR ORIGINAL

1565 090

Table 2.2-38

Fishes taken by seine on 2 August 1976 in the vicinity of TMINS.

Station	TM-AQF-1A2	TM-AQF-16A1	TM-AQF-10A2	TM-AQF-9A1	TM-AQF-9B3	Total	Catch
Time	1005	1028	1042	1108	1122		
Air Temp. (C)	20.0	19.5	20.5	20.0	20.5		
Water Temp. (C)	21.0	21.0	21.0	21.0	21.0		
Dissolved Oxygen (ppm)	8.0	7.8	8.2	8.0	7.9		
pH	8.5	8.4	8.1	8.1	8.0		
Secchi Disc (cm)	30	33	33	30	30		
River Stage (ft)	3.96	3.96	3.96	3.96	3.96		
Weather	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy		
No. of Spmn.	48	125	55	61	271	570	
No. of Sp.	7	9	6	8	7	16	
Common shiner	1	-	-	-	-	1	0.2
Common shiner	5	3	-	-	-	8	1.4
Spottail shiner	19	97	35	28	8	187	32.8
Spotfin shiner	1	-	-	-	-	1	0.2
Bluntnose minnow	-	7	2	3	-	12	2.1
Fallfish	1	2	-	-	-	3	0.5
White sucker	-	1	2	4	1	8	1.4
Northern hog sucker	-	-	-	1	-	1	0.2
Shorthead redhorse	-	4	6	4	-	14	2.5
Yellow bullhead	-	-	-	-	2	2	0.4
Brown bullhead	-	-	-	-	2	2	0.4
Channel catfish	-	-	-	3	256	259	45.4
Redbreast sunfish	-	1	-	-	-	1	0.2
Smallmouth bass	2	1	7	1	-	11	1.9
White crappie	-	-	-	-	1	1	0.2
Tessellated darter	19	9	13	17	1	59	10.4

POOR ORIGINAL

Table 2.2-39

Fishes taken by seine on 16 August 1976 in the vicinity of TMINS.

Station	TM-AQF-1A2	TM-AQF-16A1	TM-AQF-10A2	TM-AQF-9A1	TM-AQF-9B3	Total	Catch
Time	1045	1000	0945	0920	0905		
Air Temp. (C)	18.5	18.5	18.5	18.0	18.0		
Water Temp. (C)	19.5	20.0	20.0	19.5	20.0		
Dissolved Oxygen (ppm)	7.4	7.2	7.3	7.5	7.8		
pH	8.0	8.2	8.0	8.3	8.4		
Secchi Disc (cm)	28	28	28	28	28		
River Stage (ft)	4.52	4.52	4.52	4.52	4.52		
Weather	Clear	Clear	Clear	Clear	Clear		
No. of Spmn.	32	124	108	51	104	419	
No. of Sp.	9	12	10	12	8	18	
River chub	1	-	1	-	-	2	0.5
Common shiner	2	2	-	-	1	5	1.2
Common shiner	3	6	-	-	1	10	2.4
Spottail shiner	5	71	64	20	11	171	40.8
Swallowtail shiner	2	-	-	-	-	2	0.5
Spotfin shiner	12	23	3	5	12	55	13.1
Bluntnose minnow	5	3	1	1	23	33	7.9
Fallfish	-	-	1	1	-	2	0.5
White sucker	-	-	1	-	-	1	0.2
Shorthead redhorse	-	2	6	4	-	12	2.9
Rock bass	-	4	-	-	-	4	1.0
Redbreast sunfish	-	-	-	1	-	1	0.2
Pumpkinseed	-	-	1	-	-	1	0.2
Bluegill	-	1	-	2	-	3	0.7
Smallmouth bass	-	2	4	1	2	9	2.1
White crappie	-	1	-	4	18	23	5.5
Black crappie	1	1	-	-	-	2	0.5
Tessellated darter	1	8	26	12	36	83	19.8

1565 091

Table 2.2-40

Fishes taken by seine on 1 September 1976 in the vicinity of TMS.

Station	TM-AQF-1A2	TM-AQF-1A1	TM-AQF-1A2	TM-AQF-1A1	TM-AQF-1A2	Total	Catch
Time	1308	1355	1413	1445	1500		
Air Temp. (C)	24.0	26.0	25.5	26.5	25.5		
Water Temp. (C)	22.0	22.5	22.5	23.0	22.5		
Dissolved Oxygen (ppm)	8.3	8.1	7.6	7.7	8.0		
pH	8.5	8.5	8.4	8.5	8.5		
Secchi Disc (cm)	30	29	28	30	30		
River Stage (ft)	3.73	3.73	3.73	3.73	3.73		
Weather	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy		
No. of Spmn.	220	134	124	7	49	621	
No. of Spp.	8	12	11	7	4	20	
Cutlips minnow	-	-	1	-	-	1	0.2
River chub	-	-	1	-	-	1	0.2
Golden shiner	-	1	-	-	-	1	0.2
Comely shiner	-	1	2	-	-	3	0.5
Common shiner	-	7	-	-	-	7	1.1
Spottail shiner	13	104	24	12	3	156	25.1
Swallowtail shiner	8	-	-	-	-	8	1.3
Spotfin shiner	172	16	4	33	-	225	36.2
Bluntnose minnow	1	2	4	3	-	10	1.6
Creek chub	-	1	-	-	-	1	0.2
White sucker	-	4	1	1	-	6	1.0
Northern ho. sucker	2	-	-	-	-	2	0.3
Shorthead redhorse	2	-	1	3	-	6	1.0
White catfish	-	-	1	-	-	1	0.2
Channel catfish	-	-	58	28	41	127	20.5
Pumpkinseed	-	1	-	-	-	1	0.2
Bluegill	-	-	-	-	1	1	0.2
Smallmouth bass	1	3	-	-	-	4	0.6
Black crappie	-	4	-	-	-	4	0.6
Tessellated darter	21	10	7	14	4	56	9.0

Table 2.2-41

Fishes taken by seine on 13 September 1976 in the vicinity of TMS.

Station	TM-AQF-1A2	TM-AQF-1A1	TM-AQF-1A2	TM-AQF-1A1	TM-AQF-1A2	Total	Catch
Time	1432	1345	1326	1253	1240		
Air Temp. (C)	27.5	28.5	26.0	26.0	25.0		
Water Temp. (C)	21.5	21.5	21.0	22.0	22.5		
Dissolved Oxygen (ppm)	7.6	7.7	7.5	7.6	7.8		
pH	8.5	8.4	8.4	8.6	8.7		
Secchi Disc (cm)	36	46	33	53	53		
River Stage (ft)	3.48	3.48	3.48	3.48	3.48		
Weather	Clear	Clear	Clear	Clear	Clear		
No. of Spmn.	114	54	59	21	77	325	
No. of Spp.	8	10	6	5	5	14	
Comely shiner	8	1	1	-	-	10	3.1
Common shiner	-	1	-	-	-	1	0.3
Spottail shiner	16	26	18	5	-	65	20.0
Spotfin shiner	49	-	19	-	1	69	
Bluntnose minnow	13	3	3	-	-	19	3.0
Shorthead redhorse	-	2	-	-	-	2	0.6
Channel catfish	1	1	8	7	72	89	27.4
Rock bass	1	-	-	-	-	1	0.3
Redbreast sunfish	-	1	-	1	-	2	0.6
Pumpkinseed	-	-	-	-	1	1	0.3
Bluegill	-	1	-	-	1	2	0.6
Smallmouth bass	1	-	-	2	-	3	0.9
Tessellated darter	25	17	10	6	2	60	18.5
Banded darter	-	1	-	-	-	1	0.3

1565 092

Table 2.2-42

Fishes taken by seine on 29 September 1976 in the vicinity of TNINS.

Station	TM-AQF-1A2	TM-AQF-1A41	TM-AQF-10A2	TM-AQF-2A1	TM-AQF-2A3	Total	Catch
Time	1330	1415	1433	1506	1520		
Air Temp. (C)	19.5	20.0	19.5	18.5	20.0		
Water Temp. (C)	16.0	17.5	16.5	17.0	17.5		
Dissolved Oxygen (ppm)	9.6	10.0	9.8	9.8	10.0		
pH	8.3	8.2	8.1	8.1	8.1		
Secchi Disc (cm)	30	30	30	36	33		
River Stage (ft)	3.85	3.85	3.85	3.85	3.85		
Weather	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy		
No. of Spmn.	539	27	231	45	247	1104	
No. of Spp.	7	6	11	5	6	15	
Comely shiner	1	-	-	-	-	1	0.1
Spottail shiner	3	9	8	12	-	32	2.9
Swallowtail shiner	2	-	1	-	-	3	0.3
Spotfin shiner	510	-	178	-	221	909	82.3
Bluntnose minnow	2	1	10	6	-	19	1.7
Creek chub	-	1	-	-	-	1	0.1
Fallfish	-	-	2	-	-	2	0.2
Shorthead redhorse	-	-	1	-	1	2	0.2
Channel catfish	-	1	4	2	1	8	0.7
Redbreast sunfish	-	-	1	-	-	1	0.1
Pumpkinseed	-	-	-	-	1	1	0.1
Bluegill	-	-	1	-	-	1	0.1
Smallmouth bass	-	-	1	1	1	3	0.3
Tessellated darter	11	14	44	19	22	110	10.0
Banded darter	10	1	-	-	-	11	1.0

POOR ORIGINAL

Table 2.2-43

Fishes taken by seine on 19 October 1976 in the vicinity of TNINS.

Station	TM-AQF-1A2	TM-AQF-1A41	TM-AQF-10A2	TM-AQF-2A1	TM-AQF-2A3	Total	Catch
Time	1518	1437	1418	1347	1328		
Air Temp. (C)	12.0	12.0	12.5	12.0	10.0		
Water Temp. (C)	11.5	11.0	11.5	11.0	11.0		
Dissolved Oxygen (ppm)	12.4	12.7	12.7	12.5	12.4		
pH	8.1	8.1	8.1	8.1	8.2		
Secchi Disc (cm)	30	33	36	48	46		
River Stage (ft)	5.34	5.34	5.34	5.34	5.34		
Weather	Clear	Partly Cloudy	Clear	Clear	Partly Cloudy		
No. of Spmn.	49	78	265	14	90	506	
No. of Spp.	6	9	8	5	4	14	
Comely shiner	1	-	-	-	-	1	0.2
Common shiner	-	2	-	-	-	2	0.4
Spottail shiner	6	21	22	3	-	52	10.3
Swallowtail shiner	2	2	10	-	5	19	3.8
Rosyface shiner	-	1	-	-	-	1	0.2
Spotfin shiner	6	3	181	1	57	248	49.0
Bluntnose minnow	11	13	18	-	3	45	8.9
Pathead minnow	-	-	1	-	-	1	0.2
Fallfish	-	-	1	-	-	1	0.2
Rock bass	-	-	-	1	-	1	0.2
Pumpkinseed	-	-	-	1	-	1	0.2
Smallmouth bass	-	2	-	1	-	3	0.6
Tessellated darter	23	31	31	17	25	127	25.1
Banded darter	-	3	1	-	-	4	0.8

1565 093

Table 2.2-44

Fishes taken by seine on 25 October 1976 in the vicinity of TMS.

Station	TM-AQF-142	TM-AQF-14A1	TM-AQF-14A2	TM-AQF-14A1	TM-AQF-14A3	Total	Count
Time	0938	1015	1030	1105	1125		
Air Temp. (C)	9.0	9.5	9.5	9.5	10.0		
Water Temp. (C)	7.5	7.5	8.0	7.5	7.5		
Dissolved Oxygen (ppm)	12.6	12.5	12.6	12.7	12.6		
pH	8.5	8.5	8.3	8.4	8.4		
Secchi disc (cm)	18	13	13	13	13		
River Stage (ft)	7.94	7.94	7.94	7.94	7.94		
Weather	Fog	Fog	Fog	Fog	Fog		
No. of spmn.	178	72	67	91	123	531	
No. of spp.	10	10	7	6	6	13	
Common shiner	2	2	8	2	-	14	2.6
Common shiner	1	-	-	-	-	1	0.2
Spottail shiner	25	47	15	31	94	212	39.9
Swallowtail shiner	8	2	5	4	2	21	4.0
Rosyface shiner	2	3	-	-	-	5	0.9
Spotfin shiner	122	3	28	20	22	205	38.6
Bluntnose minnow	1	3	2	-	2	8	1.5
Quillback	-	-	-	-	1	1	0.2
Redbreast sunfish	-	1	-	1	-	2	0.4
Pumpkinseed	-	1	-	-	-	1	0.2
Bluegill	1	-	-	-	-	1	0.2
Smallmouth bass	1	1	2	-	-	4	0.8
Tessellated darter	5	9	7	13	2	36	6.8

POOR ORIGINAL

1565 094

Table 2.2-45

46

Summary of fishes taken at seine station TM-AQF-1A2 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	Catch
River chub	-	-	-	-	-	1	-	-	1	+
Comely shiner	-	-	-	-	4	3	4	3	14	0.7
Common shiner	-	-	-	-	5	8	-	1	14	0.5
Spottail shiner	-	-	-	407	813	24	32	31	1307	49.4
Swallowtail shiner	4	7	4	2	-	2	10	10	36	1.4
Rosyface shiner	-	1	-	-	-	-	-	2	3	0.1
Spotfin shiner	28	6	1	28	78	13	241	138	1023	38.7
Bluntnose minnow	9	-	-	-	2	5	16	12	44	1.7
Blacknose dace	1	-	-	-	2	-	-	-	3	0.1
Longnose dace	-	-	-	1	-	-	-	-	1	+
Fallfish	-	-	-	-	-	1	-	-	1	+
White sucker	-	-	-	-	4	-	-	-	4	0.2
Northern hog sucker	-	-	-	-	-	-	2	-	2	0.1
Shorthead redhorse	-	-	-	-	1	-	2	-	3	0.1
White sucker/Shorthead redhorse	-	-	-	12	-	-	-	-	12	0.5
Channel catfish	-	-	-	-	-	-	1	2	3	+
Rock bass	-	-	-	-	1	-	1	-	2	0.1
Pumpkinseed	-	-	-	1	1	-	-	-	2	0.1
Bluegill	-	-	-	-	-	-	-	1	1	+
Smallmouth bass	-	-	1	1	11	2	2	1	18	0.7
Black crappie	-	-	-	-	-	1	-	-	1	+
Tessellated darter	4	4	1	17	2	20	57	28	133	5.0
Banded darter	3	-	-	-	-	-	12	-	15	0.5
Shield darter	-	-	-	1	-	-	-	-	1	+
No. of Spmn.	49	18	4	470	924	80	573	227	2635	
No. of Spp.	6	4	4	9	12	11	12	10	29	
No. of Coll.	3	2	2	2	2	2	3	2	18	
n/Coll	16.33	9.00	2.00	235.00	462.00	40.00	291.00	113.50	146.94	
Diversity Index	1.86	1.77	2.00	0.84	0.72	2.71	1.35	1.87	1.75	

+ = less than 0.05%.

Table 2.2-46

Summary of fishes taken at seine station TM-AQF-16A1 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	Catch
Golden shiner	-	-	-	-	-	-	1	-	1	+
Comely shiner	5	-	-	3	26	2	2	2	40	1.8
Common shiner	-	-	-	-	2	9	8	2	21	0.9
Spottail shiner	3	-	-	600	483	168	139	68	1461	62.4
Swallowtail shiner	4	1	-	-	-	-	-	4	9	0.4
Rosyface shiner	-	-	-	-	-	-	-	4	4	0.2
Spotfin shiner	27	5	-	16	6	23	16	6	99	4.4
Bluntnose minnow	4	-	-	-	2	10	6	16	38	1.7
Fathead minnow	1	-	-	-	-	-	-	-	1	+
Blacknose dace	1	-	-	1	-	-	-	-	2	0.1
Longnose dace	-	-	-	1	-	-	-	-	1	+
Creek chub	-	-	-	2	-	-	2	-	4	0.2
Fallfish	-	-	-	-	4	2	-	-	6	0.3
Quillback	-	-	-	-	1	-	-	-	1	+
White sucker	-	-	-	1	3	1	4	-	9	0.4
Northern hog sucker	-	-	-	-	1	-	-	-	1	+
Shorthead redhorse	-	-	-	16	6	6	2	-	30	1.3
White sucker/Shorthead redhorse	-	-	-	324	10	-	-	-	334	14.7
Channel catfish	-	-	-	-	-	-	2	-	2	0.1
Rock bass	-	-	-	-	1	4	-	-	5	0.2
Redbreast sunfish	1	-	-	-	-	1	1	1	4	0.2
Pumpkinseed	-	-	-	-	-	-	1	1	2	0.1
Bluegill	-	-	-	-	-	1	1	-	2	0.1
Smallmouth bass	-	-	1	-	12	3	3	3	22	1.0
Largemouth bass	-	-	-	-	1	-	-	-	1	+
White crappie	-	-	-	-	-	1	-	-	1	+
Black crappie	-	-	-	-	-	1	4	-	5	0.2
Tessellated darter	10	2	-	11	18	17	41	40	139	6.1
Banded darter	-	-	-	-	3	-	2	3	8	0.4
Shield darter	-	-	-	16	-	-	-	-	16	0.7
No. of Spmn.	56	8	1	991	579	249	235	150	2269	
No. of Spp.	9	3	1	10	15	15	17	12	29	
No. of Coll.	3	2	2	2	2	2	3	2	18	
n/Coll	18.67	4.00	0.50	495.50	289.50	124.50	78.33	75.00	126.06	
Diversity Index	2.34	1.30	-	1.40	1.18	1.90	2.15	2.32	1.99	

+ = less than 0.05%.

POOR ORIGINAL

1565 095

Table 2.2-47

Summary of fishes taken at seine station TM-AQF-10A2 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	Catch
Carp	-	-	-	-	1	-	-	-	1	0.1
Cutlips minnow	-	-	-	-	-	-	1	-	1	0.1
River chub	-	-	-	-	-	1	1	-	2	0.1
Comely shiner	-	-	-	1	2	-	3	8	14	0.7
Spottail shiner	25	22	3	449	213	99	50	37	918	45.9
Swallowtail shiner	3	2	-	-	-	-	1	15	21	1.1
Rosyface shiner	4	-	-	-	-	-	-	-	4	0.2
Spotfin shiner	32	10	-	12	3	3	201	209	490	24.5
Bluntnose minnow	13	3	1	4	1	3	17	20	62	3.1
Fathead minnow	-	-	-	-	-	-	-	1	1	0.1
Longnose dace	-	-	-	2	-	-	-	-	2	0.1
Creek chub	1	-	-	9	1	-	-	-	11	0.6
Fallfish	-	-	-	-	4	1	2	1	8	0.4
White sucker	-	-	-	1	10	3	1	-	15	0.8
Shorthead redhorse	-	-	-	-	23	12	2	-	37	1.9
White sucker/Shorthead redhorse	-	-	-	133	6	-	-	-	139	7.0
White catfish	-	-	-	-	-	-	1	-	1	0.1
Channel catfish	1	-	-	-	5	-	70	-	76	3.8
Redbreast sunfish	2	-	-	2	-	-	1	-	5	0.3
Pumpkinseed	-	-	-	-	-	1	-	-	1	0.1
Bluegill	-	-	-	-	-	-	1	-	1	0.1
Smallmouth bass	-	1	1	-	5	11	1	2	21	1.1
White crappie	-	-	-	-	1	-	-	-	1	0.1
Tessellated darter	5	2	1	8	8	39	61	38	162	8.2
Dotted darter	3	1	-	-	-	-	-	1	5	0.3
Shield darter	-	-	-	1	-	-	-	-	1	0.1
No. of Spmn.	89	61	6	622	303	173	414	332	2000	
No. of Spp.	10	7	4	11	13	10	16	10	25	
No. of Coll.	3	2	2	2	2	2	3	2	18	
n/Coll.	29.67	30.50	3.00	311.00	151.50	86.50	138.00	166.00	111.11	
Diversity Index	2.48	1.77	1.79	1.24	1.49	1.90	2.20	1.53	2.45	

Table 2.2-48

Summary of fishes taken at seine station TM-AQF-9A1 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	Catch
Comely shiner	-	-	-	-	-	-	-	2	2	0.2
Spottail shiner	1	-	-	317	113	48	29	34	542	53.9
Swallowtail shiner	3	-	NO	-	-	-	-	4	7	0.7
Spotfin shiner	26	1	-	5	4	5	33	21	95	9.4
Bluntnose minnow	3	-	-	-	2	4	9	-	18	1.8
Creek chub	-	-	-	6	-	-	-	-	6	0.6
Fallfish	-	-	-	-	9	1	-	-	10	1.0
White sucker	-	-	-	-	12	4	1	-	17	1.7
Northern hog sucker	-	-	FISH	-	-	1	-	-	1	0.1
Shorthead redhorse	-	-	-	-	4	8	3	-	15	1.5
White sucker/Shorthead redhorse	-	-	-	88	-	-	-	-	88	8.7
Channel catfish	-	-	-	-	-	3	37	-	40	4.0
Rock bass	-	-	-	-	-	-	-	1	1	0.1
Redbreast sunfish	-	-	-	-	1	1	1	1	4	0.4
Pumpkinseed	-	1	TAKEN	-	-	-	-	1	2	0.2
Bluegill	-	-	-	1	-	2	-	-	3	0.3
Smallmouth bass	-	-	-	1	4	2	3	1	11	1.1
White crappie	-	-	-	-	3	4	-	-	7	0.7
Tessellated darter	5	-	-	2	10	29	39	50	135	13.4
Shield darter	-	-	-	2	-	-	-	-	2	0.2
No. of Spmn.	38	2	-	422	162	112	155	115	1006	
No. of Spp.	5	2	-	8	10	13	9	9	19	
No. of Coll.	3	2	2	2	2	2	3	2	18	
n/Coll.	12.67	1.00	-	211.00	81.00	56.00	51.67	57.50	55.89	
Diversity Index	1.48	1.00	-	1.06	1.75	2.55	2.47	2.00	2.39	

Table 2.2-49

Summary of fishes taken at seine station TM-AQF-983 during 1976.

Month	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Total	Rich
Comely shiner	-	-	-	-	-	1	-	-	1	-
Common shiner	1	-	-	-	-	1	-	-	2	0.1
Spottail shiner	1	2	1	486	212	19	3	94	1248	52.7
Swallowtail shiner	13	1	-	-	-	-	-	7	21	0.9
Spotfin shiner	14	5	-	8	-	12	222	79	365	14.8
Bluntnose minnow	1	-	-	-	1	23	-	5	30	1.2
Longnose dace	-	-	-	7	-	-	-	-	7	0.3
Creek chub	-	-	4	24	-	-	-	-	28	1.1
Fallfish	-	-	-	-	3	-	-	-	3	0.1
Quillback	-	-	-	-	-	-	-	1	1	-
White sucker	-	-	-	-	116	1	-	-	117	4.7
Shorthead redhorse	-	-	-	-	8	-	1	-	9	0.4
White sucker/Shorthead redhorse	-	-	30	100	-	-	-	-	130	5.1
Yellow bullhead	-	-	-	-	-	2	-	-	2	0.1
Brown bullhead	-	-	-	-	7	2	-	-	9	0.4
Channel catfish	-	-	-	-	2	256	114	-	372	14.7
Redbreast sunfish	1	-	-	-	-	-	-	-	1	-
Pumpkinseed	-	-	-	-	-	-	2	-	2	0.1
Bluegill	-	-	-	-	-	-	2	-	2	0.1
Smallmouth bass	-	-	1	-	6	2	1	-	10	0.4
White crappie	-	-	-	-	-	14	-	-	14	0.7
Tessellated darter	1	-	1	1	1	37	24	-	45	1.8
No. of Spmn.	57	8	17	116	374	375	373	213	2058	
No. of Spp.	7	3	5	6	9	12	8	6	24	
No. of Coll.	3	2	2	2	2	2	3	2	18	
n/Coll.	146.00	4.00	18.50	593.00	144.50	187.50	124.33	106.50	142.11	
Diversity Index	1.17	1.33	1.01	0.71	1.21	1.74	1.23	1.29	2.05	

- = less than 0.05.

Table 2.2-50

Summary of fishes taken at seine stations during March through October 1976.

Station	TM-AQF-132	TM-AQF-133	TM-AQF-134	TM-AQF-135	TM-AQF-136	Total	Rich
Carp	-	-	1	-	-	1	-
Cutlips minnow	-	-	1	-	-	1	-
River chub	1	-	2	-	-	3	-
Golden shiner	-	1	-	-	-	1	-
Comely shiner	19	40	14	2	1	76	0.7
Common shiner	14	21	-	-	2	37	0.4
Spottail shiner	1307	1461	918	542	1298	5526	52.7
Swallowtail shiner	36	9	21	7	21	94	0.9
Rosyface shiner	3	4	-	-	-	11	0.1
Spotfin shiner	1023	99	497	95	165	2072	19.8
Bluntnose minnow	14	18	62	18	30	142	1.4
Fathead minnow	-	1	1	-	-	2	-
Blacknose dace	3	2	-	-	-	5	-
Longnose dace	1	1	2	-	7	11	0.1
Creek chub	-	4	11	6	28	49	0.5
Fallfish	1	6	8	10	3	28	0.3
Quillback	-	1	-	-	1	2	-
White sucker	4	9	13	17	117	152	1.7
Northern hog sucker	2	1	-	1	-	4	-
Shorthead redhorse	3	30	37	15	4	89	0.8
White sucker/Shorthead redhorse	12	334	139	88	130	703	6.7
White catfish	-	-	1	-	-	1	-
Yellow bullhead	-	-	-	-	2	2	-
Brown bullhead	-	-	-	-	9	9	0.1
Channel catfish	1	2	76	40	377	496	4.7
Rock bass	2	5	-	1	-	8	0.1
Redbreast sunfish	-	4	5	4	1	14	0.1
Pumpkinseed	2	2	1	2	2	9	0.1
Bluegill	1	2	1	3	2	9	0.1
Smallmouth bass	18	22	21	11	10	82	0.8
Largemouth bass	-	1	-	-	-	1	-
White crappie	-	1	4	7	19	28	0.3
Black crappie	1	5	-	-	-	6	0.1
Tessellated darter	133	139	162	135	109	678	6.5
Banded darter	13	8	5	-	-	26	0.2
Shield darter	1	15	1	2	-	20	0.2
No. of Spmn.	2645	2269	2000	1006	2558	10478	
No. of Spp.	23	29	25	19	21	35	
No. of Coll.	18	18	18	18	18	90	
n/Coll.	146.94	126.06	111.11	55.89	142.11	116.52	
Diversity Index	1.25	1.99	2.25	2.39	2.35	2.36	

- = less than 0.05.

POOR ORIGINAL

Table 2.2-52

Condition factors and reproduction status (Y = young, J = juvenile, A = adult) of channel catfish taken by trammel during March through October 1976 downstream (D) and upstream (U) from the T1000 discharge. \bar{K} (number in parentheses equals specimens used in calculation).

Fork Length (5.00 to 375)	March		April		May		June		July		August		September		October	
	D	U	D	U	D	U	D	U	D	U	D	U	D	U	D	U
46-50	-	0.80(1J)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
61-65	-	0.77(1J)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
66-70	0.87(1J)	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
86-90	-	0.82(1J)	-	-	0.82(1J)	-	-	-	-	-	-	-	-	-	-	-
96-100	-	-	-	-	-	-	-	-	1.50(1J)	-	-	-	-	-	-	-
111-115	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
116-120	1.10(1J)	1.16(1J)	-	-	-	-	1.12(1J)	-	-	-	1.18(1J)	-	-	-	-	-
121-125	1.18(1J)	1.40(1J)	-	-	-	-	-	-	0.98(1J)	-	-	-	-	-	-	-
126-130	-	0.92(1J)	-	-	-	-	1.02(1J)	-	1.21(1J)	-	-	-	-	-	1.27(1J)	-
131-135	1.27(1J)	-	1.09(1J)	-	-	-	-	-	-	-	1.60(1J)	-	1.02(1J)	-	-	-
136-140	1.06(1J)	-	0.81(1J)	-	-	-	-	-	1.16(1J)	1.14(1J)	1.15(1J)	-	1.02(1J)	-	-	-
141-145	0.97(1J)	-	1.09(1J)	-	-	-	-	-	0.97(1J)	-	0.98(1J)	-	1.02(1J)	-	-	-
146-150	-	0.99(1J)	-	-	-	-	-	-	-	-	-	-	1.33(1J)	1.16(1J)	-	-
151-155	-	1.01(1J)	-	-	-	-	-	-	0.89(1J)	-	-	-	1.02(1J)	-	-	-
156-160	-	1.06(1J)	-	-	-	-	-	-	1.17(1J)	-	-	-	1.02(1J)	-	-	-
161-165	0.99(1J)	-	-	-	0.98(1J)	-	-	-	-	-	-	-	1.02(1J)	-	-	-
166-170	0.96(1J)	1.00(1J)	-	-	-	-	1.29(1J)	-	-	-	-	-	1.33(1J)	-	-	-
171-175	-	0.95(1J)	-	-	-	-	1.06(1J)	-	-	-	-	-	1.14(1J)	-	-	-
176-180	-	-	-	-	-	-	1.32(1J)	-	-	-	-	-	-	-	-	-
181-185	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
186-190	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
191-195	-	-	-	-	1.20(1A)	1.51(1A)	1.17(1A)	-	-	-	-	-	1.17(1A)	-	1.10(1J)	-
201-205	-	-	-	-	-	-	1.12(1A)	-	-	-	-	-	-	-	1.14(1A)	-
216-220	-	-	-	-	-	-	1.16(1A)	-	-	-	-	-	-	-	-	-
221-225	-	-	-	-	1.12(1A)	-	-	-	-	-	-	-	-	-	1.16(1A)	-
226-230	-	-	-	-	1.40(1A)	-	1.26(1A)	-	-	-	-	-	-	-	1.07(1A)	-
231-235	-	-	-	-	-	-	1.00(1A)	-	-	-	-	-	-	-	-	-
236-240	-	1.18(1A)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
241-245	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
246-250	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
251-255	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
256-260	-	-	1.37(2A)	-	-	-	1.33(1A)	-	-	-	-	-	-	-	-	-
261-265	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
266-270	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
271-275	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
276-280	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
281-285	-	-	-	-	1.27(1A)	-	-	-	1.31(1A)	-	-	-	-	-	1.11(1A)	-
286-290	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
291-295	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
296-300	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
301-305	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
306-310	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
311-315	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
316-320	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1565 099

Table 2.2-53

condition factors and reproductive status (V = young, J = juvenile, A = adults) of rock bass taken by trapnet during March through October 1966 westward of and upstream of the TMD. Discharge (number in parentheses equals specimens used in calculations)

Rock Length mm (KUMH)	March		April		May		June		July		August		Sept/Oct		KUMH
	V	J	V	J	V	J	V	J	V	J	V	J	V	J	
61-65	-	-	-	2.91(11)	-	-	-	-	-	-	-	-	-	-	-
96-100	-	-	-	-	-	-	-	-	-	-	-	2.24(11)	-	-	-
101-105	-	-	-	1.12(11)	-	-	-	-	-	-	-	-	-	-	-
106-110	1.95(12)	-	-	1.50(11)	-	-	-	-	-	-	-	-	1.80(11)	-	-
111-115	-	-	-	-	-	2.50(11)	-	-	-	-	-	2.17(23)	-	-	-
116-120	-	-	-	1.97(22)	-	-	-	-	-	-	-	1.76(11)	-	-	1.41(11)
121-125	-	-	-	2.00(11)	-	2.25(11)	-	-	-	-	-	-	1.95(11)	-	-
126-130	-	-	1.78(11)	-	1.73(11)	2.41(11)	2.73(11)	-	-	-	-	-	1.89(11)	2.09(11)	1.81(11)
131-135	-	-	-	-	-	-	2.32(1A)	2.19(1A)	-	-	-	-	-	-	-
136-140	-	-	-	-	-	-	2.44(1A)	-	-	-	-	-	-	2.03(1A)	-
141-145	-	-	1.71(1A)	-	2.30(1A)	-	2.71(2A)	2.31(1A)	2.89(1A)	-	-	1.81(1A)	-	-	2.04(1A)
146-150	2.07(1A)	-	-	-	-	-	2.15(2A)	2.11(1A)	-	-	-	-	-	-	2.16(1A)
151-155	-	-	-	-	-	-	-	-	2.81(1A)	-	-	-	-	-	-
156-160	-	-	-	-	2.11(2A)	-	2.37(2A)	-	2.55(2A)	-	-	2.04(1A)	1.96(1A)	-	-
161-165	-	-	1.60(1A)	-	-	-	2.17(2A)	-	-	-	-	-	1.17(1A)	-	2.04(1A)
166-170	-	-	1.87(1A)	-	-	-	2.25(1A)	-	-	-	2.32(1A)	-	2.24(1A)	2.04(1A)	-
171-175	-	-	2.09(1A)	-	2.31(2A)	-	2.25(1A)	-	-	-	-	-	2.11(2A)	-	-
176-180	-	-	-	-	-	-	-	-	2.34(1A)	-	-	-	2.14(1A)	-	-
181-185	-	-	2.05(1A)	-	2.29(1A)	2.57(1A)	2.70(2A)	-	2.40(1A)	-	-	-	2.14(1A)	-	2.15
191-195	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
196-200	-	-	2.24(1A)	-	-	2.60(1A)	-	-	-	-	-	-	2.51(1A)	-	-
206-210	-	-	2.54(1A)	-	2.32(2A)	-	-	-	-	-	-	-	-	-	-
211-215	-	-	2.39(1A)	2.92(1A)	-	-	-	-	-	-	-	-	-	-	-
216-220	-	-	-	-	-	-	-	-	-	-	-	-	1.50(1A)	-	-
221-225	2.17(1A)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
231-235	-	-	-	-	-	-	-	-	-	-	-	-	2.31(1A)	-	-
236-250	-	-	2.58(1A)	-	-	2.35(1A)	-	-	-	-	-	-	-	-	-

POOR ORIGINAL

Table 2.2-55

Condition factors and reproductive status (V = young, J = juvenile, A = adult) of pumpkinseed taken by trapnet during March through October 1976 downstream (S) and upstream (U) from the
 IRLS discharge (number in parentheses equals specimens used in calculation)

Iris length 15.33-15.33(1)	March		April		May		June		July		August		September		Total
	V	J	V	J	V	J	V	J	V	J	V	J	V	J	
56-60	0.93(1J)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
76-80	2.13(1J)	-	-	-	-	-	-	-	-	-	-	-	-	-	-
81-85	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
86-90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
91-95	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
96-100	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
101-105	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
106-110	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
111-115	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
116-120	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
121-125	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
126-130	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
131-135	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
136-140	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
141-145	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
146-150	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
151-155	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
156-160	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
161-165	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
166-170	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
171-175	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1565 101

Table 2.2-55

Condition factors and reproductive status (Y = young, I = juvenile, A = adult) of white crappie taken by trawls during March through October 1975 downstream of and upstream of the
 21155 Discharge (number in parentheses equals specimens used in calculation).

Forch month	March	April	May	June	July	August	September	October
15-48-67-122	1	2	3	4	5	6	7	8
66-70	-	-	-	-	-	-	1.46(13)	-
76-80	-	-	-	-	-	-	0.79(13)	-
86-90	-	1.51(11)	-	-	-	-	-	-
91-95	-	30	-	-	-	-	1.17(13)	2
96-100	-	-	-	-	-	-	1.79(13)	-
136-140	-	-	1.75(11)	-	-	-	-	-
161-165	-	-	-	-	-	-	1.44(13)	-
166-170	-	-	-	-	-	-	1.57(13)	-
176-180	-	-	-	-	-	-	1.99(13)	-
181-185	-	118	-	-	-	1.53(13)	-	110
186-190	-	-	-	1.56(13)	-	-	-	-
196-200	-	-	-	-	1.36(13)	-	-	-
206-210	-	-	-	-	-	-	1.45(13)	-
211-215	-	1.52(13)	-	-	-	-	-	-
226-230	-	-	-	-	-	-	1.51(13)	-
231-235	-	10018	-	-	-	-	1.51(13)	10018
261-265	-	-	-	-	-	-	1.56(13)	-
266-270	-	1.65(13)	-	-	-	-	-	-

POOR ORIGINAL

1565 102

Table 2.2-5b

Condition factors and reproductive status (Y = young, J = juvenile, A = adult) of black sturgeon taken by trawler during March through October 1976 downstream (D) and upstream (U) from the PINE discharge (number in parentheses equals specimens used in calculation)

Fork length (mm, rounded)	March		April		May		June		July		August		September		October	
	D	U	D	U	D	U	D	U	D	U	D	U	D	U	D	U
66-70	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
96-100	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
110-120	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
121-125	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
126-130	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
136-140	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
141-145	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
151-155	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
156-160	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
161-165	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
166-170	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
171-175	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
176-180	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
181-185	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
186-190	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
191-195	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
196-200	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
201-205	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
206-210	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
211-215	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
216-220	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
221-225	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
226-230	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
231-235	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
236-240	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
241-245	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
261-265	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1565 103

POOR ORIGINAL

Table 2.2-57

Condition factors and reproductive status (Y = young, J = juvenile, A = adult) of speckled shiner taken by seine during May through October 1976 at Washington, D.C. (parentheses show the FWS discharge number in parentheses equals specimens used in calculation)

Core Length 45 mm (K0042)	March		April		May		June		July		August		September		October	
	F	Y	F	Y	F	Y	F	Y	F	Y	F	Y	F	Y	F	Y
6-10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
11-15	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
16-20	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
21-25	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
26-30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
31-35	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
36-40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
41-45	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
46-50	0.904533	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
51-55	1.061119	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
56-60	1.061719	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
61-65	0.951683	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
66-70	0.901983	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
71-75	0.901983	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
76-80	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
81-85	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

POOR ORIGINAL

POOR ORIGINAL

Table 2. (cont.)

Length frequency and mean weights per 5 m group of fishes taken by trammel net in 100 m ² 10% domestic and offshore traps for the 1971-1972 season										Length frequency and mean weights per 5 m group of fishes taken by trammel net in 100 m ² 10% domestic and offshore traps for the 1973-1974 season																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																				
Species	Length (mm)	No.	Mean weight (g)	No.	Mean weight (g)	Species	Length (mm)	No.	Mean weight (g)	Species	Length (mm)	No.	Mean weight (g)	Species	Length (mm)	No.	Mean weight (g)																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
pettail shiner	81-85	1	26.7	1	26.7	white sucker	101-105	1	11.0	channel catfish	46-50	1	1.0	channel catfish	46-50	1	1.0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
	86-90	2	18.0	2	18.0		91-95	1	10.0		1	10.0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
	91-95	1	10.0	1	10.0																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									

Table 2.2-62

Length frequency and mean weights per 5 m group of fishes taken by trapnet on 27-31 March 1976 downstream and upstream from the TAMS discharge.

Species	Fork length (5 m groups)	No.	Total wt.	No.	Total wt.
CARP	266-370	80 FISH TAKEN	66.1	1000*	7.0
Spot-tail shiner	86-90	1	890.0	80 FISH TAKEN	510.0
White catfish	371-375	1	890.0	80 FISH TAKEN	510.0
Green bullhead	311-315	1	890.0	80 FISH TAKEN	510.0
Channel catfish	86-90	1	890.0	80 FISH TAKEN	510.0
Margined madtom	116-120	1	23	20	20.0
Rock bass	121-125	1	23	20	20.0
	126-130	1	23	20	20.0
	146-150	1	23	20	20.0
	151-155	1	23	20	20.0
	166-170	1	23	20	20.0
	26-100	1	23	20	20.0
	106-110	1	23	20	20.0
Pumpkinseed	166-170	1	23	20	20.0
	171-175	1	23	20	20.0
	181-185	1	23	20	20.0
	196-200	1	23	20	20.0
	206-210	1	23	20	20.0
	211-215	1	23	20	20.0
	216-220	1	23	20	20.0
	221-225	1	23	20	20.0
	226-230	1	23	20	20.0
	231-235	1	23	20	20.0
	236-240	1	23	20	20.0
	241-245	1	23	20	20.0
	246-250	1	23	20	20.0
	251-255	1	23	20	20.0
	256-260	1	23	20	20.0
	261-265	1	23	20	20.0
	266-270	1	23	20	20.0
	271-275	1	23	20	20.0
	276-280	1	23	20	20.0
	281-285	1	23	20	20.0
	286-290	1	23	20	20.0
	291-295	1	23	20	20.0
	296-300	1	23	20	20.0
	301-305	1	23	20	20.0
	306-310	1	23	20	20.0
	311-315	1	23	20	20.0
	316-320	1	23	20	20.0
	321-325	1	23	20	20.0
	326-330	1	23	20	20.0
	331-335	1	23	20	20.0
	336-340	1	23	20	20.0
	341-345	1	23	20	20.0
	346-350	1	23	20	20.0
	351-355	1	23	20	20.0
	356-360	1	23	20	20.0
	361-365	1	23	20	20.0
	366-370	1	23	20	20.0
	371-375	1	23	20	20.0
	376-380	1	23	20	20.0
	381-385	1	23	20	20.0
	386-390	1	23	20	20.0
	391-395	1	23	20	20.0
	396-400	1	23	20	20.0
	401-405	1	23	20	20.0
	406-410	1	23	20	20.0
	411-415	1	23	20	20.0
	416-420	1	23	20	20.0
	421-425	1	23	20	20.0
	426-430	1	23	20	20.0
	431-435	1	23	20	20.0
	436-440	1	23	20	20.0
	441-445	1	23	20	20.0
	446-450	1	23	20	20.0
	451-455	1	23	20	20.0
	456-460	1	23	20	20.0
	461-465	1	23	20	20.0
	466-470	1	23	20	20.0
	471-475	1	23	20	20.0
	476-480	1	23	20	20.0
	481-485	1	23	20	20.0
	486-490	1	23	20	20.0
	491-495	1	23	20	20.0
	496-500	1	23	20	20.0
	501-505	1	23	20	20.0
	506-510	1	23	20	20.0
	511-515	1	23	20	20.0
	516-520	1	23	20	20.0
	521-525	1	23	20	20.0
	526-530	1	23	20	20.0
	531-535	1	23	20	20.0
	536-540	1	23	20	20.0
	541-545	1	23	20	20.0
	546-550	1	23	20	20.0
	551-555	1	23	20	20.0
	556-560	1	23	20	20.0
	561-565	1	23	20	20.0
	566-570	1	23	20	20.0
	571-575	1	23	20	20.0
	576-580	1	23	20	20.0
	581-585	1	23	20	20.0
	586-590	1	23	20	20.0
	591-595	1	23	20	20.0
	596-600	1	23	20	20.0
	601-605	1	23	20	20.0
	606-610	1	23	20	20.0
	611-615	1	23	20	20.0
	616-620	1	23	20	20.0
	621-625	1	23	20	20.0
	626-630	1	23	20	20.0
	631-635	1	23	20	20.0
	636-640	1	23	20	20.0
	641-645	1	23	20	20.0
	646-650	1	23	20	20.0
	651-655	1	23	20	20.0
	656-660	1	23	20	20.0
	661-665	1	23	20	20.0
	666-670	1	23	20	20.0
	671-675	1	23	20	20.0
	676-680	1	23	20	20.0
	681-685	1	23	20	20.0
	686-690	1	23	20	20.0
	691-695	1	23	20	20.0
	696-700	1	23	20	20.0
	701-705	1	23	20	20.0
	706-710	1	23	20	20.0
	711-715	1	23	20	20.0
	716-720	1	23	20	20.0
	721-725	1	23	20	20.0
	726-730	1	23	20	20.0
	731-735	1	23	20	20.0
	736-740	1	23	20	20.0
	741-745	1	23	20	20.0
	746-750	1	23	20	20.0
	751-755	1	23	20	20.0
	756-760	1	23	20	20.0
	761-765	1	23	20	20.0
	766-770	1	23	20	20.0
	771-775	1	23	20	20.0
	776-780	1	23	20	20.0
	781-785	1	23	20	20.0
	786-790	1	23	20	20.0
	791-795	1	23	20	20.0
	796-800	1	23	20	20.0
	801-805	1	23	20	20.0
	806-810	1	23	20	20.0
	811-815	1	23	20	20.0
	816-820	1	23	20	20.0
	821-825	1	23	20	20.0
	826-830	1	23	20	20.0
	831-835	1	23	20	20.0
	836-840	1	23	20	20.0
	841-845	1	23	20	20.0
	846-850	1	23	20	20.0
	851-855	1	23	20	20.0
	856-860	1	23	20	20.0
	861-865	1	23	20	20.0
	866-870	1	23	20	20.0
	871-875	1	23	20	20.0
	876-880	1	23	20	20.0
	881-885	1	23	20	20.0
	886-890	1	23	20	20.0
	891-895	1	23	20	20.0
	896-900	1	23	20	20.0
	901-905	1	23	20	20.0
	906-910	1	23	20	20.0
	911-915	1	23	20	20.0
	916-920	1	23	20	20.0
	921-925	1	23	20	20.0
	926-930	1	23	20	20.0
	931-935	1	23	20	20.0
	936-940	1	23	20	20.0
	941-945	1	23	20	20.0
	946-950	1	23	20	20.0
	951-955	1	23	20	20.0
	956-960	1	23	20	20.0
	961-965	1	23	20	20.0
	966-970	1	23	20	20.0
	971-975	1	23	20	20.0
	976-980	1	23	20	20.0
	981-985	1	23	20	20.0
	986-990	1	23	20	20.0
	991-995	1	23	20	20.0
	996-1000	1	23	20	20.0

* weight greater than capacity of scale.

Table 2.2-63

Length frequency and mean weights per 5 m group of fishes taken by trapnet on 13-15 April 1976 downstream and upstream from the TAMS discharge.

Species	Fork length (5 m groups)	No.	Total wt.	No.	Total wt.
Spot-tail shiner	86-90	1	7.0	80 FISH TAKEN	510.0
	91-95	1	7.0	80 FISH TAKEN	510.0
	96-100	1	7.0	80 FISH TAKEN	510.0
	206-210	1	7.0	80 FISH TAKEN	510.0
Northern hog sucker	206-210	1	7.0	80 FISH TAKEN	510.0
White catfish	131-135	1	15.0	80 FISH TAKEN	510.0
Channel catfish	131-135	1	15.0	80 FISH TAKEN	510.0
	176-180	1	56.0	80 FISH TAKEN	510.0
	186-190	1	71.0	80 FISH TAKEN	510.0
	256-260	1	260.0	80 FISH TAKEN	510.0
	386-390	1	1000*	80 FISH TAKEN	510.0
Pack bass	496-500	1	1000*	80 FISH TAKEN	510.0
	116-120	1	2.0	80 FISH TAKEN	510.0
	166-170	1	92.0	80 FISH TAKEN	510.0
	171-175	1	112.0	80 FISH TAKEN	510.0
	181-185	1	129.0	80 FISH TAKEN	510.0
	196-200	1	176.0	80 FISH TAKEN	510.0
	206-210	1	235.0	80 FISH TAKEN	510.0
	211-215	1	232.0	80 FISH TAKEN	510.0
	216-220	1	356.0	80 FISH TAKEN	510.0
	221-225	1	19.0	80 FISH TAKEN	510.0
	226-230	1	112.0	80 FISH TAKEN	510.0
	231-235	1	85.0	80 FISH TAKEN	510.0
	236-240	1	46.0	80 FISH TAKEN	510.0
	241-245	1	85.0	80 FISH TAKEN	510.0
	246-250	1	85.0	80 FISH TAKEN	510.0
	251-255	1	85.0	80 FISH TAKEN	510.0
	256-260	1	85.0	80 FISH TAKEN	510.0
	261-265	1	85.0	80 FISH TAKEN	510.0
	266-270	1	85.0	80 FISH TAKEN	510.0
	271-275	1	85.0	80 FISH TAKEN	510.0
	276-280	1	85.0	80 FISH TAKEN	510.0
	281-285	1	85.0	80 FISH TAKEN	510.0
	286-290	1	85.0	80 FISH TAKEN	510.0
	291-295	1	85.0	80 FISH TAKEN	510.0
	296-300	1	85.0	80 FISH TAKEN	510.0
	301-305	1	85.0	80 FISH TAKEN	510.0
	306-310	1	85.0	80 FISH TAKEN	510.0
	311-315	1	85.0	80 FISH TAKEN	510.0
	316-320	1	85.0	80 FISH TAKEN	510.0
	321-325	1	85.0	80 FISH TAKEN	510.0
	326-330	1	85.0	80 FISH TAKEN	510.0
	331-335	1	85.0	80 FISH TAKEN	510.0
	336-340	1	85.0	80 FISH TAKEN	510.0
	341-345	1	85.0	80 FISH TAKEN	510.0
	346-350	1	85.0	80 FISH TAKEN	510.0
	351-355	1	85.0	80 FISH TAKEN	510.0
	356-360	1	85.0	80 FISH TAKEN	510.0
	361-365	1	85.0	80 FISH TAKEN	510.0

POOR ORIGINAL

Table 2.2-66 continued.

Species	Fork length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt.	Mean wt.	No.	Total wt.	Mean wt.
Fathead minnow	91-95	-	-	-	1	5.2	5.2
						1.9	1.9
	136-140	-	-	-	2	11.2	5.6
Pumpkinseed	146-150	1	7.6	7.6	-	-	-
	96-100	1	1.8	1.8	-	-	-
	161-165	-	-	-	2	2.4	1.2
	111-115	-	-	-	1	2.5	2.5
	116-120	2	6.7	3.35	5	21.1	4.22
	121-125	1	6.2	6.2	1	5.4	5.4
	126-130	-	-	-	2	11.8	5.9
	131-135	3	3.9	1.3	-	-	-
	136-140	2	1.6	0.8	1	0.8	0.8
	141-145	-	-	-	2	12.6	6.3
	146-150	1	7.6	7.6	-	-	-

* weight greater than capacity of scales.

Table 2.2-66

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 26-28 April 1976 downstream and upstream from the TONG discharge.

Species	Fork length 5 mm groups	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Channel catfish							
Brown bullhead	246-300	-	-	-	1	378	378.0
	316-320	2	1072	536.0	-	-	-
	321-325	2	1080	540.0	-	-	-
	326-330	1	444	444.0	-	-	-
	346-350	1	550	550.0	-	-	-
	351-355	-	-	-	1	24	24.0
	356-360	-	-	-	1	20	20.0
	361-365	1	30	30.0	-	-	-
	366-370	1	44	44.0	-	-	-
	371-375	1	69	69.0	-	-	-
Rock bass	101-105	-	-	-	1	13	13.0
	106-110	1	26	26.0	-	-	-
	121-125	-	-	-	1	39	39.0
	126-130	1	39	39.0	-	-	-
	141-145	1	52	52.0	-	-	-
	146-150	1	77	77.0	-	-	-
	151-155	1	101	101.0	-	-	-
	156-160	1	130	130.0	-	-	-
	161-165	2	162	81.0	-	-	-
	166-170	2	119	59.5	-	-	-
Rock bass	176-180	-	-	-	1	62	62.0
	191-195	1	72	72.0	-	-	-
	206-210	1	260	260.0	-	-	-
	211-215	-	-	-	1	8	8.0
	216-220	-	-	-	1	13	13.0
	221-225	-	-	-	-	-	-
	226-230	1	39	39.0	-	-	-
	231-235	-	-	-	1	39	39.0
	236-240	1	52	52.0	-	-	-
	241-245	1	77	77.0	-	-	-

Table 2.2-65 continued.

Species	Fork Length (5 mm groups)	No.	DOWNSTREAM		UPSTREAM	
			Total wt. (g)	Mean wt. (g)	Total wt. (g)	Mean wt. (g)
Pumpkinseed	91-95	1	20	20.0	1	19.0
	101-105	4	103	25.8	-	-
	106-110	-	-	-	58	29.0
	111-115	2	69	34.5	2	35.5
	116-120	4	170	42.5	2	44.0
	121-125	1	52	52.0	-	-
	126-130	7	402	57.5	1	60.0
	131-135	1	58	58.0	-	-
	136-140	2	156	78.0	-	-
	141-145	2	160	80.0	-	-
Blackchin shiner	91-95	-	-	-	1	20.0
	111-115	1	30	30.0	-	-
	131-135	1	55	55.0	-	-
	136-140	1	4	4.0	-	-
	141-145	1	68	68.0	20	100.0
	146-150	1	20	20.0	20	100.0
	151-155	1	28	28.0	-	-
	156-160	2	88	44.0	-	-
	161-165	1	115	115.0	-	-

POOR ORIGINAL

Table 2.2-65

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 10-12 May 1976 downstream and upstream from the fish discharge.

Species	Fork Length (5 mm groups)	No.	DOWNSTREAM		UPSTREAM	
			Total wt. (g)	Mean wt. (g)	Total wt. (g)	Mean wt. (g)
Spot tail shiner	91-95	1	70	70.0	7	7.0
Golden shiner	111-115	1	20	20.0	191	100.0
Yellow perch	111-115	2	812	406.0	20	100.0
Brown bullhead	131-135	1	16	16.0	-	-
Channel catfish	131-135	1	36	36.0	-	-
	161-165	-	-	-	1	44
	166-170	1	62	62.0	-	-
	171-175	1	378	378.0	-	-
	176-180	1	38	38.0	2	60.5
	181-185	1	70	70.0	-	-
	186-190	2	171	85.5	-	-
	191-195	1	110	110.0	-	-
	196-200	2	430	215.0	-	-
Rock bass	106-110	1	25	25.0	-	-
	111-115	-	-	-	1	39
	121-125	-	-	-	1	47
	131-135	1	66	66.0	1	52
	136-140	2	164	82.0	-	-
	141-145	3	233	77.7	2	151
	146-150	-	-	-	1	80
	151-155	2	204	102.0	1	93
	156-160	5	512	102.4	-	-
	166-170	2	238	119.0	-	-
	171-175	1	106	106.0	-	-
	191-195	1	180	180.0	-	-
	196-200	1	180	180.0	-	-

Table 2.2-66

Length frequency and mean weights per 5 mm group of fishes taken by trawpnet on 26-28 May 1976 downstream and upstream from the TULEE Discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Spotfin shiner							
	71-75	1	4	4.0			
	76-80	1	7	7.0			
50 FISH TAKEN							
	86-90	1	12	12.0			
	91-95	1	7	7.0			
	96-100	1	16	16.0			
	101-105	1	18	18.0			
20 FISH TAKEN							
White sucker	256-270	1	265	265.0			
Brown bullhead	316-320	-	-	-	1	547	547.0
50 FISH TAKEN							
	326-340	-	-	-	1	619	619.0
20 FISH TAKEN							
Channel catfish	351-355	1	788	788.0			
	86-90	-	-	-	1	6	6.0
	136-140	-	-	-	1	31	31.0
	186-190	-	-	-	1	97	97.0
20 FISH TAKEN							
	221-225	1	128	128.0	-	-	-
	226-230	1	170	170.0	-	-	-
50 FISH TAKEN							
Margined madtom	126-130	-	-	-	1	70	70.0
Rock bass	111-115	-	-	-	1	38	38.0
	121-125	-	-	-	1	50	50.0
	126-130	-	-	-	1	53	53.0
	171-175	1	140	140.0	-	-	-
	181-185	1	145	145.0	1	163	163.0
	196-200	-	-	-	1	208	208.0
20 FISH TAKEN							
Redbreast sunfish	206-210	-	-	-	1	324	324.0
	166-170	1	108	108.0	1	94	94.0
	151-155	2	213	106.5	1	121	121.0
	156-160	1	178	178.0	1	115	115.0
20 FISH TAKEN							
	161-165	2	70	35.0	1	10	10.0
	106-110	1	33	33.0	-	-	-
	111-115	2	86	43.0	-	-	-
	116-120	1	59	59.0	1	46	46.0
20 FISH TAKEN							
	126-130	-	-	-	2	121	60.5
	131-135	2	137	68.5	-	-	-
	136-140	1	238	238.0	2	154	77.0
	141-145	1	90	90.0	2	166	83.0
	146-150	-	-	-	1	90	90.0
	156-160	-	-	-	1	126	126.0

Table 2.2-67

Length frequency and mean weights per 5 mm group of fishes taken by trawpnet on 7-9 June 1976 downstream and upstream from the TULEE Discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	wt.	Mean wt.	No.	wt.	Mean wt.
White sucker	211-215	1	1000	1000			
Brown bullhead	251-255	1	108	108.0			
Channel catfish	316-320	1	591	591.0			
Rock bass	166-170	1	51	51.0			
	126-130	1	100	100.0			
	161-165	1	75	75.0			
	156-160	1	100	100.0			
	161-165	1	-	-	1	100	100.0
	166-170	1	110	110.0			
Redbreast sunfish	116-120	1	56	56.0			
	121-125	-	-	-	1	56	56.0
	126-130	1	55	55.0			
	131-135	-	-	-	1	62	62.0
Pumpkinseed	166-170	1	17	17.0			
	161-165	1	42	42.0			
	126-130	1	63	63.0			
	141-145	1	82	82.0	1	83	83.0
	146-150	1	97	97.0			
	151-155	-	-	-	1	108	108.0
Black crappie	161-165	1	101	101.0			
	171-175	1	104	104.0			
	181-185	1	128	128.0			

* weight greater than capacity of scales.

POOR ORIGINAL

Table 7.7-6B cont (Contd.)

Species	Fork Length (5 yr groups)	DOWNSTREAM			UPSTREAM		
		No.	wt. (g)	Mean wt. (g)	No.	wt. (g)	Mean wt. (g)
Rock bass	111-135	-	-	-	1	1000*	-
	136-160	1	90	90.0	-	-	-
	161-185	2	145	72.5	1	72	72.0
	186-210	1	102	102.0	-	-	-
	211-235	-	-	-	1	95	95.0
Redbreast sunfish	111-135	1	120	120.0	-	-	-
	136-160	2	279	139.5	-	-	-
	161-185	1	82	82.0	-	-	-
	186-210	-	-	-	1	103	103.0
Pumpkinseed	111-135	-	-	-	1	158	158.0
	136-160	1	50	50.0	-	-	-
	161-185	1	68	68.0	1	59	59.0
	186-210	1	62	62.0	1	57	57.0
	211-235	2	153	76.5	1	66	66.0
Smallmouth bass	111-135	-	-	-	1	90	90.0
	136-160	1	100	100.0	-	-	-
	161-185	1	71	71.0	-	-	-
	186-210	2	162	81.0	-	-	-
	211-235	1	106	106.0	-	-	-
	236-260	1	148	148.0	-	-	-
	261-285	1	192	192.0	-	-	-
	286-310	2	430	215.0	-	-	-
	311-335	1	270	270.0	-	-	-
White sucker	111-135	-	-	-	1	53	53.0
	136-160	-	-	-	1	53	53.0
	161-185	-	-	-	1	53	53.0
	186-210	-	-	-	1	53	53.0
	211-235	-	-	-	1	53	53.0
	236-260	-	-	-	1	53	53.0
	261-285	-	-	-	1	53	53.0
	286-310	-	-	-	1	53	53.0
	311-335	-	-	-	1	53	53.0
	336-360	-	-	-	1	53	53.0
	361-385	-	-	-	1	53	53.0
	386-410	-	-	-	1	53	53.0
	411-435	-	-	-	1	53	53.0
	436-460	-	-	-	1	53	53.0
	461-485	-	-	-	1	53	53.0
	486-510	-	-	-	1	53	53.0
	511-535	-	-	-	1	53	53.0
	536-560	-	-	-	1	53	53.0
	561-585	-	-	-	1	53	53.0
	586-610	-	-	-	1	53	53.0
	611-635	-	-	-	1	53	53.0
	636-660	-	-	-	1	53	53.0
	661-685	-	-	-	1	53	53.0
	686-710	-	-	-	1	53	53.0
	711-735	-	-	-	1	53	53.0
	736-760	-	-	-	1	53	53.0
	761-785	-	-	-	1	53	53.0
	786-810	-	-	-	1	53	53.0
	811-835	-	-	-	1	53	53.0
	836-860	-	-	-	1	53	53.0
	861-885	-	-	-	1	53	53.0
	886-910	-	-	-	1	53	53.0
	911-935	-	-	-	1	53	53.0
	936-960	-	-	-	1	53	53.0
	961-985	-	-	-	1	53	53.0
	986-1010	-	-	-	1	53	53.0
	1011-1035	-	-	-	1	53	53.0
	1036-1060	-	-	-	1	53	53.0
	1061-1085	-	-	-	1	53	53.0
	1086-1110	-	-	-	1	53	53.0
	1111-1135	-	-	-	1	53	53.0
	1136-1160	-	-	-	1	53	53.0
	1161-1185	-	-	-	1	53	53.0
	1186-1210	-	-	-	1	53	53.0
	1211-1235	-	-	-	1	53	53.0
	1236-1260	-	-	-	1	53	53.0
	1261-1285	-	-	-	1	53	53.0
	1286-1310	-	-	-	1	53	53.0
	1311-1335	-	-	-	1	53	53.0
	1336-1360	-	-	-	1	53	53.0
	1361-1385	-	-	-	1	53	53.0
	1386-1410	-	-	-	1	53	53.0
	1411-1435	-	-	-	1	53	53.0
	1436-1460	-	-	-	1	53	53.0
	1461-1485	-	-	-	1	53	53.0
	1486-1510	-	-	-	1	53	53.0
	1511-1535	-	-	-	1	53	53.0
	1536-1560	-	-	-	1	53	53.0
	1561-1585	-	-	-	1	53	53.0
	1586-1610	-	-	-	1	53	53.0
	1611-1635	-	-	-	1	53	53.0
	1636-1660	-	-	-	1	53	53.0
	1661-1685	-	-	-	1	53	53.0
	1686-1710	-	-	-	1	53	53.0
	1711-1735	-	-	-	1	53	53.0
	1736-1760	-	-	-	1	53	53.0
	1761-1785	-	-	-	1	53	53.0
	1786-1810	-	-	-	1	53	53.0
	1811-1835	-	-	-	1	53	53.0
	1836-1860	-	-	-	1	53	53.0
	1861-1885	-	-	-	1	53	53.0
	1886-1910	-	-	-	1	53	53.0
	1911-1935	-	-	-	1	53	53.0
	1936-1960	-	-	-	1	53	53.0
	1961-1985	-	-	-	1	53	53.0
	1986-2010	-	-	-	1	53	53.0
	2011-2035	-	-	-	1	53	53.0
	2036-2060	-	-	-	1	53	53.0
	2061-2085	-	-	-	1	53	53.0
	2086-2110	-	-	-	1	53	53.0
	2111-2135	-	-	-	1	53	53.0
	2136-2160	-	-	-	1	53	53.0
	2161-2185	-	-	-	1	53	53.0
	2186-2210	-	-	-	1	53	53.0
	2211-2235	-	-	-	1	53	53.0
	2236-2260	-	-	-	1	53	53.0
	2261-2285	-	-	-	1	53	53.0
	2286-2310	-	-	-	1	53	53.0
	2311-2335	-	-	-	1	53	53.0
	2336-2360	-	-	-	1	53	53.0
	2361-2385	-	-	-	1	53	53.0
	2386-2410	-	-	-	1	53	53.0
	2411-2435	-	-	-	1	53	53.0
	2436-2460	-	-	-	1	53	53.0
	2461-2485	-	-	-	1	53	53.0
	2486-2510	-	-	-	1	53	53.0
	2511-2535	-	-	-	1	53	53.0
	2536-2560	-	-	-	1	53	53.0
	2561-2585	-	-	-	1	53	53.0
	2586-2610	-	-	-	1	53	53.0
	2611-2635	-	-	-	1	53	53.0
	2636-2660	-	-	-	1	53	53.0
	2661-2685	-	-	-	1	53	53.0
	2686-2710	-	-	-	1	53	53.0
	2711-2735	-	-	-	1	53	53.0
	2736-2760	-	-	-	1	53	53.0
	2761-2785	-	-	-	1	53	53.0
	2786-2810	-	-	-	1	53	53.0
	2811-2835	-	-	-	1	53	53.0
	2836-2860	-	-	-	1	53	53.0
	2861-2885	-	-	-	1	53	53.0
	2886-2910	-	-	-	1	53	53.0
	2911-2935	-	-	-	1	53	53.0
	2936-2960	-	-	-	1	53	53.0
	2961-2985	-	-	-	1	53	53.0
	2986-3010	-	-	-	1	53	53.0
	3011-3035	-	-	-	1	53	53.0
	3036-3060	-	-	-	1	53	53.0
	3061-3085	-	-	-	1	53	53.0
	3086-3110	-	-	-	1	53	53.0
	3111-3135	-	-	-	1	53	53.0
	3136-3160	-	-	-	1	53	53.0
	3161-3185	-	-	-	1	53	53.0
	3186-3210	-	-	-	1	53	53.0
	3211-3235	-	-	-	1	53	53.0
	3236-3260	-	-	-	1	53	53.0
	3261-3285	-	-	-	1	53	53.0
	3286-3310	-	-	-	1	53	53.0
	3311-3335	-	-	-	1	53	53.0
	3336-3360	-	-	-	1	53	53.0
	3361-3385	-	-	-	1	53	53.0
	3386-3410	-	-	-	1	53	53.0
	3411-3435	-	-	-	1	53	53.0
	3436-3460	-	-	-	1	53	53.0
	3461-3485	-	-	-	1	53	53.0
	3486-3510	-	-	-	1	53	53.0
	3511-3535	-	-	-	1	53	53.0
	3536-3560	-	-	-	1	53	53.0
	3561-3585	-	-	-	1	53	53.0
	3586-3610	-	-	-	1	53	53.0
	3611-3635	-	-	-	1	53	53.0
	3636-3660	-	-	-	1	53	53.0
	3661-3685	-	-	-	1	53	53.0
	3686-3710	-	-	-	1	53	53.0
	3711-3735	-	-	-	1	53	53.0
	3736-3760	-	-	-	1	53	53.0
	3761-3785	-	-	-	1	53	53.0
	3786-3810	-	-	-	1	53	53.0
	3811-3835	-	-	-	1	53	53.0
	3836-3860	-	-	-	1	53	53.0
	3861-3885	-	-	-	1	53	53.0
	3886-3910	-	-	-	1	53	53.0
	3911-3935	-	-	-	1	53	53.0
	3936-3960	-	-	-	1	53	53.0
	3961-3985	-	-	-	1	53	53.0
	3986-4010	-	-	-	1	53	53.0
	4011-4035	-	-	-	1	53	53.0
	4036-4060	-	-	-	1	53	53.0
	4061-4085	-	-	-	1	53	53.0
	4086-4110	-	-	-	1	53	53.0
	4111-4135	-	-	-	1	53	53.0
	4136-4160	-	-	-	1	53	53.0
	4161-4185	-	-	-	1	53	53.0
	4186-4210	-	-	-	1	53	53.0
	4211-4235	-	-	-	1	53	53.0
	4236-4260	-	-	-	1	53	53.0
	4261-4285	-	-	-	1	53	53.0
	4286-4310	-	-	-	1	53	53.0
	4311-4335	-	-	-	1	53	53.0
	4336-4360	-	-	-	1	53	53.0
	4361-4385	-	-	-	1	53	53.0
	4386-4410	-	-	-	1	53	53.0
	4411-4435	-	-	-	1	53	53.0
	4436-4460	-	-	-	1	53	53.0
	4461-4485	-	-	-	1	53	53.0
	448						

POOR ORIGINAL

Table 2.2-69 continued.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Pumpkinseed	100-110	-	62.2	62.2	1	12	12.0
	110-120	1	50.0	50.0	1	64	64.0
	121-125	-	-	-	1	50	50.0
	126-130	4	236	59.5	4	254	63.5
	131-135	2	120	60.0	-	-	-
	136-140	2	128	64.0	-	-	-
	141-145	3	225	75.0	-	-	-
	146-150	-	-	-	1	100	100.0
	151-155	1	100	100.0	-	-	-
	156-160	1	76	76.0	-	-	-
Blackchin shiner	156-160	2	240	120.0	-	-	-
	161-165	1	90	90.0	-	-	-
	166-170	1	110	110.0	-	-	-
	171-175	1	62	62.0	-	-	-
Largemouth bass	161-165	1	82	82.0	-	-	-
	166-170	1	120	120.0	-	-	-
	171-175	1	120	120.0	-	-	-
	176-180	1	166	166.0	-	-	-
	181-185	1	120	120.0	-	-	-
	186-190	1	166	166.0	-	-	-
	191-195	1	166	166.0	-	-	-
	196-200	1	166	166.0	-	-	-
	201-205	1	166	166.0	-	-	-
	206-210	1	166	166.0	-	-	-
	211-215	1	166	166.0	-	-	-
	216-220	1	166	166.0	-	-	-
	221-225	1	166	166.0	-	-	-
	226-230	1	166	166.0	-	-	-
	231-235	1	166	166.0	-	-	-
	236-240	1	166	166.0	-	-	-
	241-245	1	166	166.0	-	-	-
	246-250	1	166	166.0	-	-	-
	251-255	1	166	166.0	-	-	-
	256-260	1	166	166.0	-	-	-
	261-265	1	166	166.0	-	-	-
	266-270	1	166	166.0	-	-	-
	271-275	1	166	166.0	-	-	-
	276-280	1	166	166.0	-	-	-
	281-285	1	166	166.0	-	-	-
	286-290	1	166	166.0	-	-	-
	291-295	1	166	166.0	-	-	-
	296-300	1	166	166.0	-	-	-
	301-305	1	166	166.0	-	-	-
	306-310	1	166	166.0	-	-	-
	311-315	1	166	166.0	-	-	-
	316-320	1	166	166.0	-	-	-
	321-325	1	166	166.0	-	-	-
	326-330	1	166	166.0	-	-	-
	331-335	1	166	166.0	-	-	-
	336-340	1	166	166.0	-	-	-
	341-345	1	166	166.0	-	-	-
	346-350	1	166	166.0	-	-	-
	351-355	1	166	166.0	-	-	-
	356-360	1	166	166.0	-	-	-
	361-365	1	166	166.0	-	-	-
	366-370	1	166	166.0	-	-	-
	371-375	1	166	166.0	-	-	-
	376-380	1	166	166.0	-	-	-
	381-385	1	166	166.0	-	-	-
	386-390	1	166	166.0	-	-	-
	391-395	1	166	166.0	-	-	-
	396-400	1	166	166.0	-	-	-
	401-405	1	166	166.0	-	-	-
	406-410	1	166	166.0	-	-	-
	411-415	1	166	166.0	-	-	-
	416-420	1	166	166.0	-	-	-
	421-425	1	166	166.0	-	-	-
	426-430	1	166	166.0	-	-	-
	431-435	1	166	166.0	-	-	-
	436-440	1	166	166.0	-	-	-
	441-445	1	166	166.0	-	-	-
	446-450	1	166	166.0	-	-	-
	451-455	1	166	166.0	-	-	-
	456-460	1	166	166.0	-	-	-
	461-465	1	166	166.0	-	-	-
	466-470	1	166	166.0	-	-	-
	471-475	1	166	166.0	-	-	-
	476-480	1	166	166.0	-	-	-
	481-485	1	166	166.0	-	-	-
	486-490	1	166	166.0	-	-	-
	491-495	1	166	166.0	-	-	-
	496-500	1	166	166.0	-	-	-
	501-505	1	166	166.0	-	-	-
	506-510	1	166	166.0	-	-	-
	511-515	1	166	166.0	-	-	-
	516-520	1	166	166.0	-	-	-
	521-525	1	166	166.0	-	-	-
	526-530	1	166	166.0	-	-	-
	531-535	1	166	166.0	-	-	-
	536-540	1	166	166.0	-	-	-
	541-545	1	166	166.0	-	-	-
	546-550	1	166	166.0	-	-	-
	551-555	1	166	166.0	-	-	-
	556-560	1	166	166.0	-	-	-
	561-565	1	166	166.0	-	-	-
	566-570	1	166	166.0	-	-	-
	571-575	1	166	166.0	-	-	-
	576-580	1	166	166.0	-	-	-
	581-585	1	166	166.0	-	-	-
	586-590	1	166	166.0	-	-	-
	591-595	1	166	166.0	-	-	-
	596-600	1	166	166.0	-	-	-
	601-605	1	166	166.0	-	-	-
	606-610	1	166	166.0	-	-	-
	611-615	1	166	166.0	-	-	-
	616-620	1	166	166.0	-	-	-
	621-625	1	166	166.0	-	-	-
	626-630	1	166	166.0	-	-	-
	631-635	1	166	166.0	-	-	-
	636-640	1	166	166.0	-	-	-
	641-645	1	166	166.0	-	-	-
	646-650	1	166	166.0	-	-	-
	651-655	1	166	166.0	-	-	-
	656-660	1	166	166.0	-	-	-
	661-665	1	166	166.0	-	-	-
	666-670	1	166	166.0	-	-	-
	671-675	1	166	166.0	-	-	-
	676-680	1	166	166.0	-	-	-
	681-685	1	166	166.0	-	-	-
	686-690	1	166	166.0	-	-	-
	691-695	1	166	166.0	-	-	-
	696-700	1	166	166.0	-	-	-
	701-705	1	166	166.0	-	-	-
	706-710	1	166	166.0	-	-	-
	711-715	1	166	166.0	-	-	-
	716-720	1	166	166.0	-	-	-
	721-725	1	166	166.0	-	-	-
	726-730	1	166	166.0	-	-	-
	731-735	1	166	166.0	-	-	-
	736-740	1	166	166.0	-	-	-
	741-745	1	166	166.0	-	-	-
	746-750	1	166	166.0	-	-	-
	751-755	1	166	166.0	-	-	-
	756-760	1	166	166.0	-	-	-
	761-765	1	166	166.0	-	-	-
	766-770	1	166	166.0	-	-	-
	771-775	1	166	166.0	-	-	-
	776-780	1	166	166.0	-	-	-
	781-785	1	166	166.0	-	-	-
	786-790	1	166	166.0	-	-	-
	791-795	1	166	166.0	-	-	-
	796-800	1	166	166.0	-	-	-
	801-805	1	166	166.0	-	-	-
	806-810	1	166	166.0	-	-	-
	811-815	1	166	166.0	-	-	-
	816-820	1	166	166.0	-	-	-
	821-825	1	166	166.0	-	-	-
	826-830	1	166	166.0	-	-	-
	831-835	1	166	166.0	-	-	-
	836-840	1	166	166.0	-	-	-
	841-845	1	166	166.0	-	-	-
	846-850	1	166	166.0	-	-	-
	851-855	1	166	166.0	-	-	-
	856-860	1	166	166.0	-	-	-
	861-865	1	166	166.0	-	-	-
	866-870	1	166	166.0	-	-	-
	871-875	1	166	166.0	-	-	-
	876-880	1	166	166.0	-	-	-
	881-885	1	166	166.0	-	-	-
	886-890	1	166	166.0	-	-	-
	891-895	1	166	166.0	-	-	-
	896-900	1	166	166.0	-	-	-
	901-905	1	166	166.0	-	-	-
	906-910	1	166	166.0	-	-	-
	911-915	1	166	166.0	-	-	-
	916-920	1	166	166.0	-	-	-
	921-925	1	166	166.0	-	-	-
	926-930	1	166	166.0	-	-	-
	931-935	1	166	166.0	-	-	-
	936-940	1	166	166.0	-	-	-
	941-945	1	166	166.0	-	-	-
	946-950	1	166	166.0	-	-	-
	951-955	1	166	166.0	-	-	-
	956-960	1	166	166.0	-	-	-
	961-965	1	166	166.0	-	-	-
	966-970	1	166	166.0	-	-	-
	971-975	1	166	166.0	-	-	-
	976-980	1	166	166.0	-	-	-
	981-985	1	166	166.0	-	-	-
	986-990	1	166	166.0	-	-	-
	991-995	1	166	166.0	-	-	-
	996-1000	1	166	166.0	-	-	-

* weight greater than capacity of scales.

Table 2.2-69

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 6-8 July 1976 downstream and upstream from the ENR discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Carp	526-580	1	1000*	-	-	507 ± 11.1	507.0
Golden shiner	151-155	-	-	-	1	56	56.0
	156-160	-	-	-	-	-	-
	161-165	-	-	-	-	-	-
	166-170	-	-	-	-	-	-
	171-175	1	62	62.0	-	-	-
	176-180	-	-	-	-	-	-
Quillback	326-330	-	-	-	1	552	552.0
	336-340	-	-	-	-	-	-
	346-350	-	-	-	-	-	-
	356-360	-	-	-	-	-	-
	366-370	-	-	-	-	-	-
	376-380	-	-	-	-	-	-
Brown bullhead	291-295	1	578	578.0	-	-	-
	296-300	-	-	-	1	1000*	-
	306-310	-	-	-	-	-	-
	311-315	1	534	534.0	-	-	-
	316-320	-	-	-	-	-	-
Channel catfish	121-125	1	26	26.0	-	-	-
	126-130	-	-	-	-	-	-
	131-135	-	-	-	1	28	28.0
	136-140	-	-	-	-	-	-
	141-145	-	-	-	-	-	-
	146-150	-	-	-	-	-	-
	151-155	-	-	-	-	-	-
	156-160	-	-	-	-	-	-
	161-165	-	-	-	-	-	-
	166-170	-	-	-	-	-	-
	171-175	-	-	-	-	-	-
	176-180	-	-	-	-	-	-
	181-185	-	-	-	-	-	-
	186-190	-	-	-	-	-	-
	191-195	-	-	-	-	-	-
	196-200	-	-	-	-	-	-
	201-205	-	-	-	-	-	-
	206-210	-	-	-	-	-	-
	211-215	-	-	-	-	-	-
	216-220	-	-	-	-	-	-
	221-225	-	-	-	-	-	-
	226-230	-	-	-	-	-	-
	231-235	-	-	-	-	-	-
	236-240	-	-	-	-	-	-
	241-245	-	-	-	-	-	-
	246-250	-	-	-	-	-	-
	251-255	-	-	-	-	-	-
	256-260	-	-	-	-	-	-
	261-265	-	-	-	-	-	-
	266-270	-	-	-	-	-	-
	271-275	-	-	-	-	-	-
	276-280	-	-	-	-	-	-
	281-285	-	-	-	-	-	-
	286-290	-	-	-	-	-	-
	291-295	-	-	-	-	-	-
	296-300	-	-	-	-	-	-
	301-305	-	-	-	-	-	-
	306-310	-	-	-	-	-	-
	311-315	-	-	-	-	-	-
	316-320	-	-	-	-	-	-
	321-325	-	-	-	-	-	-
	326-330	-	-	-	-	-	-
	331-335	-	-	-	-	-	-
	336-340	-	-	-	-	-	-
	341-345	-	-	-	-	-	-
	346-350	-	-	-	-	-	-
	351-355	-	-	-	-	-	-
	356-360	-	-	-	-	-	-
	361-365	-	-	-	-	-	-
	366-370	-	-	-	-	-	-
	371-375	-	-	-	-	-	-
	376-380	-	-	-	-	-	-
	381-385	-	-	-	-	-	-
	386-390	-	-	-	-	-	-
	391-395	-	-	-	-	-	-
	396-400	-	-	-	-	-	-
	401-405	-	-	-	-	-	-
	406-410	-	-	-	-	-	-
	411-415	-	-	-	-	-	-
	416-420	-	-	-	-	-	-
	421-425	-	-	-	-	-	-
	426-430	-	-	-	-	-	-
	431-435	-	-	-	-	-	-
	436-440	-	-	-	-	-	-
	441-445	-	-	-	-	-	-
	446-450	-	-	-	-	-	-
	451-455	-	-	-	-	-	-
	456-460	-	-	-	-	-	-
	461-465	-	-	-	-	-	-
	466-470	-	-	-	-	-	-
	471-475	-	-	-	-	-	-
	476-480	-	-	-	-	-	-
	481-485	-	-	-	-	-	-
	486-490	-	-	-	-	-	-
	491-495	-	-	-	-	-	-
	496-500	-	-	-	-	-	-
	501-505	-	-	-	-	-	-
	506-510	-	-	-	-	-	-
	511-515	-	-	-	-	-	-
	516-520	-	-	-	-	-	-
	521-525	-	-	-	-	-	-
	526-530	-	-	-	-	-	-
	531-535	-	-	-	-	-	-
	536-540	-	-	-	-	-	-
	541-545	-	-	-	-	-	-
	546-550	-	-	-	-	-	-
	551-555	-	-	-	-	-	-
	556-560	-	-	-	-	-	-
	561-565	-	-	-	-	-	-
	566-570	-	-	-	-	-	-
	571-575	-	-	-	-	-	-
	576-580	-	-	-	-	-	-
	581-585	-	-	-	-	-	-
	586-590	-	-	-	-	-	-
	591-595	-	-	-	-	-	-
	596-600	-	-	-	-	-	-
	601-605	-	-	-	-	-	-
	606-610	-	-	-	-	-	-
	611-615	-	-	-	-	-	-
	616-620	-	-	-	-	-	-
	621-625	-	-	-	-	-	-
	626-630	-	-	-	-	-	-
	631-635	-	-	-	-	-	-
	636-640	-	-	-	-	-	-
	641-645	-	-	-	-	-	-
	646-650	-	-	-	-	-	-
	651-655	-	-	-	-	-	-
	656-660	-	-	-	-	-	-
	661-665	-	-	-	-	-	-
	666-670	-	-	-	-	-	-
	671-675	-	-	-	-	-	-
	676-680	-	-	-	-	-	-
	681-685	-	-	-	-	-	-
	686-690	-	-	-	-	-	-
	691-695	-	-	-	-	-	-
	696-700	-	-	-	-	-	-
	701-705	-	-	-	-	-	-
	706-710	-	-	-	-	-	-
	711-715	-	-	-	-	-	-
	716-720	-	-	-	-	-	-
	721-725	-	-	-	-	-	-
	726-730	-	-	-	-	-	-
	731-735	-	-	-	-	-	-
	736-740	-	-	-	-	-	-
	741-745	-	-	-	-	-	-
	746-750	-	-	-	-	-	-
	751-755	-	-	-	-	-	-
	756-760	-	-	-	-	-	-
	761-765	-	-	-	-	-	-
	766-770	-	-	-	-	-	-
	771-775	-	-	-	-	-	-
	776-780	-	-	-	-	-	-
	781-785	-	-	-	-	-	-
	786-790	-	-	-	-	-	-
	791-795	-	-	-	-	-	-
	796-800	-	-	-	-	-	-
	801-805	-	-	-	-	-	-
	806-810	-	-	-	-	-	-
	811-815	-	-	-	-	-	-
	816-820	-	-	-	-	-	-
	821-825	-	-	-	-	-	-
	826-830	-	-	-	-	-	-
	831-835	-	-	-	-	-	-
	836-840	-	-	-	-	-	-
	841-845	-	-	-	-	-	-
	846-850	-	-	-	-	-	-
	851-855	-	-	-	-	-	-
	856-860	-	-	-	-	-	-
	861-865	-	-	-	-	-	-
	866-870	-	-	-	-	-	-
	871-875	-	-	-	-	-	-
	876-880	-	-	-	-	-	-
	881-885	-	-	-	-	-	-
	886-890	-	-	-	-	-	-
	891-895	-	-	-	-	-	-
	896-900	-	-	-	-	-	-
	901-905	-	-	-	-	-	-
	906-910	-	-	-	-	-	-
	911-915	-	-	-	-	-	-
	916-920	-	-	-	-	-	-
	921-925	-	-	-	-	-	-
	926-930	-	-	-	-	-	-
	931-935	-	-	-	-	-	-
	936-940	-	-	-	-	-	-
	941-945	-	-	-	-	-	-
	946-950	-	-	-	-	-	-
	951-955	-	-	-	-	-	-
	956-960	-	-	-	-	-	-
	961-965	-	-	-	-	-	-
	966-970	-	-	-	-	-	-
	971-975	-	-	-	-	-	-
	976-980	-	-	-	-	-	-
	981-985	-	-	-	-	-	-
	986-990	-	-	-	-	-	-
	991-995	-	-	-	-	-	-
	996-1000	-	-	-	-	-	-
	1001-1005	-	-	-	-	-	-
	1006-1010	-	-	-	-	-	-
	1011-1015	-	-	-	-	-	-
	1016-1020	-	-	-	-	-	-
	1021-1025	-	-	-	-	-	-
	1026-1030	-	-	-	-	-	-
	1031-1035	-	-	-	-	-	-
	1036-1040	-	-	-	-	-	-
	1041-1045	-	-	-	-	-	-
	1046-1050	-	-	-	-	-	-
	1051-1055	-	-	-	-	-	-
	1056-1060	-	-	-	-	-	-
	1061-1065	-	-	-	-	-	-
	1066-1070	-	-	-	-	-	-
	1071-1075	-	-	-	-	-	-
	1076-1080	-	-	-	-	-	-
	1081-1085	-	-	-	-	-	-
	1086-1090	-	-	-	-	-	-
	1091-1095	-	-	-	-	-	-
	1096-1100	-	-	-	-	-	-
	1101-1105	-	-	-	-	-	-
	1106-1110	-	-	-	-	-	-
	1111-1115	-	-	-	-	-	-
	1116-1120	-	-	-	-	-	-
	1121-1125	-	-	-	-	-	-
	1126-1130	-	-	-	-	-	-
	1131-1135	-	-	-	-	-	-
	1136-1140	-	-	-	-	-	-
	1141-1145	-	-	-	-	-	-
	1146-1150	-	-	-	-	-	-
	1151-1155	-	-	-	-	-	-
	1156-1160	-	-	-	-	-	-
	1161-1165	-	-	-	-	-	-
	1166-1170	-	-	-	-	-	-
	1171-1175	-	-	-	-	-	-
	1176-1180	-	-	-	-	-	-
	1181-1185	-	-	-	-	-	-
	1186-1190	-	-	-	-	-	-
	1191-1195	-	-	-	-	-	-
	1196-1200	-	-	-	-	-	-
	1201-1205	-	-	-	-	-	-
	1206-1210	-	-	-	-	-	-
	1211-1215	-	-	-	-	-	-
	1216-1220	-	-	-	-	-	-
	1221-1225	-	-	-	-	-	-
	1226-1230	-	-	-	-	-	-
	1231-1235	-	-	-	-	-	-
	1236-1240	-	-				

Table 2.2-70

Length frequency and mean weights per 5 mm group of fishes taken by trammel net on 19-21 July 1976 downstream and upstream from the Miller discharge.

Species	Length (5 mm group)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. g	Mean wt. g	No.	Total wt. g	Mean wt. g
Carp	551-555	1	1000	-	-	-	-
Golden shiner	296-300	1	1000	-	-	-	-
	161-165	1	160	160.0	-	-	-
Spottail shiner	176-180	1	60	60.0	-	-	-
	96-100	1	13	13.0	-	-	-
White sucker	166-170	1	9	9.0	-	-	-
	156-160	1	175	175.0	-	-	-
White catfish	221-225	1	250	250.0	-	-	-
Brown bullhead	211-215	1	115	115.0	-	-	-
	241-245	1	202	202.0	-	-	-
	266-270	-	-	-	1	295	295.0
	281-285	1	275	275.0	-	-	-
	306-310	-	-	-	1	570	570.0
Channel catfish	116-120	-	-	-	-	-	-
	96-100	1	15	15.0	-	-	-
	116-120	1	17	17.0	-	-	-
	121-125	2	67	33.5	-	-	-
	131-135	1	29	29.0	-	-	-
	141-145	2	59	29.5	-	-	-
	151-155	1	33	33.0	-	-	-
	281-285	1	303	303.0	-	-	-

Table 2.2-70 continued.

Species	Length (5 mm group)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. g	Mean wt. g	No.	Total wt. g	Mean wt. g
Largehead sucker	151-155	1	62	62.0	-	-	-
Rock bass	151-155	2	272	136.0	-	-	-
	156-160	1	102	102.0	-	-	-
Pumpkinseed	91-95	-	-	-	1	10	10.0
	116-120	1	49	49.0	-	-	-
	121-125	-	-	-	1	21	21.0
	126-130	3	160	53.3	-	-	-
	131-135	2	138	69.0	-	-	-
	136-140	1	80	80.0	-	-	-
	156-160	1	92	92.0	-	-	-
	151-155	1	98	98.0	-	-	-
Bluegill	136-140	-	-	-	-	-	-
	151-155	1	71	71.0	-	-	-
Black crappie	126-130	1	72	72.0	-	-	-
	161-165	2	159	79.5	-	-	-
	166-170	2	185	92.5	-	-	-
	171-175	1	118	118.0	-	-	-
	186-190	2	267	133.5	-	-	-
	186-190	1	117	117.0	-	-	-
	196-200	2	378	189.0	-	-	-
	201-205	2	139	69.5	-	-	-
	206-210	1	120	120.0	-	-	-
	211-215	1	167	167.0	-	-	-
Yellow perch	186-190	1	97	97.0	-	-	-
	196-200	1	111	111.0	-	-	-

* weight greater than capacity of scales.

Table 2.2-71

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 2-4 August 1976 downstream and upstream from the PINE RIVER barge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Carp	511-515	-	-	-	1	1000*	-
Golden shiner	321-325	1	1000*	-	-	-	-
	356-360	1	57	57.0	-	-	-
	161-165	1	72	72.0	50	1130	22.6
Quillback	191-195	1	107	107.0	-	-	-
	376-380	1	1000*	-	-	-	-
Brown bullhead	601-605	1	1000*	-	30	1130	37.7
	251-255	-	-	-	1	211	211.0
	291-295	1	366	366.0	-	-	-
Channel catfish	301-305	-	-	-	1	661	661.0
	111-115	1	21	21.0	-	-	-
	121-125	2	50	25.0	-	-	-
	126-130	3	78	26.0	1	22	22.0
	131-135	-	-	-	5	152	30.4
	136-140	-	-	-	2	52	26.0
Rock bass	396-400	-	-	-	1	372	372.0
	96-100	-	-	-	2	65	32.5
	111-115	-	-	-	1	30	30.0
	116-120	-	-	-	1	57	57.0
	161-165	-	-	-	1	43	43.0
Redbreast sunfish	156-160	-	-	-	1	82	82.0
	151-155	-	-	-	2	160	80.0
	156-160	1	100	100.0	-	-	-
	161-165	1	110	110.0	-	-	-
	171-175	-	-	-	1	120	120.0

Table 2.2-71 continued.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Pumpkinseed	86-90	-	-	-	1	17	17.0
	111-115	-	-	-	1	37	37.0
	116-120	-	-	-	1	117	117.0
	121-125	2	95	47.5	2	79	39.5
	126-130	1	52	52.0	1	139	139.0
	131-135	-	-	-	4	226	56.5
	136-140	-	-	-	1	68	68.0
	141-145	1	83	83.0	-	-	-
	146-150	1	82	82.0	1	91	91.0
Bluegill	111-115	1	91	91.0	-	-	-
	126-130	1	50	50.0	50	1130	22.6
	156-160	1	112	112.0	-	-	-
White crappie	166-170	1	175	175.0	-	-	-
	181-185	1	97	97.0	20	1230	61.5
Black crappie	176-180	1	107	107.0	-	-	-
	191-195	1	128	128.0	-	-	-
	196-200	2	118	59.0	50	1130	22.6
	211-215	2	123	61.5	-	-	-
	226-230	1	202	202.0	-	-	-
Galley	121-125	1	21	21.0	50	1130	22.6

* weight greater than capacity of scales.

POOR ORIGINAL

Table 2.2-72

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 16-18 August 1976 downstream and upstream from the INHS Discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Carp	530-550	1	10000	-	30	1150	38.3
Quillback	371-375	-	-	-	1	10000	-
	411-415	-	-	-	1	10000	-
	436-440	1	10000	-	-	-	-
White sucker	271-275	1	250	250.0	-	-	-
	321-325	1	280	280.0	1	550	550.0
Channel catfish	256-260	1	208	208.0	-	-	-
	366-370	-	-	-	1	550	550.0
	411-415	-	-	-	1	0	0.0
Rock bass	106-110	1	115	115.0	1	21	21.0
Redbreast sunfish	91-95	-	-	-	-	-	-
	166-170	2	210	105.0	-	-	-
Pumpkinseed	106-110	-	-	-	1	21	21.0
	126-130	1	68	68.0	1	63	63.0
	141-145	1	68	68.0	-	-	-
	151-155	1	83	83.0	-	-	-
Bluegill	161-165	1	117	117.0	30	1150	38.3
Black crappie	176-180	1	96	96.0	1	106	106.0
	206-210	1	162	162.0	-	-	-
	216-220	1	198	198.0	-	-	-
Yellow perch	216-220	1	131	131.0	30	1150	38.3

Weight greater than capacity of scales.

Table 2.2-73

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 1-3 September 1976 downstream and upstream from the INHS Discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Quillback	371-375	1	10000	-	30	1150	38.3
	411-415	-	-	-	1	10000	-
	436-440	-	-	-	-	-	-
White catfish	216-220	30	1150	38.3	1	10000	-
Yellow perch	276-280	30	1150	38.3	1	300	300.0
Channel catfish	126-130	-	-	-	2	65	32.5
	141-145	-	-	-	1	0	0.0
	146-150	-	-	-	1	0	0.0
	151-155	-	-	-	2	77	38.5
	161-165	1	60	60.0	-	-	-
	171-175	-	-	-	-	-	-
Rock bass	106-110	-	-	-	1	10000	-
	126-130	1	63	63.0	-	-	-
	146-150	-	-	-	1	100	100.0
	171-175	1	118	118.0	-	-	-
	216-220	1	160	160.0	-	-	-
Pumpkinseed	141-145	-	-	-	1	38	38.0
	146-150	-	-	-	2	114	57.0
	151-155	30	1150	38.3	2	106	53.0
	166-170	1	96	96.0	1	66	66.0
	171-175	1	83	83.0	1	83	83.0
Bluegill	166-170	1	NA	NA	30	1150	38.3
	171-175	1	51	51.0	-	-	-
White crappie	161-165	1	67	67.0	30	1150	38.3
Black crappie	161-165	2	148	74.0	-	-	-
	181-185	1	104	104.0	-	-	-
	186-190	1	150	150.0	-	-	-
	206-210	-	-	-	1	152	152.0
	221-225	-	-	-	1	24	24.0
	241-245	1	30	30.0	-	-	-
Yellow perch	121-125	30	1150	38.3	1	67	67.0
	161-165	1	104	104.0	1	83	83.0

Weight greater than capacity of scales.

NA = Not Available.

POOR ORIGINAL

1565 115

Table 2.2-74 continued.

Species	Fork length (5 mm group)	DOWNSTREAM			UPSTREAM		
		No.	Total wt.	Mean wt.	No.	Total wt.	Mean wt.
Pumpkinseed	90-95	1	17	17.0	1	18.7	18.7
	95-100	2	35	17.5	1	29	29.0
	96-100	-	-	-	1	23	23.0
	116-120	-	-	-	1	45	45.0
	121-125	-	-	-	1	45	45.0
	126-130	-	-	-	1	57	57.0
	136-140	-	-	-	3	189	63.0
	141-145	1	76	76.0	-	-	-
	146-150	1	86	86.0	1	90	90.0
	151-155	1	106	106.0	-	-	-
Bluegill	146-170	1	130	130.0	-	-	-
	96-100	1	17	17.0	-	-	-
	106-110	2	75	37.5	-	-	-
	131-135	1	50	50.0	-	-	-
White crappie	166-170	1	125	125.0	-	-	-
	76-80	1	4	4.0	-	-	-
	176-180	1	76	76.0	-	-	-
	226-230	-	-	-	-	-	-
	231-235	1	184	184.0	-	-	-
Black crappie	136-140	-	-	-	1	40	40.0
	141-145	-	-	-	1	50	50.0
	161-165	-	-	-	1	66	66.0
	166-170	1	90	90.0	-	-	-
	171-175	-	-	-	1	95	95.0
	181-185	1	117	117.0	-	-	-
	186-190	1	103	103.0	-	-	-
	191-195	1	136	136.0	-	-	-
	201-205	-	-	-	2	293	146.5
	206-210	1	163	163.0	-	-	-
	216-220	1	199	199.0	-	-	-
	226-230	1	200	200.0	-	-	-
	231-235	1	276	276.0	1	220	220.0
Walleye	211-215	1	104	104.0	30	30 FISH TAKEN	-

* weights taken after preservation in 10% formalin.
 * weight greater than capacity of scales.

Table 2.2-76

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 13-15 September 1976 downstream and upstream from the PINE discharge.

Species	Fork length (5 mm group)	DOWNSTREAM			UPSTREAM		
		No.	Total wt.	Mean wt.	No.	Total wt.	Mean wt.
Alexlife	136-140	1	18	18.0	16	16.0	16.0
Musellunge	496-500	2	71	35.5	20	20 FISH TAKEN	-
Carp	431-435	-	-	-	1	1000*	-
	501-505	-	-	-	1	1000*	-
	551-555	1	1000*	-	-	-	-
	621-625	1	1000*	-	-	-	-
	641-645	1	1000*	-	-	-	-
Golden shiner	166-170	2	153	76.5	-	-	-
	176-180	1	93	93.0	-	-	-
	186-190	1	108	108.0	-	-	-
Quillback	196-200	1	130	130.0	-	-	-
	316-320	1	672	672.0	-	-	-
	346-350	1	819	819.0	-	-	-
	356-360	1	730	730.0	-	-	-
	371-375	1	106*	-	-	-	-
White sucker	181-185	1	1000*	-	-	-	-
	336-340	1	530	530.0	1	608	608.0
Brown bullhead	321-325	1	530	530.0	-	-	-
Channel catfish	161-165	1	26	26.0	20	20 FISH TAKEN	-
Rock bass	131-135	-	-	-	1	50	50.0
	166-170	1	110	110.0	-	-	-
	171-175	1	108	108.0	-	-	-
	176-180	2	269	134.5	-	-	-
	181-185	1	161	161.0	-	-	-
Redbreast sunfish	231-235	1	300	300.0	-	-	-
	76-80	1	6	6.0	-	-	-
	91-95	1	12	12.0	-	-	-
	146-150	1	77	77.0	-	-	-
	151-155	1	100	100.0	-	-	-
	171-175	3	393	131.0	-	-	-

POOR ORIGINAL

Table 2.2-75

Length frequency and mean weights per 5 mm group of fishes taken by trapnet on 27-29 September 1976 downstream and upstream from the INESS Discharge.

Species	Fork Length (5 mm group)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Golden shiner	161-165	1	76	76.0			
	181-185	3	300	100.0	30 FISH TAKEN		
Quillback	216-220	1	176	176.0			
	351-355	1	1000*	-	80 FISH TAKEN		
Yellow bullhead	276-280	1	867	867.0			
	306-310	1	448	448.0	335 FISH TAKEN		
Brown bullhead	311-315	1	510	510.0	30 FISH TAKEN		
	186-190	1	80	80.0			
Channel catfish	251-255	1	172	172.0	80 FISH TAKEN		
	121-125	1	38	38.0			
Rock bass	126-130	-	-	-	1	46	46.0
	156-160	1	80	80.0	-	-	-
161-165	-	-	-	-	1	93	93.0
	176-180	1	140	140.0	-	-	-
181-185	-	1	174	174.0	-	-	-
	191-195	1	186	186.0	-	-	-
Redbreast sunfish	86-90	-	-	-	1	9	9.0
	116-120	-	-	-	1	37	37.0
156-160	-	1	110	110.0	-	-	-
	166-170	1	176	176.0	-	-	-
Pumpkinseed	86-90	-	-	-	1	12	12.0
	96-100	1	23	23.0	-	-	-
101-105	-	1	22	22.0	-	-	-
	106-110	1	27	27.0	1	30	30.0
116-120	-	2	83	41.5	-	-	-
	121-125	-	-	-	1	50	50.0
126-130	-	-	-	-	1	50	50.0
	131-135	-	-	-	2	123	61.5
136-140	-	-	-	-	1	82	82.0
	146-150	2	164	82.0	3	276	92.0
151-155	-	1	106	106.0	-	-	-
	156-160	1	102	102.0	-	-	-
171-175	-	-	-	-	1	12	12.0

Table 2.2-75 continued.

Species	Fork Length (5 mm group)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Bluntnose	171-175	1	159	159.0	30 FISH TAKEN		
	66-70	1	5	5.0			
White crappie	91-95	1	10	10.0			
	96-100	1	10	10.0			
166-170	-	1	77	77.0	30 FISH TAKEN		
	206-210	1	136	136.0			
221-225	-	2	390	195.0			
	251-255	1	230	230.0			
Black crappie	66-70	-	-	-	1	4	4.0
	96-100	-	-	-	1	18	18.0
191-195	-	1	136	136.0	-	-	-
	201-205	1	143	143.0	-	-	-
216-220	-	1	188	188.0	-	-	-
	231-235	1	230	230.0	-	-	-
236-240	-	1	317	317.0	-	-	-
	251-255	1	284	284.0	-	-	-

* weight greater than capacity of scales.

Table 2.2-76

Length frequency and mean weights per 5 m group of fishes taken by trawnet on 18-20 October 1976 downstream and upstream from the T112 discharge.

Species	Fork length (5 m groups)	DOWNSTREAM			UPSTREAM		
		No.	wt. (g)	Mean (g)	No.	wt. (g)	Mean (g)
Carp	671-675	1	1000*	-	-	-	-
Channel catfish	726-730	1	1000*	-	-	-	-
	1-1-125	-	-	-	1	25	25.0
	166-150	-	-	-	1	39	39.0
	151-155	1	55	52.0	-	-	-
	156-160	1	62	62.0	-	-	-
	161-165	3	164	54.7	-	-	-
	171-175	1	68	68.0	-	-	-
	181-185	-	-	-	1	86	86.0
	216-220	-	-	-	1	124	124.0
	221-225	-	-	-	1	117	117.0
	226-230	1	130	130.0	-	-	-
	246-250	-	-	-	1	175	175.0
	256-260	2	425	212.5	-	-	-
	266-270	-	-	-	1	230	230.0
	276-280	1	106	106.0	-	-	-
	281-285	1	256	256.0	1	282	282.0
	296-300	1	309	309.0	-	-	-
	301-305	1	368	368.0	-	-	-
	311-315	-	-	-	1	307	307.0
	316-320	1	175	175.0	1	106	106.0
Rock bass	331-335	1	500	500.0	-	-	-
	116-120	-	-	-	1	31	31.0
	126-130	-	-	-	1	62	62.0
	136-140	-	-	-	1	62	62.0
	141-145	30 FISH TAKEN	-	-	1	66	66.0
	161-165	-	-	-	1	91	91.0
	181-185	-	-	-	1	153	153.0
	201-205	1	152	152.0	-	-	-
	246-250	1	163	163.0	-	-	-

* weight greater than capacity of scales.

Table 2.2-77

Length frequency and mean weights per 5 m group of fishes taken by trawnet on 25-27 October 1976 downstream and upstream from the T112 discharge.

Species	Fork length (5 m groups)	DOWNSTREAM			UPSTREAM		
		No.	wt. (g)	Mean (g)	No.	wt. (g)	Mean (g)
Channel catfish	156-160	1	11	11.0	-	-	-
	161-165	1	48	48.0	-	-	-
	166-170	1	136	136.0	-	-	-
	171-175	1	58	58.0	-	-	-
	191-195	1	88	88.0	-	-	-
	206-210	1	182	182.0	-	-	-
	256-260	1	204	204.0	-	-	-
	276-280	1	220	220.0	-	-	-
	301-305	1	12	12.0	-	-	-
	306-310	1	116	116.0	-	-	-

POOR ORIGINAL

Table 2.2-78

Length frequency and mean weights per 5 mm group of fishes taken by seine on 2 March 1976 downstream and upstream from the TRINIS discharge.

Species	Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Conely shiner	31-35	1	0.3	0.30			
	56-60	30 FISH TAKEN					
Common shiner	61-65	1	2.0	2.00			
	21-25	1	2.3	2.30			
Spottail shiner	46-50	1	0.2	0.20			
	61-65	1	1.1	1.10			
	66-70	-	-	-			
Swallowtail shiner	16-20	1	2.8	2.80			
	21-25	1	0.1	0.10			
	31-35	6	0.8	0.13			
	36-40	1	0.4	0.40			
	41-45	1	0.6	0.60			
Spotfin shiner	16-20	7	0.6	0.09			
	21-25	19	2.1	0.11			
	26-30	8	1.3	0.16			
	31-35	3	0.9	0.30			
	36-40	1	0.6	0.60			
	41-45	-	-	-			
Bluntnose minnow	21-25	1	0.2	0.20			
	31-35	1	0.4	0.40			
	36-40	-	-	-			
	41-45	-	-	-			
Blacknose dace	51-55	-	-	-			
Redbreast sunfish	31-35	30 FISH TAKEN					
	36-40	1	1.0	1.00			
Tessellated darter	41-45	1	0.5	0.50			
	56-60	2	2.5	1.25			
Banded darter	36-40	30 FISH TAKEN					
	41-45	1	0.2	0.20			
	51-55	2	1.2	0.60			

Table 2.2-79

Length frequency and mean weights per 5 mm group of fishes taken by seine on 15 March 1976 downstream and upstream from the TRINIS discharge.

Species	Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Common shiner	31-35	2	2.6	1.30			
Spottail shiner	46-50	1	1.6	1.60			
	56-60	1	1.8	1.80			
	66-70	1	3.2	3.20			
Swallowtail shiner	16-20	1	0.1	0.10			
	21-25	-	-	-			
	26-30	2	0.5	0.25			
	31-35	-	-	-			
	36-40	1	0.6	0.60			
	46-50	1	1.0	1.00			
Spotfin shiner	16-20	2	5.6	2.80			
	21-25	8	1.0	0.13			
	26-30	8	1.6	0.20			
	31-35	13	4.3	0.33			
	36-40	5	1.9	0.38			
	41-45	2	1.5	0.75			
	51-55	2	3.7	1.85			
	56-60	3	3.7	1.23			
Bluntnose minnow	31-35	1	1.4	1.40			
	41-45	1	0.7	0.70			
	46-50	2	2.4	1.20			
	51-55	2	3.1	1.55			
	56-60	1	0.7	0.70			
Redbreast sunfish	46-50	1	0.7	0.70			
Tessellated darter	41-45	1	1.3	1.30			
	51-55	1	1.3	1.30			
	56-60	1	1.3	1.30			

POOR ORIGINAL

1565 119

Table 2.2-81

Length frequency and mean weights per 5 m group of fishes taken by seine on 11 April 1976 downstream and upstream from the LPS discharge.

Species	Length (5 m groups)	Downstream			Upstream		
		No.	Total wt.	Mean wt.	No.	Total wt.	Mean wt.
Spot-tail shiner							
	51-55	1	1.6	1.6	1	1.6	1.6
	56-60	6	12.5	2.08	6	12.5	2.08
	61-65	1	7.6	7.6	1	7.6	7.6
	66-70	2	1.8	0.9	2	1.8	0.9
	71-75	2	3.4	1.7	2	3.4	1.7
Spot-tail shiner 16-20							
	26-30	2	0.2	0.1	2	0.2	0.1
Rock bass shiner							
	61-65	2	0.4	0.2	2	0.4	0.2
Spot-tail shiner							
	16-20	1	0.1	0.1	1	0.1	0.1
	21-25	1	0.3	0.3	1	0.3	0.3
	26-30	1	0.3	0.3	1	0.3	0.3
	31-35	1	1.4	1.4	1	1.4	1.4
	36-40	1	1.7	1.7	1	1.7	1.7
Bluntnose minnow							
	21-25	1	0.1	0.1	1	0.1	0.1
	26-30	1	0.1	0.1	1	0.1	0.1
Pumpkinseed							
	36-40	1	0.2	0.2	1	0.2	0.2
Tessellated darter							
	51-55	1	0.9	0.9	1	0.9	0.9
	56-60	1	0.2	0.2	1	0.2	0.2
Banded darter							
	56-60	1	0.2	0.2	1	0.2	0.2

* 0.1 means reading was below the accuracy of the scales.

Table 2.2-80

Length frequency and mean weights per 5 m group of fishes taken by seine on 19 March 1976 downstream and upstream from the LPS discharge.

Species	Length (5 cm groups)	Downstream			Upstream		
		No.	Total wt.	Mean wt.	No.	Total wt.	Mean wt.
Spot-tail shiner							
	56-60	2	2.5	1.25	2	2.5	1.25
	61-65	2	3.7	1.85	-	-	-
	66-69	6	14.0	2.33	-	-	-
	61-65	6	15.6	2.60	-	-	-
	66-70	2	6.0	3.00	1	1.2	1.20
Spot-tail shiner							
	21-25	1	1.2	1.2	-	-	-
	31-35	1	0.5	0.50	-	-	-
Spotfin shiner							
	21-25	1	0.1	0.10	2	0.3	0.15
	26-30	4	1.0	0.25	1	0.3	0.30
	31-35	1	0.4	0.40	-	-	-
	36-40	9	5.1	0.57	3	1.8	0.60
	41-45	2	1.8	0.90	2	1.5	0.75
	46-50	2	2.1	1.05	1	0.9	0.90
Bluntnose minnow							
	16-20	-	-	-	1	0.6	0.60
	41-45	6	5.1	0.85	2	1.9	0.95
Bluntnose darter							
	51-55	2	1.3	0.65	-	-	-
	56-60	1	2.0	2.00	1	2.2	2.20
Pumpkinseed							
	16-20	1	0.2	0.20	-	-	-
Channel catfish							
	46-70	1	3.0	3.00	-	-	-
Tessellated darter							
	41-45	2	1.3	0.65	6	2.5	0.43
	46-50	1	0.8	0.80	3	2.5	0.83
	51-55	3	3.0	1.00	6	7.0	1.17
	56-60	-	-	-	1	1.3	1.30
Banded darter							
	16-20	2	1.0	0.50	-	-	-
	21-25	1	0.7	0.70	-	-	-

POOR ORIGINAL

POOR ORIGINAL

Table 2.2-83

Length frequency and mean weights per 5 m group of fishes taken by seine on 10 May 1976 downstream and upstream from the TREC discharge.

Species	Length (5 m group)	Downstream			Upstream		
		No.	Mean wt. (g)	Total wt. (g)	No.	Mean wt. (g)	Total wt. (g)
Bluntnose shiner	41-65	1	1.90	1.90	1	1.90	1.90
Tessellated darter	41-65	1	1.90	1.90	1	1.90	1.90

Table 2.2-82

Length frequency and mean weights per 5 m group of fishes taken by seine on 30 April 1976 downstream and upstream from the TREC discharge.

Species	Length (5 m group)	Downstream			Upstream		
		No.	Mean wt. (g)	Total wt. (g)	No.	Mean wt. (g)	Total wt. (g)
Spotfin shiner	41-65	2	1.90	3.80	1	1.90	1.90
Spotfin shiner	56-60	2	2.50	5.00	1	2.50	2.50
Spotfin shiner	61-65	2	2.80	5.60	1	2.80	2.80
Spotfin shiner	16-20	1	0.60	0.60	2	1.10	2.20
Spotfin shiner	21-25	2	0.10	0.20	3	0.40	1.20
Spotfin shiner	26-30	1	0.20	0.20	3	0.70	2.10
Spotfin shiner	36-40	2	1.10	2.20	1	1.10	1.10
Spotfin shiner	41-65	3	2.90	8.70	1	2.90	2.90
Spotfin shiner	66-70	6	1.10	6.60	1	1.10	1.10
Spotfin shiner	71-75	2	1.90	3.80	1	1.90	1.90
Spotfin shiner	86-90	1	2.20	2.20	1	2.20	2.20
Spotfin shiner	91-95	2	3.20	6.40	1	3.20	3.20
Spotfin shiner	96-100	2	6.00	12.00	1	6.00	6.00
Spotfin shiner	101-105	1	2.10	2.10	1	2.10	2.10
Spotfin shiner	106-110	1	1.10	1.10	1	1.10	1.10
Tessellated darter	36-40	1	0.60	0.60	1	0.60	0.60
Tessellated darter	41-45	1	0.60	0.60	1	0.60	0.60

* 0.1 means rounding was below the accuracy of the scales.

Table 2.2-85

Length frequency and mean weights per 5 mm group of fishes taken by seine on 7 June 1976 downstream and upstream from the THINS Discharge.

Species	Fork Length (5 mm groups)	No.	DOWNSTREAM		UPSTREAM	
			Total wt. (g)	Mean wt. (g)	Total wt. (g)	Mean wt. (g)
Spottail shiner	6-10	72	0.8	0.01	46	1.0
	11-15	327	8.5	0.03	265	5.4
	16-20	16	1.0	0.06	16	0.7
Swallowtail shiner	36-40		NO FISH TAKEN		1	0.8
	41-45				1	1.0
Spotfin shiner	21-25	2	0.4	0.20		
	26-30	4	1.0	0.25		
	31-35				3	1.3
	36-40				5	2.8
	41-45				7	5.8
	46-50				9	10.0
	51-55				10	16.8
Bluntnose minnow	11-15	4	2.0	0.50	NO FISH TAKEN	
	16-20	5	0.4	0.08	1	0.1
	21-25	6	0.7	0.18		
Creek chub	11-15	4	0.2	0.05	NO FISH TAKEN	
	16-20	5	0.5	0.10		
	21-25	7	1.1	0.16		
White sucker/	11-15	14	2.5	0.17	6.4	0.4
Jointhead redbreast	16-20	4	2.1	0.53	6	1.5
	21-25	7	9.8	1.41	5	0.7
	26-30	29	6.5	0.22	2	0.5
Redbreast nontish	51-55	2	6.8	3.40	NO FISH TAKEN	
Bluegill	66-70	1	6.8	6.80	NO FISH TAKEN	
Tessellated darter	11-15	1	0.1*	-	3	0.1
	16-20	1	0.1	0.10	2	0.1
	21-25	1	0.1	0.10		
Shield darter	6-10				1	0.1*
	11-15	2	0.1*	-	15	0.5

* 0.1 means reading was below the accuracy of the scales.

Table 2.2-84

Length frequency and mean weights per 5 mm group of fishes taken by seine on 24 May 1976 downstream and upstream from the THINS Discharge.

Species	Fork Length (5 mm groups)	No.	DOWNSTREAM		UPSTREAM	
			Total wt. (g)	Mean wt. (g)	Total wt. (g)	Mean wt. (g)
Spottail shiner	11-15	1	0.1*	-	NO FISH TAKEN	
	61-65	2	6.7	3.35		
	66-70	1	3.6	3.60		
Swallowtail shiner	31-35	1	0.2	0.20	1	0.5
Spotfin shiner	21-25	1	0.1*	-	NO FISH TAKEN	
Creek chub	11-15	3	0.2	0.07	NO FISH TAKEN	
	16-20	25	1.3	0.05	NO FISH TAKEN	
White sucker/	16-20					
Shorthead redbreast	21-25	5	0.4	0.08		
Smallmouth bass	86-90				1	9.0
	106-110	1	19.8	19.80		
	111-115				1	22.8
	151-155	1	56.2	56.20		
Tessellated darter	16-20	1	0.1*	-	NO FISH TAKEN	

* 0.1 means reading was below the accuracy of the scales.

Table 2.2-87

Length frequency and mean weights per 5 mm group of fishes taken by seine on 7 July 1976 downstream and upstream from the TMS discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Common shiner							
11-15	11-15	5	0.2	0.04	5	0.2	0.04
16-20	16-20	8	0.6	0.08	8	0.6	0.08
21-25	21-25	10	1.0	0.10	10	1.0	0.10
Spot-tail shiner							
11-15	11-15	6	0.6	0.07	17	0.6	0.06
16-20	16-20	168	11.8	0.07	83	5.6	0.07
21-25	21-25	186	22.6	0.12	20	8.6	0.12
26-30	26-30	46	9.7	0.21	28	5.7	0.20
31-35	31-35	6	1.7	0.28	2	0.6	0.30
36-40	36-40	2	1.1	0.55	-	-	-
Spot-tail shiner							
11-15	11-15	-	-	-	1	0.1*	-
16-20	16-20	-	-	-	2	0.2	0.10
31-35	31-35	-	-	-	1	0.6	0.60
36-40	36-40	1	0.8	0.80	9	5.8	0.65
41-45	41-45	-	-	-	11	16.8	0.83
46-50	46-50	-	-	-	5	5.6	1.06
51-55	51-55	-	-	-	7	1.5	1.75
56-60	56-60	-	-	-	1	2.6	2.60
Bluntnose minnow							
21-25	21-25	1	5.3	5.30	-	-	-
36-40	36-40	-	-	-	-	-	-
Bluegill							
11-15	11-15	1	2.9	2.90	-	-	-
16-20	16-20	2	1.3	0.65	1	1.2	1.20
21-25	21-25	4	3.2	0.80	-	-	-
26-30	26-30	7	7.0	1.00	1	1.0	1.00
31-35	31-35	2	2.6	1.30	-	-	-
White sucker							
16-20	16-20	3	0.4	0.13	-	-	-
21-25	21-25	5	1.0	0.20	-	-	-
26-30	26-30	16	4.6	0.28	-	-	-
31-35	31-35	66	27.9	0.42	-	-	-
36-40	36-40	51	27.2	0.53	-	-	-
41-55	41-55	8	8.2	1.03	-	-	-
46-50	46-50	7	10.0	1.43	-	-	-
White sucker							
36-40	36-40	1	2.5	2.50	1	0.6	0.60
Northern hog sucker							
16-20	16-20	1	0.1	0.10	2	0.2	0.10
21-25	21-25	4	0.8	0.20	3	0.6	0.20
26-30	26-30	10	1.9	0.19	5	1.1	0.22
31-35	31-35	1	0.3	0.30	-	-	-
36-40	36-40	1	0.1*	0.1*	-	-	-
Brown bullhead							
26-30	26-30	2	0.2	0.10	-	-	-
31-35	31-35	2	0.0	0.00	-	-	-

Table 2.2-86

Length frequency and mean weights per 5 mm group of fishes taken by seine on 21 June 1976 downstream and upstream from the TMS discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Common shiner							
	11-15	1	0.1*	-	3	0.1	0.03
	16-20	-	-	-	-	-	-
Spot-tail shiner							
	11-15	47	1.3	0.03	51	1.5	0.03
	16-20	106	7.1	0.07	67	6.2	0.07
	21-25	110	16.8	0.13	67	8.9	0.13
	26-30	15	3.1	0.21	17	7.6	0.45
	31-35	1	0.1	0.10	-	-	-
	36-40	1	0.2	0.20	-	-	-
Spot-tail shiner							
	11-15	2	0.7	0.35	1	0.5	0.50
	16-20	4	2.0	0.50	5	3.0	0.60
	21-25	5	4.7	0.94	1	1.0	1.00
	26-30	3	4.4	1.47	1	1.5	1.50
	31-35	2	3.6	1.80	1	1.7	1.70
	36-40	-	-	-	1	2.9	2.90
Bluntnose minnow							
	21-25	1	6.1	6.10	-	-	-
Bluegill							
	21-25	1	0.2	0.20	2	0.4	0.20
	26-30	5	1.5	0.30	-	-	-
	31-35	13	6.4	0.49	-	-	-
	36-40	2	1.2	0.60	-	-	-
White sucker							
	26-40	1	2.8	2.80	1	0.6	0.60
Shorthead redhorse							
	21-25	1	1.2	1.20	-	-	-
	26-30	-	-	-	8	0.5	0.06
	31-35	-	-	-	6	0.8	0.13
	36-40	-	-	-	2	0.6	0.30
White sucker							
	11-15	2	0.1	0.05	-	-	-
	16-20	19	1.1	0.06	3	0.7	0.07
	21-25	2	0.3	0.15	1	0.1	0.10
	26-30	2	0.6	0.30	-	-	-
	31-35	2	0.7	0.35	-	-	-
	36-40	1	0.1	0.10	-	-	-
Shorthead redhorse							
	11-15	-	-	-	1	0.1*	0.1*
	16-20	-	-	-	2	0.1	0.05
	21-25	1	0.2	0.20	9	1.7	0.19
	26-30	6	1.6	0.27	10	1.9	0.19
	31-35	-	-	-	1	0.3	0.30
	36-40	-	-	-	1	0.1*	0.1*
White sucker							
	11-15	-	-	-	-	-	-
	16-20	-	-	-	-	-	-
	21-25	-	-	-	-	-	-
	26-30	-	-	-	-	-	-
	31-35	-	-	-	-	-	-
	36-40	-	-	-	-	-	-

POOR ORIGINAL

Table 2.2-87 continued.

Species	Fork Length (5 age groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Channel catfish	16-20	3	0.4	0.13	NO FISH TAKEN		
	21-25	9	1.6	0.18			
Rock bass	6-10	NO FISH TAKEN			1	0.1*	
Smallmouth bass	16-20	1	0.3	0.30	2	0.4	0.20
	21-25	1	0.3	0.30	5	1.0	0.20
	26-30	6	2.1	0.35	5	1.6	0.32
	31-35	1	0.7	0.70	1	0.5	0.50
	46-50	-	-	-	1	2.0	2.00
White crappie	16-20	2	2.6	1.30	NO FISH TAKEN		
	21-25	1	0.2	0.20			
	26-30	2	0.7	0.35			
Tessellated darter	16-20	-	-	-	4	6.8	0.20
	21-25	2	0.3	0.15	5	0.9	0.18
	26-30	9	2.1	0.23	-	-	-
	31-35	6	1.8	0.30	2	0.6	0.30
Banded darter	21-25	NO FISH TAKEN			2	0.2	0.10

* 0.1 means reading was below the accuracy of the scales.

Table 2.2-88

Length frequency and mean weights per 5 age group of fishes taken by seine on 19 July 1976 downstream and upstream from the TWIN discharge.

Species	Fork Length (5 age groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
SAFP	21-25	1	2.0	2.00	NO FISH TAKEN		
Common shiner	26-30	1	0.2	0.20	7	1.6	0.23
	31-35	1	0.4	0.40	-	-	-
Common shiner	21-25	NO FISH TAKEN			3	0.9	0.30
	26-30	3	0.9	0.30	-	-	-
	36-40	1	0.8	0.80	-	-	-
Spottail shiner	16-20	7	0.6	0.09	1	0.9	0.09
	21-25	28	3.9	0.14	25	10.2	0.41
	26-30	60	14.2	0.24	95	22.0	0.23
	31-35	41	14.9	0.36	39	13.8	0.35
	36-40	11	6.8	0.62	7	6.1	0.87
	41-45	-	-	-	2	1.8	0.90
Spotfin shiner	36-40	-	-	-	1	2.0	0.67
	41-45	1	0.7	0.70	17	14.1	0.83
	46-50	2	2.8	1.40	18	21.0	1.17
	51-55	1	1.7	1.70	10	17.6	1.76
	61-65	2	-	-	1	2.9	2.90
Bluntnose minnow	21-25	1	2.2	2.20	1	6.9	6.90
	16-20	1	0.1*	-	-	-	-
	26-30	-	-	-	1	0.3	0.30
	31-35	-	-	-	1	0.5	0.50
	46-50	1	1.9	1.90	-	-	-
Blacknose dace	26-30	1	2.2	2.20	-	-	-
	36-40	1	0.7	0.70	NO FISH TAKEN		
Creek chub	46-50	1	1.5	1.50	2	2.9	1.45
Fallfish	51-55	-	-	-	1	1.7	1.70
Golden shiner	41-45	NO FISH TAKEN			1	1.7	1.70

Table 2.2-88 continued.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
White sucker	26-30	1	0.3	0.30	-	-	-
	31-35	-	-	-	1	0.6	0.60
	36-40	5	3.3	0.66	4	2.9	0.73
	41-45	2	5.0	1.00	1	1.1	1.10
	46-50	2	2.2	1.10	1	1.1	1.10
Shorthead redhorse	26-30	1	0.3	0.30	2	0.7	0.35
	31-35	10	4.5	0.45	1	0.4	0.40
	36-40	12	7.5	0.63	2	1.2	0.60
	41-45	5	4.2	0.84	2	2.1	1.05
	46-50	1	1.2	1.20	-	-	-
Brown bullhead	31-35	1	0.5	0.50	-	-	-
Rock bass	41-45	2	2.3	1.15	-	-	-
	46-50	-	-	-	1	0.2	0.20
Redbreast sunfish	26-30	1	4.1	4.10	-	-	-
Pumpkinseed	46-50	-	-	-	1	1.3	1.30
Smallmouth bass	26-30	-	-	-	1	0.5	0.50
	36-40	-	-	-	2	1.5	0.75
	41-45	1	1.3	1.30	4	4.5	1.13
	46-50	1	1.5	1.50	-	-	-
	61-65	-	-	-	1	3.5	3.50
	66-70	1	4.6	4.60	-	-	-
	71-75	-	-	-	1	5.3	5.30
	76-80	2	16.9	7.65	-	-	-
Largemouth bass	41-45	-	-	-	1	1.0	1.00
	46-50	-	-	-	1	0.1	0.10
	51-55	1	0.2	0.20	1	0.1	0.10
	56-60	2	0.5	0.25	5	1.0	0.20
	61-65	10	3.0	0.30	2	0.5	0.25
	66-70	1	0.6	0.60	-	-	-
	71-75	-	-	-	1	0.3	0.30
	76-80	-	-	-	-	-	-

* 0.1 means reading was below the accuracy of the scales.

Table 2.2-89

Length frequency and mean weights per 5 mm group of fishes taken by seine on 2 August 1976 downstream and upstream from the TINS Discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Common shiner	41-45	-	-	-	1	0.8	0.80
	46-50	-	-	-	4	1.3	0.33
	51-55	3	1.6	0.53	3	1.6	0.53
	56-60	-	-	-	1	0.7	0.70
Spottail shiner	21-25	2	0.5	0.25	4	0.8	0.20
	26-30	11	3.0	0.27	43	11.5	0.27
	31-35	36	15.3	0.43	43	17.7	0.41
	36-40	18	11.7	0.65	24	16.4	0.60
	41-45	4	3.6	0.90	2	1.7	0.85
Spotfin shiner	51-55	-	-	-	1	1.8	1.80
	56-60	2	0.3	0.15	1	0.2	0.20
	61-65	1	0.3	0.30	6	1.5	0.25
	66-70	1	0.5	0.50	-	-	-
	71-75	-	-	-	1	0.6	0.60
Fallfish	41-45	-	-	-	1	0.6	0.60
	46-50	-	-	-	1	0.6	0.60
White sucker	41-45	3	3.3	1.10	2	2.3	1.15
	46-50	2	2.6	1.30	1	1.6	1.60
	51-55	-	-	-	1	1.2	1.20
	56-60	1	2.5	2.50	-	-	-
	61-65	1	3.3	3.30	-	-	-
	66-70	1	1.0	1.00	-	-	-
Northern hog sucker	41-45	-	-	-	1	1.0	1.00
	46-50	-	-	-	1	1.0	1.00
Shorthead redhorse	31-35	1	0.4	0.40	2	1.2	0.60
	36-40	5	3.7	0.74	-	-	-
	41-45	3	3.2	1.07	2	1.8	0.90
	46-50	1	1.2	1.20	-	-	-
Yellow perch	26-30	1	0.4	0.40	1	0.5	0.50
	31-35	1	0.5	0.50	-	-	-
	36-40	2	2.1	1.05	2	2.0	1.00
	41-45	3	0.7	0.23	-	-	-
Channel catfish	21-25	16	4.8	0.30	-	-	-
	26-30	51	25.8	0.51	-	-	-
	31-35	29	18.6	0.64	-	-	-
	36-40	3	2.7	0.90	-	-	-
	41-45	1	1.7	1.70	-	-	-
	46-50	-	-	-	-	-	-

POOR ORIGINAL

Table 2.2-89 continued.

Species	Fork Length (5 yr group)	JPMF:TH:AS			JF:TH:AM		
		No.	wt. (g)	Mean wt. (g)	Total wt. (g)	Mean wt. (g)	Mean wt. (g)
Redbreast sunfish	19-20	2	2.2	1.10	-	-	-
Smallmouth bass	36-60	1	1.3	1.30	1	1.2	1.20
	41-65	1	1.3	1.30	1	1.2	1.20
	51-55	2	4.0	2.00	-	-	-
	61-65	2	7.7	3.85	1	3.5	3.50
	66-70	1	4.9	4.90	-	-	-
White crappie	76-80	1	0.3	0.30	1	7.2	7.20
Tessellated darter	26-30	1	-	-	3	0.7	0.23
	31-35	15	4.5	0.30	10	2.7	0.27
	36-60	12	5.8	0.48	13	5.5	0.42
	51-55	5	2.2	0.28	2	1.2	0.60

* 0.1 means reading was below the accuracy of the scales.

Table 2.2-90

Length frequency and mean weights per 5 mm group of fishes taken by seine on 16 August 1976 downstream and upstream from the MTR discharge.

Species	Fork Length (5 yr group)	JPMF:TH:AS			JF:TH:AM		
		No.	wt. (g)	Mean wt. (g)	Total wt. (g)	Mean wt. (g)	Mean wt. (g)
River chub	46-50	1	2.2	2.20	-	-	-
	51-55	1	0.2	0.20	-	-	-
Common shiner	26-30	-	-	-	2	0.6	0.30
	46-50	1	0.3	0.30	2	2.1	1.05
	51-55	-	-	-	-	-	-
Spottail shiner	26-30	1	0.4	0.40	7	2.2	0.31
	31-35	7	3.0	0.43	14	6.3	0.45
	36-60	23	15.8	0.69	26	17.0	0.65
	41-45	29	28.0	0.97	18	16.4	0.91
	46-50	27	25.5	0.94	10	13.4	1.34
	51-55	7	11.8	1.69	1	2.0	2.00
Swallowtail shiner	91-95	1	11.0	11.00	-	-	-
	26-30	-	-	-	1	0.4	0.40
Spottin shiner	46-50	8	0.6	0.08	1	1.5	1.50
	51-55	5	0.7	0.14	4	0.6	0.15
	26-30	1	0.3	0.30	19	4.7	0.25
	31-35	-	-	-	7	2.8	0.40
	36-60	-	-	-	1	0.5	0.50
	46-50	1	1.5	1.50	1	1.1	1.10
	51-55	1	1.7	1.70	-	-	-
	56-60	1	2.5	2.50	-	-	-
	61-65	3	7.9	2.63	-	-	-
Bluntnose minnow	21-25	4	0.7	0.18	1	0.2	0.20
	26-30	15	7.3	0.26	2	0.7	0.35
	31-35	5	1.7	0.34	1	0.5	0.50
	36-60	1	0.8	0.80	1	0.7	0.70
	41-45	-	-	-	2	1.9	0.95
	46-50	1	1.5	1.50	-	-	-
	51-55	-	-	-	1	2.0	2.00

1565 126

Table 2.2-90 continued.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Fallfish	51-55	2	4.4	2.20	NO FISH TAKEN	-	-
White sucker	51-55	1	1.7	1.70	NO FISH TAKEN	-	-
Shorthead redhorse	36-40	1	0.9	0.90	-	-	-
	41-45	-	-	-	-	-	-
	46-50	6	7.8	1.30	1	1.0	1.00
	51-55	2	3.3	1.65	1	2.0	2.00
	56-60	1	2.5	2.50	-	-	-
Rock bass	31-35	NO FISH TAKEN	-	-	2	1.6	0.80
	36-40	-	-	-	1	1.0	1.00
Redbreast sunfish	46-50	1	13.0	13.00	NO FISH TAKEN	-	-
Bluegill	71-75	1	7.6	7.60	NO FISH TAKEN	-	-
	76-80	2	0.9	0.45	1	0.4	0.40
Smallmouth bass	36-40	1	1.0	1.00	-	-	-
	46-50	2	3.1	1.55	1	1.8	1.80
	51-55	2	4.3	2.15	1	2.0	2.00
White crappie	41-45	2	7.0	3.50	-	-	-
	46-50	8	3.8	0.48	-	-	-
	51-55	4	4.9	0.61	-	-	-
	56-60	4	3.5	0.88	-	-	-
Black crappie	46-50	2	2.3	1.15	1	1.2	1.20
	51-55	NO FISH TAKEN	-	-	1	1.7	1.70
	56-60	-	-	-	-	-	-
Tessellated darter	26-30	1	0.2	0.20	1	2.7	2.70
	31-35	18	5.1	0.28	2	0.4	0.20
	36-40	39	16.0	0.41	3	0.9	0.30
	41-45	14	8.0	0.57	4	1.8	0.45
	46-50	2	1.8	0.90	-	-	-

Table 2.2-91

Length frequency and mean weights per 5 mm group of fishes taken by seine on 1 September 1976 downstream and upstream from the THIR discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Lucioperca minnow	41-45	1	1.0	1.00	NO FISH TAKEN	-	-
River chub	51-55	1	1.9	1.90	NO FISH TAKEN	-	-
Golden shiner	51-55	NO FISH TAKEN	-	-	1	1.6	1.60
Common shiner	31-35	2	0.7	0.35	-	-	-
	36-40	-	-	-	1	0.2	0.20
	41-45	NO FISH TAKEN	-	-	4	2.7	0.68
Spottail shiner	31-35	-	-	-	1	0.7	0.70
	36-40	6	4.1	0.68	45	30.1	0.67
	41-45	18	15.6	0.87	34	39.8	1.17
	46-50	8	10.0	1.25	20	26.4	1.22
	51-55	6	9.0	1.50	12	19.7	1.64
	56-60	1	2.2	2.20	5	10.5	2.10
Swallowtail shiner	26-30	-	-	-	1	0.2	0.20
	31-35	3	1.1	0.37	1	1.1	0.37
	36-40	NO FISH TAKEN	-	-	1	0.5	0.50
	41-45	1	1.0	1.00	1	1.0	1.00
	46-50	1	1.2	1.20	1	1.2	1.20
	51-55	-	-	-	1	1.5	1.50
Spotfin shiner	16-20	-	-	-	18	0.5	0.03
	21-25	9	1.3	0.14	17	0.7	0.04
	26-30	19	4.3	0.23	63	14.5	0.23
	31-35	5	1.7	0.34	38	14.4	0.38
	36-40	3	1.8	0.60	12	6.6	0.55
	41-45	-	-	-	1	0.8	0.80
Bluntnose minnow	41-45	1	2.1	2.10	-	-	-
	46-50	1	0.2	0.20	-	-	-
	51-55	1	0.2	0.20	-	-	-
	56-60	1	0.5	0.50	2	1.0	0.50
Creek chub	36-40	4	2.6	0.65	1	0.8	0.80
	41-45	NO FISH TAKEN	-	-	1	1.7	1.70
White sucker	41-45	1	0.8	0.80	-	-	-
	56-60	1	2.5	2.50	-	-	-
	66-70	-	-	-	1	4.0	4.00
	71-75	-	-	-	1	5.0	5.00
	76-80	-	-	-	2	12.3	6.15

Table 2.2-91 continued.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPPER/AV		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Northern hog sucker	61-65	1	2.7	2.70	1	2.7	2.70
	66-70	1	3.7	3.70			
Shorthead redhorse	46-50	-	-	-			
	51-55	2	3.9	1.95	1	1.3	1.30
	56-60	1	2.5	2.50	1	1.9	1.90
	61-65	1	3.3	3.30	-	-	-
White catfish	56-60	1	2.2	2.20	NO FISH TAKEN		
Channel catfish	26-30	1	0.6	0.60			
	31-35	3	1.3	0.43			
	36-40	2	1.5	0.75			
	41-45	9	8.7	0.97			
	46-50	57	73.7	1.29	NO FISH TAKEN		
	51-55	43	70.6	1.64			
	56-60	8	17.7	2.21			
	61-65	2	4.9	2.45			
	66-70	1	3.4	3.40			
	71-75	1	3.9	3.90			
Pumpkinseed	61-65	1	0.1	0.10	1	1.5	1.50
Bluegill	21-25	1	0.1	0.10	1	1.2	1.20
Smallmouth bass	41-45	1	0.1	0.10			
	56-60	1	2.9	2.90			
	66-70	1	3.9	3.90			
Black crappie	31-35	1	12.0	12.00			
	51-55	1	2.2	2.20			
	56-60	1	3.1	3.10			
Translocated darter	46-50	2	8.5	4.25			
	51-55	3	1.0	0.33			
	56-60	4	2.0	0.50			
	61-65	14	8.3	0.59			
	66-70	4	2.7	0.68			

Table 2.2-92

Length frequency and mean weights per 5 mm group of fishes taken by seine on 13 September 1976 downstream and upstream from the 1918S discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPPER/AV		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Common shiner	21-25	1	0.2	0.20	1	0.3	0.30
	26-30	-	-	-			
	31-35	-	-	-	1	0.6	0.60
	36-40	-	-	-	1	1.0	1.00
	41-45	-	-	-	3	3.9	1.30
	46-50	-	-	-	2	4.1	2.05
	51-55	-	-	-	1	2.2	2.20
Common shiner	56-60	1	1.5	1.50			
Spot-tail shiner	21-25	8	7.6	0.95	2	1.6	0.80
	26-30	8	10.1	1.26	8	7.6	0.95
	31-35	11	5.6	0.51	14	17.4	1.24
	36-40	4	2.4	0.60	11	19.1	1.74
	41-45	2	5.6	2.80	5	10.8	2.16
Spot-tail shiner	46-50	1	0.1	0.10	1	0.1	0.10
	51-55	6	1.1	0.18	12	1.6	0.13
	56-60	9	2.3	0.26	12	2.9	0.24
	61-65	4	1.1	0.28	7	2.7	0.41
	66-70	-	-	-	9	5.2	0.58
	71-75	-	-	-	4	3.4	0.85
	76-80	-	-	-	1	1.4	1.40
Bluntnose minnow	21-25	1	0.2	0.20	1	0.9	0.90
	26-30	-	-	-	4	0.9	0.23
	31-35	1	0.5	0.50	1	0.3	0.30
	36-40	1	0.6	0.60	1	0.7	0.70
	41-45	-	-	-	7	6.7	0.96
	46-50	-	-	-	1	1.1	1.10
Shorthead redhorse	56-60	1	4.2	4.20			

POOR ORIGINAL

Table 2.2-93

Length frequency and mean weights per 5 mm group of fishes taken by seine on 29 September 1976 downstream and upstream from the INLSE discharge.

Species	Fork length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Common shiner	36-40	1	0.9	0.90	1	0.2	0.20
Spot tail shiner	41-45	2	2.0	1.00	2	2.4	1.20
	46-50	5	6.3	1.26	2	2.4	1.20
	51-55	3	4.8	1.60	3	5.5	1.83
	56-60	4	9.2	2.30	5	10.6	2.08
	61-65	5	15.9	3.18	-	-	-
Swallowtail shiner	16-20	1	0.1	0.10	-	-	-
Spotfin shiner	11-15	25	0.7	0.03	2	0.9	0.45
	16-20	80	4.7	0.06	30	1.7	0.06
	21-25	125	16.5	0.13	46	5.2	0.11
	26-30	45	9.8	0.22	10	2.0	0.20
	31-35	3	1.0	0.33	2	0.9	0.45
Bluntnose minnow	26-30	-	-	-	1	6.0	6.00
	16-20	2	0.1	0.05	-	-	-
	21-25	3	0.6	0.13	-	-	-
	26-30	4	0.7	0.18	-	-	-
	36-40	3	1.6	0.53	-	-	-
	41-45	2	2.0	1.00	-	-	-
	46-50	1	1.1	1.10	1	1.3	1.30
	51-55	-	-	-	2	1.8	0.90
	56-60	1	2.1	2.10	1	3.9	3.90
Green chub	26-30	-	-	-	-	-	-

Table 2.2-92 continued.

Species	Fork length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Channel catfish	31-35	2	1.1	0.55	-	-	-
	36-40	7	4.8	0.69	-	-	-
	41-45	21	21.6	1.03	-	-	-
	46-50	20	27.4	1.37	-	-	-
	51-55	24	41.6	1.73	1	1.7	1.70
	56-60	12	26.5	2.21	1	2.0	2.00
	61-65	1	2.2	2.20	-	-	-
Rock bass	41-45	-	-	-	1	1.9	1.90
Redbreasted sunfish	21-25	1	3.1	3.10	-	-	-
Blackchin shiner	21-25	1	0.3	0.30	-	-	-
Bluegill	21-25	1	0.2	0.20	-	-	-
Smallmouth bass	41-45	-	-	-	1	1.3	1.30
	56-60	1	3.2	3.20	1	2.8	2.80
	61-65	1	3.5	3.50	-	-	-
Tessellated darter	26-30	3	1.6	0.53	-	-	-
	41-45	10	6.6	0.66	10	17.8	0.59
	46-50	5	3.9	0.78	12	9.8	0.82
Banded darter	26-30	20	17.8	0.89	1	0.3	0.30

* 0.1 is a reading was below the accuracy of the scales.

Table 2.2-93 continued.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Fallfish	66-70	1	3.9	3.90	NO FISH TAKEN		
Shorthead redhorse	81-85	1	6.5	6.50	NO FISH TAKEN		
	56-60	1	1.9	1.90	NO FISH TAKEN		
Channel catfish	61-65	1	3.0	3.00			
	46-50	2	3.1	1.55			
	51-55	5	8.8	1.76			
Redbreast sunfish	71-75				1	4.5	4.50
	61-65	1	1.6	1.60	NO FISH TAKEN		
Bluegill	36-40	1	0.9	0.90	NO FISH TAKEN		
	61-65	1	4.5	4.50	NO FISH TAKEN		
Smallmouth bass	61-65	1	3.5	3.50			
	56-70	1	4.7	4.70	NO FISH TAKEN		
Tessellated darter	91-95	1	10.0	10.00			
	36-40	4	2.0	0.50	2	1.0	0.50
	41-45	16	11.1	0.69	6	3.9	0.65
	46-50	49	41.4	0.84	10	8.3	0.83
	51-55	13	14.1	1.08	7	7.9	1.13
	56-60	3	4.3	1.43			
Handed darter	31-35				3	1.2	0.40
	36-40	NO FISH TAKEN			6	3.1	0.52
	41-45				2	1.4	0.70

Table 2.2-94

Length frequency and mean weights per 5 mm group of fishes taken by seine on 19 October 1976 downstream and upstream from the PINS discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM			UPSTREAM		
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)
Common shiner	56-60	NO FISH TAKEN			1	2.0	2.00
Common shiner	46-50	2	2.0	1.00			
Spottail shiner	36-40	6	6.8	1.13	4	3.7	0.93
	41-45	6	9.8	1.63	4	4.6	1.15
	46-50	5	9.1	1.82	9	14.3	1.59
	51-55	1	2.7	2.70	2	15.4	7.70
	56-60	5	14.1	2.82	2	5.2	2.60
	61-65	2			1	1.2	1.20
Smallmouth shiner	21-25	5	0.6	0.12			
	26-30	1	0.2	0.20			
	31-35	4	1.8	0.45			
	36-40	2	1.7	0.85			
	41-45	2	1.0	0.50			
Rock bass	16-20	6	0.5	0.08	4	0.3	0.08
Spot, in shiner	21-25	41	5.5	0.13	1	0.2	0.20
	26-30	51	11.2	0.22			
	31-35	30	10.5	0.35			
	36-40	21	10.9	0.52			
	41-45	6	4.3	0.72			
	46-50	2	2.0	1.00			
Bluntnose minnow	26-30	1	2.0	2.00			
	31-35	3	1.2	0.40	1	0.4	0.40
	36-40	5	3.4	0.68	3	2.1	0.70
	41-45	6	5.2	0.87	10	9.0	0.90
	46-50	1	1.4	1.40	7	8.7	1.24
	51-55	1	1.7	1.70	1	1.4	1.40
	56-60	2	4.2	2.10	1	1.9	1.90
	61-65	1	2.3	2.30			

1565 130

POOR ORIGINAL

Table 2.2-94 continued.

Species	Fork Length (5 mm groups)	DOWNSTREAM				UPSTREAM			
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)		
Fathead minnow	36-40	1	0.8	0.80	NO FISH TAKEN				
Fathead minnow	41-45	1	4.9	4.90	NO FISH TAKEN				
Fathead minnow	46-50	1	4.6	4.60	NO FISH TAKEN				
Rock bass	31-35	1	0.6	0.60	NO FISH TAKEN				
Smallmouth bass	56-60	-	-	-	1	3.3	3.30		
	66-70	-	-	-	1	5.2	5.20		
	71-75	1	6.0	6.00	-	-	-		
Tessellated darter	31-35	3	1.1	0.37	1	0.5	0.50		
	36-40	4	2.1	0.53	4	2.0	0.50		
	41-45	15	9.6	0.64	16	8.6	0.61		
	46-50	30	24.8	0.83	24	20.6	0.86		
	51-55	17	17.6	1.04	10	12.3	1.23		
	56-60	4	5.2	1.30	1	1.7	1.70		
Banded darter	21-25	1	0.2	0.20	-	-	-		
	31-35	-	-	-	1	0.5	0.50		
	36-40	-	-	-	2	1.2	0.60		

POOR ORIGINAL

Table 2.2-95

Length frequency and mean weights per 5 mm group of fishes taken by seine on 25 October 1976 downstream and upstream from the Tullis discharge.

Species	Fork Length (5 mm groups)	DOWNSTREAM				UPSTREAM			
		No.	Total wt. (g)	Mean wt. (g)	No.	Total wt. (g)	Mean wt. (g)		
Common shiner	51-55	3	4.5	1.50	-	-	-		
	56-60	4	7.4	1.85	1	1.8	1.80		
	61-65	3	6.8	2.27	1	7.1	2.37		
Common shiner	86-90	30	FISH TAKEN	-	1	6.2	6.20		
Spottail shiner	36-40	2	1.5	0.75	-	-	-		
	41-45	9	8.7	0.97	-	-	-		
	46-50	29	37.1	1.28	6	7.0	1.17		
Spottail shiner	51-55	42	65.9	1.57	15	24.4	1.63		
	56-60	22	46.0	2.09	16	36.6	2.29		
	61-65	28	74.0	2.64	27	74.5	2.76		
Spottail shiner	66-70	8	26.7	3.34	6	27.9	3.69		
	71-75	-	-	-	2	0.2	0.10		
	76-80	-	-	-	2	0.3	0.15		
Spottail shiner	81-85	1	0.3	0.30	1	0.3	0.30		
	86-90	6	2.5	0.42	3	1.2	0.40		
	91-95	4	2.3	0.58	2	1.2	0.60		
Rosyface shiner	51-55	50	FISH TAKEN	-	3	4.4	1.47		
Spottail shiner	56-60	-	-	-	2	3.6	1.80		
	61-65	1	0.1*	-	11	0.4	0.04		
	66-70	17	2.4	0.14	41	4.9	0.12		
Spottail shiner	71-75	11	2.2	0.20	11	2.1	0.19		
	76-80	18	6.8	0.38	2	0.9	0.45		
	81-85	12	6.4	0.53	-	-	-		
Spottail shiner	86-90	6	5.1	0.85	2	2.0	1.00		
	91-95	2	2.1	1.05	-	-	-		
	96-100	1	1.5	1.50	-	-	-		
Bluntnose minnow	61-65	1	3.9	3.90	1	2.9	2.90		
	66-70	1	0.2	0.20	-	-	-		
	71-75	1	0.9	0.90	-	-	-		
Bluntnose minnow	76-80	2	2.1	1.05	3	2.8	0.93		
	81-85	-	-	-	1	2.7	2.70		
	86-90	1	13.9	13.90	30	FISH TAKEN	-		
Quillback	91-95	1	1.0	1.00	30	FISH TAKEN	-		
Redbreast sunfish	36-40	1	0.2	0.20	1	0.2	0.20		
Rock bass	96-100	30	FISH TAKEN	-	1	22.1	22.10		
Bluegill	26-30	1	0.2	0.20	1	0.2	0.20		
Smallmouth bass	86-90	1	10.7	10.70	-	-	-		
	91-95	-	-	-	1	2.7	2.70		
	101-105	-	-	-	1	22.4	22.40		
Tessellated darter	31-35	1	0.2	0.20	-	-	-		
	36-40	5	2.3	0.46	2	0.9	0.45		
	41-45	20	11.5	0.58	2	1.5	0.75		
Tessellated darter	46-50	13	10.5	0.81	5	4.2	0.85		
	51-55	3	2.1	0.70	2	1.2	0.60		
	56-60	-	-	-	-	-	-		

* 0.1 means reading was below the accuracy of the scales.

* 0.1 means reading was below the accuracy of the scales.

1565 131

Table 2.3-1

Summary of number of specimens, number of species, number of collections, and number per collection of fishes taken by trapnet and seine upstream and downstream from the TWINS Discharge during 1974 through 1976.

		Number of Specimens	Number of Species	Number of Collections	Number per Collection
TRAPNET					
Upstream	1974	713	19	36	*19.81
	1975	324	20	51	6.35
	1976	295	19	36	8.19
Downstream	1974	1257	20	72	17.46
	1975	1286	21	87	14.78
	1976	527	24	108	4.88
Total	1974	1970	22	108	18.24
	1975	1610	22	138	11.67
	1976	822	25	144	5.71
SEINE					
Upstream	1974	3460	24	28	123.57
	1975	1558	24	30	51.93
	1976	4914	30	36	136.50
Downstream	1974	5127	28	45	113.93
	1975	5016	29	45	111.47
	1976	5564	31	54	103.04
Total	1974	8587	30	73	117.63
	1975	6574	30	75	87.65
	1976	10478	35	90	116.42

1565 132

POOR ORIGINAL

Table 2.3-2

Kendall-tau correlation coefficients for species composition at trapnet and seine stations during 1974-1975, 1974-1976, and 1975-1976.

	1974-1975	1974-1976	1975-1976
TRAPNET			
1A3	0.65	0.47	0.47
11A2	0.69	0.59	0.70
11A3	0.59	0.35	0.48
9B2	0.62	0.57	0.51
SEINE			
1A2	0.55	0.40	0.22*
16A1	0.60	0.42	0.44
10A2	0.57	0.25	0.26
9A1	0.37	0.34	0.26*
9B3	0.61	0.52	0.37

*Significant at 95% level.

1565 133

POOR ORIGINAL




Figure 2.1-1. Location of trapnet and seine stations in the vicinity of TMINS.

3.0 MACROINVERTEBRATES

The ETS Section 4.1.1E requires semimonthly samples to be taken at five stations in the vicinity of TMINS April through October. Replicate samples were taken both inside and outside of the thermal plume.

3.1 METHODS

Location and description of macroinvertebrate sampling stations are given in Table 3.1-1 and Figure 3.1-1. Four quantitative samples were taken at each station with a 23 x 23 cm (529 cm²) Ponar grab sampler. Samples were returned to the lab, washed through U.S. Standard No. 30 mesh screens, and preserved in a mixture of 10% formalin and rose bengal stain. The stain facilitates sorting of macroinvertebrates from the detritus or sediment present in the sample (Nason and Yevich 1967). Macroinvertebrates were sorted from each sample with unaided eye. Oligochaetes and chironomids were re-sorted under a stereo microscope (10X-15X). Many oligochaetes were damaged when the samples were washed; those with a complete anterior end were counted. Every tenth Limnodrilus picked from the sample was selected for species determination. Tubificid worms used for identification were cleared in Arman's lactophenol and mounted on microscope slides in CMC-9 or Hoyer's mounting media. Naidid worms were cleared when mounted directly in CMC-9 or Hoyer's mounting media. Chironomids were cleared in a warm 5 to 10% solution of KOH and mounted on microscope slides in CMC-9 or Hoyer's mounting media.

POOR ORIGINAL

Organisms were dried in a drying oven at 55 C for 24 hr, cooled in a desiccator, and weighed to the nearest 0.1 mg on a Mettler H31 balance. Gastropods were decalcified in a 7 to 8 M solution of HCL and pelecypod shells were removed manually prior to weight determinations. Weights were not determined for individuals less than 0.1 mg or those retained for taxonomic purposes.

3.1.1 TAXONOMIC TREATMENT

Specimens were identified to genus or species when possible. For diversity calculations, taxonomic comparisons were made by the number of taxa, where a taxon represents the lowest level to which an organism was identified.

Flatworms (Turbellaria) contract when preserved directly in formalin, which renders species identification difficult. For this report, they were determined to the class Turbellaria. The phyla Nemertinea and Nematoda were treated similarly. Numbers and biomass were not calculated for entoprocts and bryozoans because of their colonial nature.

Identification of oligochaetes is dependent on external and internal organs. Most Enchytraeidae, Naididae, and the tubificids Aulodrilus, Branchiura, Pelosclex, and Limnodrilus udekenianus can be identified by their somatic chaetae (hair-like structures present on all oligochaetes except the order Branchiobdellida) or external structures at all stages in their development. Sexually mature specimens are needed for species identifications of most Limnodrilus, Monodrilus, and Tubifex. Limnodrilus was the only tubificid collected possessing only bifurcate crotchet chaetae;

1565 136

88 POOR ORIGINAL

Ilyodrilus and Tubifex additionally possess capilliform chaetae. All immature tubificids without capilliform chaetae were assumed to be immature Limnodrilus. Of the sexually mature Limnodrilus encountered during the sampling period, 88% were L. hoffmeisteri; immature Limnodrilus were grouped with L. hoffmeisteri for interpretation of data. Immature Ilyodrilus and Tubifex collected were recorded as immature tubificids with capilliform chaetae. For data analysis, these were grouped with I. templetoni and T. tubifex and calculated on a percentage basis from the number of sexually mature specimens collected at each station.

Certain species of earthworms (limicolous) inhabit the margins of streams, ponds, lakes, etc. (Reynolds 1975). Limicolous earthworms encountered unidentifiable to the family Lumbriculidae or Sparganophilidae were recorded as "megadriles" (a term which collectively encompasses many families of earthworms).

Leeches (Hirudinea) contract when preserved directly in formalin, which makes identification difficult. Large, mature specimens are needed for genus and species determinations in the family Erpobdellidae; immature or badly distorted specimens were identified only as Erpobdellidae. Mooreobdella microstoma was the only erpobdellid collected identifiable to species. Individuals of M. microstoma were grouped with Erpobdellidae for diversity calculations.

Numbers of larval and pupal Diptera were combined for data interpretation.

If keys were unavailable or incomplete for species separation in some genera, apparently different species of a genus were given letter designators (i.e., Eukiefferiella sp. A, sp. B, etc.). Some genera or species are

POOR ORIGINAL

separable only to groups. The Palpomyia group (Diptera: Ceratopogonidae) contains the genera Bezzia, Probezzia, and Palpomyia, which are inseparable in their larval stages at the present time.

Species identifications in the Chironomidae are difficult. Roback (1976) found many identifications of immatures and adults (by various authors, including himself) were incorrect or nomenclatorially out of date. Major genera and groups constantly undergo revision. Consequently, some specimens identified previously may have been misdetermined.

Misdeterminations in Kennedy (1975) and Polk and Epler (1976) were:

Limnodrilus profundicola (1975, 1976) = L. hoffmeisteri (this report)

Trepobates sp. = Metrobates hesperius

Micropsectra sp. = Tanytarsus sp.

Synonomies from Polk and Epler (1976) are:

Athripsodes = Ceraclea (cf. Morse and Wallace 1975)

Ablabesmyia auriensis = A. mallochi (cf. Roback 1971)

Chironomus attenuatus (as Chironomus sp.) = C. decorus (cf. Sublette and Sublette 1974).

Specimens identified to tribe Tanytarsini in Polk and Epler (1976) were Tanytarsus sp.

Macroinvertebrate taxa found in the Susquehanna River in the vicinity of Three Mile Island are listed in Table 3.1-2. Identification of benthic organisms was aided with keys and descriptions in Beck and Beck (1966), Beck and Beck (1969), Boesel (1974), Brinkhurst and Jamieson (1971), Brown (1972), Burch (1973), Burks (1975), Curry (1958), Edmondson (1959), Foster

1565 138

(1972), Harman and Berg (1971), Hilsenhoff (1975), Hiltunen (1973), Holsinger (1972), Johannsen (1969), Klemm (1972), Mason (1973), Needham and Westfall (1955), Reynolds (1975), Roback (1957, 1974b, 1976), Ross (1972), Saether (1975, 1976), Sawyer (1972), Usinger (1956), Walker (1958), and Walker and Corbet (1975).

3.1.2 DATA ANALYSIS

Densities (number/m²) were calculated for taxa that comprised more than one percent of the total number of organisms collected. Biomass (mg/m²) of selected taxa was determined.

Diversity indices are used to analyze community structure. Monthly estimates of diversity (information per individual) as defined by Shannon's formula:

$$D = \sum_{i=1}^s n_i / n \log_2 n_i / N$$

were computed for each station using the formula:

$$D = C/N (N \log_{10} N - \sum n_i \log_{10} n_i)$$

where D = information per individual, C = 3.321928 (converts log₁₀ to log₂), N = total number of individuals, n_i = total number of individuals in the ith species, s = the number of species in the sample for a station (Lloyd et al. 1968). This index summarizes the number of taxa present and the distribution of individuals among the taxa. Low D values result from a small number of taxa and uneven distribution of individuals. Large numbers of taxa and even distribution of individuals result in high D values.

Benthic communities at each station may have similar D values, but differ in species composition. The similarity in species composition between stations was investigated by an index of percent similarity (Whittaker and Fairbanks 1958). This index is expressed as:

$$PSc = 100 - 0.5 \sum |a - b|$$

where PSc is the percent similarity and a and b are the percentages of a species in samples A and B. This is a quantitative measure of the relative similarity of species composition in two samples. The PSc is only an empirical measure and is not an estimate of a statistical parameter of the population from which the samples are drawn. Values range from 0 to 100. A value of 0 indicates the species composition is entirely different and a value of 100 indicates complete similarity between the two samples.

Analysis of variance (ANOVA), randomized block design (Sokal and Rohlf 1969), was performed to compare numbers of taxa between stations and assess changes over time. The Student-Newman-Keuls multirange test (Woolf 1968) was performed if significant differences existed in the ANOVA. The multirange test was designed to specify which sample means (\bar{y}) differed significantly from each other.

Changes in populations of Limnodrilus hoffmeisteri for the 1974 through 1976 sample dates were examined; a three-factor ANOVA was used (Sokal and Rohlf 1969). Comparisons were made for years, sample dates, the macroinvertebrate stations, and interactions between these factors. No quantitative data was available for April 1974 at Station 11A2. The ANOVA was performed

on data collected from May through October 1974, 1975, and 1976. Numbers of L. profundicola misdetermined in 1974 and 1975 were included with L. hoffmeisteri in the analysis.

The distribution of benthic organisms is clumped rather than random; a logarithmic transformation $[\log_{10}(y+1)]$ was used on densities of L. hoffmeisteri for each replicate to normalize the data (Elliot 1971). Logarithmic transformation prevents attention being focused on small variations in numbers which may not be statistically significant (Brinkhurst and Jamieson 1971). The Student-Newman-Keuls multirange test ($P = 0.05$) was employed if differences were indicated.

3.2 RESULTS

Results of April through October macroinvertebrate collections are reported in Tables 3.2-1 through 3.2-35. A total of 90,567 specimens (112 taxa) was taken during 1976 (Table 3.2-36). The three most abundant taxa (84.8% of the specimens) were Limnodrilus hoffmeisteri (65,516 specimens, 72.3%), Chironomus decorus (9,084, 10.0%), and Nais elinguis (2,265, 2.5%). Other taxa that comprised more than 1% of the total were Pisidium sp. (1,646 specimens, 1.8%), Procladius sp. (1,199, 1.3%), Gammarus fasciatus (1,084, 1.2%), L. clapparedianus (1,059, 1.2%), and L. udekemianus (912, 1.0%).

Limnodrilus hoffmeisteri was usually the most abundant organism collected at each station (Table 3.2-36 and Figure 3.2-1). Nais elinguis was most abundant at Station 1A1 in May and Chironomus decorus was most abundant at Station 9B1 in May and at 1A2 in June.

1565 141

The freshwater polychaete, Manayunkia speciosa, was collected for the first time in 1976. M. speciosa has been taken from the Susquehanna River in Conowingo Pond (D. Wahl, Ichthyological Associates, Inc., personal communication)*. Formerly believed to be rare, recent papers (Hiltunen 1965; Poe and Stefan 1974; Spencer 1976) have shown M. speciosa to be common and abundant (over 45,000/m² in Lake Erie) in many areas. Its small size (<3 mm) probably caused it to be overlooked.

High numbers of the limpet Ferrissia were collected at Stations 1A1 and 1A2 on 19 October 1976. The reason for the high numbers was not readily apparent. The limpets may have drifted to Stations 1A1 and 1A2 from rocks located upstream.

3.2.1 DENSITY

Densities of benthic macroinvertebrates were dominated by a few taxa (Table 3.2-37). Limnodrilus hoffmeisteri was most abundant and ranged from 390/m² (Station 1A2 in April) to 9,776/m² (Station 11A1 in October). Peak densities occurred in June at Stations 1A1, 11A2, and 9B1 and in September at 1A2. Lowest densities occurred in April at all stations.

Peak densities of Chironomus decorus occurred at all stations in June. The greatest density (4,863/m²) was at Station 9B1. No C. decorus were collected at Stations 1A2, 11A1, and 9B1 in April, and at 11A1 in October.

Nais elinguis reached a peak density of 4,612/m² at Station 1A1 in May. No N. elinguis were collected after June.

3.2.2 BIOMASS

At most stations and months the pleurocerid snail, Coniobasis virginica, comprised the greatest biomass and peaked at $8,734.1 \text{ mg/m}^2$ at Station 11A2 in September (Table 3.2-38). The greatest biomass ($2,103.3 \text{ mg/m}^2$) for L. hoffmeisteri occurred at 11A2 in June. C. decorus outweighed all organisms at Station 9B1 in May and June, and peaked ($3,022.4 \text{ mg/m}^2$) in June.

3.2.3 DIVERSITY (D)

Monthly D values at each station ranged from 0.66 to 2.76 (Table 3.2-39). The lowest D value (0.66) occurred at Station 9B1 in April; L. hoffmeisteri comprised 91.3% of the organisms in the sample. The highest D value (2.76) occurred at 1A2 in May; L. hoffmeisteri accounted for only 42.8% of the sample. D values were generally higher at Station 1A1.

3.2.4 PERCENT SIMILARITY COMPOSITION

Numbers and percent composition of benthic organisms and calculated PSc indices are presented in Tables 3.2-36 and 3.2-40, respectively. The PSc values were in the intermediate to high affinity range (>50% similarity). Index values indicated that Stations 1A2 and 9B1 were the most similar; Stations 1A1 and 11A1 were the least similar. The PSc values indicated that 1A2, 11A1, 11A2, and 9B1 were more similar in their faunal associations with each other than with Station 1A1.

3.2.6 ANALYSIS OF VARIANCE

Results of the ANOVA (randomized block design) for number of taxa revealed significant differences ($P = 0.05$) between stations and sample

POOR ORIGINAL

dates (Table 3.2-41). Application of the Student-Newman-Keuls multirange test ($P = 0.05$) between station means (\bar{y}) indicated Station 1A1 was significantly different from the other stations and possessed the greatest mean number of taxa (Table 3.2-42). Station 1A2 was similar to 9B1 and Station 11A1 was similar to 11A2 and 9B1. Station 1A2 differed significantly from 11A1 and 11A2 in number of taxa.

The multirange test performed on numbers of taxa between sample dates showed no significant difference between the two sample dates for each month (Table 3.2-43). Comparison of number of taxa by month revealed that the 6 April date was significantly different from the 4 and 18 May dates and the 20 April date differed from 4 May, but was similar to 18 May. The number of taxa taken on 4 May was different from the number taken on 1 June, and similar to 15 June. The 18 May date was similar to both June dates. Numbers of taxa collected on 1 June differed from both July dates and 15 June was similar to both July dates. No significant difference was indicated when comparing July to August, August to September, and September to October. The lowest number of taxa was recorded on both April dates. Numbers of taxa collected in May and June were highest.

Results of the three-factor ANOVA for populations of Limnodrilus hoffmeisteri revealed significant differences ($P = 0.05$) between years, benthic sample dates, stations, and the interaction between these factors. The station factor had the greatest effect on densities of L. hoffmeisteri (Table 3.2-44).

1565 144

The Student-Newman-Keuls multirange test performed on year means revealed all three years were significantly different. Mean densities of L. hoffmeisteri were highest in 1976 and lowest in 1974 (Table 3.2-45).

Application of the multirange test to sample dates showed no significant differences between densities on the two dates sampled each month (Table 3.2-46). Comparison of densities by month revealed that the first and second May sampling dates differed from the first and second June dates. Both June dates had densities similar to the two July dates. Densities for the first July date were significantly different from the first August date and the second July date differed from both August dates. The density for the first August date differed from both September dates and the second August date was similar to the September dates. Both September densities differed from the two October densities. Densities of L. hoffmeisteri were low during October, May, and August. High densities recorded in June, July, and September were not significantly different.

The multirange test applied to densities at the macroinvertebrate stations revealed all were different, except Stations 11A1 and 11A2 (Table 3.2-47). The two stations were similar and had the highest mean densities of the macroinvertebrate stations. Station 1A1 had the lowest mean density of L. hoffmeisteri.

POOR ORIGINAL

1565 145

3.3 DISCUSSION

The nature of the substrate has been found to be the most important factor that determines species distribution (Brinkhurst and Jamieson 1971; Hynes 1972; Harman 1974). Chutter (1969) and deMarch (1976) have demonstrated that changes in substrate result in changes of macroinvertebrate populations. Hynes (1972), Hoopes (1974), Polk et al. (1976), and Massengill (1976) concluded that high flows scour substrates and wash away many organisms. Massengill (1976) and Tebo (1970) found this particularly true of substrates composed of sand and silt (mud) where finer sediments are washed away, leaving sand. Shifting substrates create an unstable habitat (Chutter 1969; Tebo 1970). Massengill (1976) found that ice affects substrates, particularly those composed of sand.

Substrates at Stations 1A1 and 1A2 were basically similar (Table 3.1-1). Substrates at Stations 11A1 and 11A2 were composed of compacted mud with coal and fine sand over a gravel substratum. The substrate at Station 9B1 consisted of mud. During high flows, silt and allochthonous materials were washed away, leaving behind the heavier materials. The substrate of Station 1A1 was more susceptible to changes in river flow.

Ice exerts a heavy influence on the substrate at Stations 1A1 and 1A2. Effects of ice were not as severe at Stations 11A1 and 11A2, where operation of TMINS leaves the area free of ice. In the spring, ice break-up subjects Stations 11A1 and 11A2 to ice scouring. Conditions at Station 9B1 vary from open water to ice cover; the substrate is subjected to scouring.

The dominant organism at all stations was Limnodrilus hoffmeisteri. Brinkhurst and Jamieson (1971) found L. hoffmeisteri to be the most abundant organism in shoreline areas with soft sediments. Carr and Hiltunen (1965) found that abundance of tubificid worms was related to organic enrichment of the substrate. Polk et al. (1976) and results of the three-factor ANOVA indicated that Station 1A1 produced the lowest density and was significantly different from the other stations. The unstable, abrasive, and organically poor sand and coal substrate at 1A1 may account for the low density. Similar densities at Stations 1A2 and 9B1 (Polk et al. 1976) could be attributed to the similarity of habitat; both stations are backwater areas during low flow. Stations 11A1 and 11A2 produced the highest densities and the three-factor ANOVA revealed they were not significantly different. The close proximity and like substrate may account for the similarities at these stations.

The three-factor ANOVA revealed populations of L. hoffmeisteri were highest in early June. Densities were lowest in May and October; periods most affected by ice or high river flow. Kennedy (1966) found that ice cover caused cessation of winter breeding activity in L. hoffmeisteri. Stations significantly affected by flood waters were 1A1, 1A2, and 9B1 (Polk et al. 1976).

Chironomid populations were affected by substrate and reproductive cycles. Polk et al. (1976) found Chironomus populations were affected by the September 1975 flood waters (Hurricane Eloise). Chironomids are holometabolous insects, and population fluctuations are caused in part by pupal

POOR ORIGINAL

emergence (adults are not aquatic). Reproductive cycles for some chironomids range from several generations a year to one every two years (Roback 1974a). Interspecific competition is reduced by staggered breeding periods (Reid 1961; Grant and Mackay 1969), and could account for high numbers of Orthocladinae in the spring, and Chironominae later in the year. Low numbers of C. decorus were observed at Station 11A1, April through October 1976. Reasons for low numbers of C. decorus were not apparent.

Leeches were most abundant at Stations 11A1 and 11A2. Members of the genus Helobdella are known to feed on mollusks (Sawyer 1972; Klemm 1975, 1976). Specimens of Helobdella were often found inside the shells of the snail Coniobasis virginica. This snail possesses an operculum, a hard shell-like disc which closes the shell aperture. Klemm (personal communication) believes the leeches may use the shells as an attachment substrate, as snails preyed upon by Helobdella are non-operculate. Maloney and Chandler (1976) reported that H. lineata feeds on Coniobasis under laboratory conditions. The greatest number of Helobdella collected coincided with the greatest number of Coniobasis collected at 11A1 and 11A2 in 1976.

Erpobdellid leeches were collected only at Stations 11A1 and 11A2. Sawyer (1974) states "no other single factor is more important in restricting the distribution of freshwater leeches than the availability of food organisms". Erpobdellids are predacious on oligochaetes and insect larvae. Oligochaete numbers were highest at 11A1 and 11A2.

1565 148

100 POOR ORIGINAL

Spatial (station) comparisons made with 1976 data indicated that Station 1A1 differed from all other stations in number of taxa and species composition. This was due to substrate variability and drift from upstream locations. Station 1A2 was similar to 9B1 in number of taxa and species composition (91% similar); this may have resulted from similarities in habitat most of the year. Stations 11A1 and 11A2 were similar in number of taxa and species composition (89% similar); this may be caused by station proximity and substrate similarity.

The benthic communities had high numbers of taxa in May and June (Figure 3.3-1). This was attributed to seasonal abundance of naidid worms and orthocladiid Chironomidae. Naidids, especially the genus Nais, mature simultaneously in the spring. Populations reach a maximum in June, then die off rapidly due to degeneration of the digestive tract caused by reproductive activity (Loden 1976). The seasonality of naidids at Three Mile Island concurs with Loden's work.

Wihlm (1970) collected data from numerous authors and found, in general, that diversity (D) values less than one were indicative of "stressed" communities. Values greater than three were indicative of communities associated with "good-excellent" water quality. Ischinger and Nalepa (1966) attributed low diversity of benthic fauna to low habitat diversity rather than poor water quality. Trends in D values along Three Mile Island from 1974 to 1976 indicated "poor" to "good" water quality or habitat diversity (Table 3.3-1, Figure 3.3-2). D values were generally higher at Station 1A1, where conditions were "good". Stations 1A2 and 9B1 displayed generally "fair" values, while 11A1 and 11A2 fluctuated from "poor" to "fair".

No apparent effects on benthic communities at Stations 11A1, 11A2, and 9B1 were noted due to the shutdown of TMINS for refueling (21 February to 27 May 1976) or the dredging operations in front of the Intakes for Units 1 and 2 (August through September 1976).

Trace amounts of oil were present in most samples at all stations. No effects of oil on macroinvertebrates have been observed.

In summary, the variability of the substrate influenced by river flow and ice exerts more stress on the benthic communities along Three Mile Island than the operation of TMINS. This completes the macroinvertebrate requirements of the ETS for Unit 1; the program will be continued as specified in the ETS for Unit 2.

- Beck, E.C., and W.M. Beck, Jr. 1969. Chironomidae (Diptera) of Florida. III. The Harnischia complex (Chironominae). Bull. Fla. St. Mus. 13(5):277-313.
- Beck, W.M., Jr., and E.C. Beck. 1966. Chironomidae (Diptera) of Florida: I. Pentaneurini (Tanypodinae). Bull. Fla. St. Mus. 10(8):305-379.
- Boesel, M.W. 1974. Observations on the Coelotanypodini of the northeastern states, with keys to the known stages (Diptera:Chironomidae:Tanypodinae). J. Kans. Ent. Soc. 47(4):417-432.
- Brinkhurst, R.O., and B.G.M. Jamieson. 1971. Aquatic Oligochaeta of the world. University of Toronto Press, Toronto. 860 pp.
- Brown, H.P. 1972. Biota of freshwater ecosystems identification manual No. 6. Aquatic dryopoid beetles (Coleoptera) of the United States. U.S. Government Printing Office, Washington, D.C. 82 pp.
- Burch, J.B. 1973. Biota of freshwater ecosystems identification manual No. 11. Freshwater unionacean clams (Mollusca:Pelecypoda) of North America. U.S. Government Printing Office, Washington, D.C. 176 pp.

- Burks, B.D. 1975. The mayflies, or Ephemeroptera, of Illinois. [First published in 1953 as Illinois Nat. Hist. Survey, Bull. 26(1)]. Reprinted by Entomological Reprint Specialists, Los Angeles, California. 216 pp.
- Carr, J.F., and J.K. Hiltunen. 1965. Changes in the bottom fauna of western Lake Erie from 1930 to 1961. *Limnol. and Oceanogr.* 10(4):551-569.
- Chutter, F.M. 1969. The effects of silt and sand on the invertebrate fauna of streams and rivers. *Hydrobiologia* 34(1):57-76.
- Curry, L.L. 1958. Larvae and pupae of the species of Cryptochironomus (Diptera) in Michigan. *Limnol. and Oceanogr.* 3(4):427-442.
- deMarch, B.G.E. 1976. Spatial and temporal patterns in macrobenthic stream diversity. *J. Fish. Res. Board Can.* 33:1261-1270.
- Edmondson, W.T. (editor). 1959. Freshwater biology. Second edition. John Wiley and Sons, New York, New York. 1248 pp.
- Elliot, J.M. 1971. Some methods for the statistical analysis of samples of benthic invertebrates. Freshwater Biological Association Scientific Publication No. 25. Ambleside, Westmorland, U.K. 144 pp.
- Foster, N. 1972. Biota of freshwater ecosystems manual No. 4. Freshwater polychaetes (Annelida) of North America. U.S. Government Printing Office, Washington, D.C. 15 pp.
- Grant, P.R., and R.J. Mackay. 1969. Ecological segregation of systematically related stream insects. *Can. J. Zool.* 47:691-694.
- Harman, W.N. 1974. Snails (Mollusca:Gastropoda), pp. 275-312. In C.W. Hart Jr. and S.L.H. Fuller (editors). Pollution ecology of freshwater invertebrates. Academic Press, Inc., New York, New York.
- Harman, W.N., and C.O. Berg. 1971. The freshwater snails of central New York with illustrated keys to the genera and species. Cornell Univ. Ag. Exp. Sta., Ithaca, New York 1(4):1-68.
- Hilsenhoff, W.L. 1975. Aquatic insects of Wisconsin. Technical Bull. No. 89. Dept. of Natural Resources, Madison, Wisconsin. 53 pp.
- Hiltunen, J.K. 1965. Distribution and abundance of the polychaete, Manayunkia speciosa Leidy, in western Lake Erie. *Ohio. J. Sci.* 65(4):183-185.
- _____. 1973. Keys to the tubificid and naidid Oligochaeta of the Great Lakes region. Second edition. Great Lakes Fishery Laboratory, Ann Arbor, Michigan. 25 pp.

1565 151

POOR ORIGINAL

- Holsinger, J.R. 1972. Biota of freshwater ecosystems identification manual No. 5. The freshwater amphipod crustaceans (Gammaridae) of North America. U.S. Government Printing Office, Washington, D.C. 89 pp.
- Hoopes, R.L. 1974. Flooding, as the result of Hurricane Agnes; and its effect on a macrobenthic community in an infertile headwater stream in central Pennsylvania. *Limnol. and Oceanogr.* 19(5):853-857.
- Hynes, H.B.N. 1972. The ecology of running waters. University of Toronto Press, Toronto. 555 pp.
- Ischinger, L.S., and T.F. Nalepa. 1966. Water Pollution: Freshwater Macroinvertebrates. *J. Water Pollut. Contr. Fed.* 48(6):1318-1335.
- Johannsen, O.A. 1969. Aquatic Diptera. (First published in 1934, 1935, 1937, and 1937 as Parts I through IV. *Memoirs* 164, 177, 205, and 210 Cornell Univ. Exp. Station). Reprinted by Entomological Reprint Specialists, Los Angeles, California. 369 pp.
- Kennedy, C.R. 1966. The life history of Limnodrilus hoffmeisteri Clap. (Oligochaeta: Tubificidae) and its adaptive significance. *Oikos* 17:158-168.
- Kennedy, J.H. 1975. Macroinvertebrates. pp. 339-386. In W.A. Potter and Associates. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Annual Report for 1974. Ichthyological Associates, Inc. 468 pp.
- Klemm, D.J. 1972. Biota of freshwater ecosystems identification manual No. 8. Freshwater leeches (Annelida: Hirudinea) of North America. U.S. Government Printing Office, Washington, D.C. 53 pp.
- _____. 1975. Studies on the feeding relationships of leeches (Annelida: Hirudinea) as natural associates of mollusks. *Sterkiana* (58):1-50; (59):1-20.
- _____. 1976. Leeches (Annelida: Hirudinea) found in North America mollusks. *Malac. Rev.* 9:63-76.
- Lloyd, M., J.H. Zar, and J.R. Karr. 1968. On the calculation of information-theoretical measures of diversity. *Amer. Midl. Nat.* 79(7):257-272.
- Loden, M.S. 1976. Life history patterns of naidid oligochaetes in Indiana streams. Paper presented at the 24th meeting of North American Benthological Society, LaCrosse, Wisconsin, March 24-26, 1976.
- Maloney, S.D., and C.M. Chandler. 1976. Leeches (Hirudinea) in the upper Stones River drainage of middle Tennessee. *Amer. Midl. Nat.* 95(1): 42-48.

- Mason, W.T. 1973. An introduction to the identification of chironomid larvae. Environmental Protection Agency, Cincinnati, Ohio. 90 pp.
- Mason, W.T., and P.P. Yevich. 1967. The use of phloxine B and rose bengal stains to facilitate sorting benthic samples. Trans. Amer. Micros. Soc. 86(2):221-223.
- Massengill, R.R. 1976. Benthic fauna:1965-1967 versus 1968-1972 pp. 39-53. In D. Merriman and L.M. Thorpe. The Connecticut River Study. The Impact of a Nuclear Power Plant. Am. Fish. Soc. Monogr. 1. 252 pp.
- Morse, J.C., and I.D. Wallace. 1975. Athripsodes Billberg and Ceraclea Stephens, distant genera of long-horned caddis-flies (Trichoptera: Leptoceridae). Proc. of the First Int. Symp. on Trichoptera. Junk. The Hague. pp. 33-40.
- Needham, J.G., and M.J. Westfall, Jr. 1955. A manual of the dragonflies of North America (Anisoptera), including the Greater Antilles and the provinces of the Mexican border. Univ. of California Press, Berkeley and Los Angeles. 615 pp.
- Poe, T.P., and D.C. Stefan. 1974. Several environmental factors influencing the distribution of the fresh-water polychaete, Manayunkia speciosa Leidy. Chesapeake Sci. 15(4):235-237.
- Polk, J.L., and J.H. Epler. 1976. Macroinvertebrates. pp. 241-298. In W.A. Potter and Associates. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Annual Report for 1975. Ichthyological Associates, Inc. 395 pp.
- Polk, J.L., J.H. Kennedy, W.A. Potter, and J.H. Epler. 1976. The effects of flooding on populations of Chironomus Meigen (Diptera:Chironomidae) and Limnodrilus Claparede (Oligochaeta:Tubificidae) in the Susquehanna River in the vicinity of Three Mile Island, Pennsylvania. Proc. Pa. Acad. Sci. 50(1):91-95.
- Reid, G.K. 1961. Ecology of inland waters and estuaries. Van Nostrand Reinhold Co., New York, New York. 375 pp.
- Reynolds, J.W. 1975. Sparganophilus pearsei n. sp. (Oligochaeta:Sparganophilidae) a nearctic earthworm from western North Carolina. Megadrilologica. 2(2):9-11.
- Roback, S.S. 1957. The immature tendipedids of the Philadelphia area. Monogr. Acad. Nat. Sci. Philad. No. 9. 180 pp.
- _____. 1971. The adults of the subfamily Tanypodinae (= Pelopiinae) in North America (Diptera:Chironomidae). Monogr. Acad. Nat. Sci. Philad. No. 17. 410 pp.

- _____. 1974a. Insects (Arthropoda:Insecta), pp. 313-376. In C.W. Hart, Jr. and S.L.H. Fuller (editors). Pollution ecology of freshwater invertebrates. Academic Press, Inc., New York, New York.
- _____. 1974b. The immature stages of the genus Coelotanypus (Chironomidae: Tanypodinae:Coelotanypodini) in North America. Proc. Acad. Nat. Sci. Philad. 126(2):9-19.
- _____. 1976. The immature chironomids of the eastern United States. I. Introduction and Tany, Tiniinae-Coelotanypodini. Proc. Acad. Nat. Sci. Philad. 127(14):147-210.
- Ross, H.H. 1972. The caddis flies, or Trichoptera, of Illinois. [First published in 1944 as Illinois Nat. Hist. Survey, Bull. 23(1).] Reprinted by Entomological Reprint Specialists, Los Angeles, California. 326 pp.
- Saether, O.A. 1975. Nearctic and Palaearctic Heterotrissocladius (Diptera: Chironomidae). Bull. Fish. Res. Board Can. 193. 65 pp.
- _____. 1976. Keys to larvae and pupae of Orthocladiinae and Telmatogetoninae. 61 pp. (Unpublished).
- Sawyer, R.T. 1972. North American freshwater leeches, exclusive of the Piscicolidae, with a key to all species. Illinois Bio. Monogr. 46. 154 pp.
- _____. 1974. Leeches (Annelida:Hirudinea) pp. 81-142. In C.W. Hart, Jr. and S.L.H. Fuller (editors). Pollution ecology of freshwater invertebrates. Academic Press, Inc., New York, New York.
- Sokal, R.R., and F.J. Rohlf. 1969. Biometry, the principles and practice of statistics in biological research. W.H. Freeman, San Francisco. 776 pp.
- Spencer, D.R. 1976. Occurrence of Manavunkia speciosa (Polychaeta:Sabellidae) in Cayuga Lake, New York, with additional notes on its North American distribution. Trans. Amer. Micros. Soc. 95(1):127-128.
- Sublette, J.E., and M.F. Sublette. 1974. A review of the genus Chironomus (Diptera:Chironomidae) V. The maturus-complex. Stud. Nat. Sci. 1(8):1-42.
- Tebo, L.B., Jr. 1970. Effects of siltation, resulting from improper logging, on the bottom fauna of a small trout stream in the southern Appalachians. The Prog. Fish-Cul. 17(1955):64-70.
- Usinger, R.L. (editor). 1956. Aquatic insects of California with keys to North American genera and California species. Univ. of California Press. Berkeley and Los Angeles. 508 pp.

- Walker, E.M. 1958. The Odonata of Canada and Alaska. Volume Two. Part III: the Anisoptera - four families. Univ. of Toronto Press, Toronto. 318 pp.
- Walker, E.M., and P.S. Corbet. 1975. The Odonata of Canada and Alaska. Volume Three. Part III: the Anisoptera - three families. Univ. of Toronto Press, Toronto and Buffalo. 307 pp.
- Whittaker, R.H., and T.W. Fairbanks. 1958. A study of plankton copepod communities in the Columbia Basin, southeastern Washington. Ecology 39:46-65.
- Wilhm, J.L. 1970. Range of diversity index in benthic macroinvertebrate populations. J. Water Pollut. Contr. Fed. 42(5) Part 2:221-224.
- Woolf, C.M. 1968. Principles of biometry. Van Nostrand Co., Ltd., Toronto, Canada. 359 pp.

1565 155

Table 3.1-1

Location and description of macroinvertebrate stations.

Station Number	Location and Description
TM-AQI-1A1*	40° 09' 52" N, 76° 43' 26" W. North end of Sand Beach Island, 30 to 75 m offshore. Water depth varied from 0.5 to 2.0 m. Substrate composed of sand, coal particles, and detritus; sometimes with mud and/or clay. Trace amounts of oil sometimes present.
TM-AQI-1A2	40° 09' 36" N, 76° 43' 30" W. Southwest St. Johns Island, 1 to 15 m offshore at mouth of channel between TMI and St. Johns Island. Water depth varied from 0.5 to 1.5 m. Substrate composed mostly of sand, coal particles, and detritus; sometimes with gravel or clay. Trace amounts of oil sometimes present.
TM-AQI-11A1	40° 09' 09" N, 76° 43' 39" W. 1 to 10 m downstream from TMI Discharge, 1 to 15 m offshore. Water depth varied from 0.25 to 1.0 m. Substrate composed of mud mixed with coal particles, fine sand, and detritus; sometimes with muck, clay, or gravel. Trace amounts of oil present.
TM-AQI-11A2	40° 09' 07" N, 76° 43' 39" W. 70 to 75 m downstream from TMI Discharge, 1 to 15 m offshore. Water depth varied from 0.25 to 1.5 m. Substrate composed of mud with fine sand, some coal particles, and detritus; sometimes with clay. Trace amounts of oil present.
TM-AQI-9B1	40° 08' 03" N, 76° 43' 33" W. 1900 m downstream from TMI Discharge, 5 to 15 m offshore. Water depth varied from 0.75 to 1.5 m. Substrate composed of mud with muck, detritus, and fine sand, with some coal particles. Trace amounts of oil present.

*Polar coordinate prefix TM-AQI- deleted from all station numbers for discussion in text.

Macroinvertebrates taken from the Susquehanna River in the vicinity of Three Mile Island.

Coelenterata	Nirudinea
Hydrozoa	Rhynchobdellida
Hydroida	Clossiphoniidae
Hydridae	<i>Melobdella elongata</i> (Castle)
<i>Hydra</i> sp.	<i>H. lineata</i> (Verrill)
Lymnemedusa	<i>Placobdella manitifica</i> Moore
Olinidiidae	<i>Z. parasitica</i> (Say)
<i>Craspedacusta sowerbyi</i> Lankester	Piscicolidae
Platyhelminthes	<i>Microbdella lugubris</i> Leidy
Turbellaria	Pharyngobdellida
Tricladida	Erpobdellidae
Planariidae	<i>Erpobdella punctata</i> (Leidy)
<i>Dugesia tigrina</i> (Girard)	<i>Microbdella microstoma</i> (Moore)
Nemertinea	Arthropoda
Enopla	Arachnida
Hoploneurini	Araneae
Tetrastemmatidae	Fissauridae
<i>Prostoma rubrum</i> (Leidy)	<i>Polonides triton</i> (Walckenaer)
Nematoda	Hydracarina
Acanthocephala	Crustacea
Palaecanthocephala	Copepoda
Echinorhynchida	Lernaeidae
<i>Lentorhynchoides thersites</i> (Linton)	<i>Lernaea</i> sp.
Entoprocta	Branchiura
Urnatellida	<i>Argulus catostomi</i> Dana and Herrick
Urnatellidae	Isopoda
<i>Urnatella gracilis</i> Leidy	Asellidae
Bryozoa	<i>Asellus communis</i> Say
Phylactolaemata	Amphipoda
Lophopodidae	Gammaridae
<i>Lophopodella carteri</i> (Hyatt)	<i>Gammarus</i> sp.
<i>Pectinatella magnifica</i> (Leidy)	<i>Gammarus fasciatus</i> Say
Plumatellidae	Decapoda
<i>Hyalinella punctata</i> (Hancock)	Astacidae
<i>Plumatella repens</i> (Linnaeus)	<i>Orconectes obscurus</i> (Hagen)
Cynolaemata	<i>O. rusticus</i> (Girard)
Ctenostomida	Insecta
Paludicellidae	Collembola
<i>Paludicella arctica</i> ? (Ehrenberg)	Ephemeroptera
Amelida	Siphonuridae
Polychaeta	<i>Isomura</i> sp.
Errantia	<i>Siphonurus</i> sp.
Scabellidae	Baetidae
<i>Manayunkia speciosa</i> Leidy	<i>Baetis</i> sp.
Oligochaeta	Heptageniidae
Plesiopora	<i>Stenocranus carolina</i> (Banks)
Enchytraeidae	<i>S. interruptum</i> (Say)
Naididae	<i>Stenocranus ares</i> Burks
<i>Arctonais lomondi</i> (Martin)	<i>S. pulchellum</i> (Walsh)
<i>Aulophorus furcatus</i> (Müller)	Leptophlebiidae
<i>Chaetogaster diaphanus</i> (Gruithuisen)	<i>Leptophlebia</i> sp.
<i>Nais brethescheri</i> Michaelson	Ephemereidae
<i>N. elongata</i> Müller	<i>Ephemerella aestiva</i> McDunnough
<i>N. variabilis</i> Piquet	<i>E. serrata</i> group sp.
<i>Ophidionis serpentina</i> (Müller)	<i>E. variabilis</i> McDunnough
<i>Paranais iridi</i> Hrab	Tricorythidae
<i>Slavina appendiculata</i> (d'Udekem)	<i>Tricorythodes</i> sp.
Tubificidae	Caenidae
<i>Aulodrilus pluriseta</i> (Piquet)	<i>Caenis riggsi</i> Burks
<i>Branchiura sowerbyi</i> Reddard	Potamanthidae
<i>Hydrodrilus completioni</i> (Southern)	<i>Potamanthus</i> sp.
<i>Limnodrilus cervix</i> Brinkhurst	Ephemeridae
<i>L. claredei</i> Ratzei	<i>Hemaphysalis limba</i> (Serville)
<i>L. hoffmeisteri</i> Claparède	<i>H. riggsi</i> McDunnough
<i>L. spiralis</i> (Eisen)	Baetiscidae
<i>L. udekemianus</i> Claparède	<i>Baetisca</i> sp.
<i>Poloscolex lerox</i> (Eisen)	Odonata
<i>P. multisetosus</i> (Smith)	Aeshnidae
<i>Tubifex tubifex</i> (Müller)	<i>Anax junius</i> (Drury)
Trochopora	Comptidae
Lumbricidae	<i>Comptosia spinosus</i> Selys
Branchiobdellida	<i>Comptosia (Artemesia) villosipes</i> Selys
Spisthopora	<i>C. (Comptosia) variegata</i> Walsh
Sparganophilidae	<i>C. (Comptosia) livida</i> Selys
<i>Sparganophilus eiseni</i> Smith	<i>C. (Artemesia) spiniceps</i> (Walsh)
	Macronitidae
	<i>Macronia illinoensis</i> Walsh
	Corduliidae
	<i>Epitheca pringens</i> Hagen

1565 157

POOR ORIGINAL

Table 3.1-2 continued.

Arthropoda - continued

Insecta - continued

Odonata - continued

Libellulidae

- Libellula luctuosa Burmeister
L. ludia Drury
L. pulchella Drury
Pachydiplax longipennis (Burmeister)
Pantala flavescens (Fabricius)
P. lunata (Say)
Perithemis tenera (Say)

Calopterygidae

- Metaeterna americana (Fabricius)

Coenagrionidae

- Argia apicalis (Say)
A. rufa (Hagen)
A. translata (Hagen)
Enallagma civile (Hagen)
E. esculans (Hagen)
Ischnura verticalis (Say)

Plecoptera

Pteronarcidae

- Pteronarcys sp.

Taeniopterygidae

- Brachyptera fasciata (Burmeister)
Taeniopteryx sp.

Nemouridae

- Nemoura delosa Ricker

Capniidae

- Allocaenia sp.

Perlidae

- Acronuria sp.
Perlusia placida (Hagen)

Perlodidae

- Isoperla bilineata (Say)

Hemiptera

Corixidae

- Sigara alternata (Say)
Trichocorixa calva (Say)

Notonectidae

- Notonecta undulata Say

Belostomatidae

- Belostoma sp.

Nepidae

- Nepa nigris Herrich-Schaeffer

Gelastocoridae

- Gelastocoris oculatus (Fabricius)

Cerridae

- Cerris argenticollis Parshley
C. confinis (Uhler)
Metrobates hesperius Uhler
Rheumatobates gilvi Bergroth

Veliidae

- Microvelia americana (Uhler)
Chagovelia obesa Uhler

Mesoveliidae

- Mesovelia mulsanti White

Saldidae

- Pentacora ligata (Say)
Saldula sp.

Megaloptera

Sialidae

- Sialis sp.

Corydalidae

- Corydalus cornutus (Linnaeus)

Neuroptera

Sisyridae

- Climacia areolaris (Hagen)

Trichoptera

Psychomyiidae

- Psychomyia flavida Hagen

Polycentropodidae

- Neureclipsis sp.
Polycentropus ciferri Hagen

Hydropsychidae

- Cheumatopsyche campyla Ross
Hydropsyche atrata Ross
H. bifida group sp.
H. morosa Hagen
H. phalerata Hagen
Macronema carolina Banks
M. zebra Hagen

Glossosomatidae

- Protophila palina (Ross)

Hydroptilidae

- Hydroptila spatulata Norton
H. wickhami Letten

Limnephilidae

- genus nr. Trisialis Letten
Prempavola subulata (Say)

Leptoceridae

- Ceraclea pennellata (Letten)
C. flava (Letten)
C. tarsitarsata (Letten)
Leptocera sp.
Metacera sp.
Metacera sp.

Lepidoptera

Pyralidae

- Parapala sp.
Parapala sp.

Coleoptera

Halipidae

- Halipia sp.
Feltonia sp.

Dytiscidae

- Copelatus sp.
Laccophilus sp.

Gyrinidae

- Dineutus discolor Aubé
D. normis Roberts
Gyrinus sp.

Hydrophilidae

- Hermes sp.
Isopisthus sp.
I. nigratus (Say)

Psephenidae

- Psephenus sp.

Elmidae

- Amblyderus sp.
Elmidae sp.
E. vittata (Melsheimer)
Macronema sp.
Oniscus sp.
Oniscus sp.
Oniscus sp.
Oniscus sp.
Oniscus sp.
Oniscus sp.

Chrysomelidae

- Donacia sp.

Diptera

Tipulidae

- Anopheles sp.
Hematomia sp.
Tipula sp.

Psychodidae

- Pericoma sp.
Psychoda sp.
Telescopus sp.

Culicidae

- Aedes sp.

Chaoboridae

- Chaoborus sp.

Chironomidae

Tanyptinidae

- Tanypterus sp.
Procladius sp.
Psectrocladius sp.
Coelotanypterus sp.
C. scutellaris (Loew)
Ablabesmus sp.
A. n. nigratus (Loew)
Thienemannia group sp.

Diametidae

- Diametia sp.

Orthocladidae

- Brillia sp.
Cricotopus sp.
C. silvestris group sp.
Cricotopus sp.
Eukiefferiella sp.
Heterotanypterus sp.
Orthocladus sp.
Orthocladus sp.
Parametoponema sp.
Psectrocladius sp.
Smittia sp.
 genus nr. Leptocladus Steffer

Table 3.1-2 continued.

Arthropoda - continued	Mollusca
Insecta - continued	Gastropoda
Diptera - continued	Basommatophora
Chironomidae - continued	Physidae
Chironominae	<i>Physa</i> sp.
<i>Chironomus texensis</i> Johannsen	Lymnaeidae
<i>Cryptochironomus</i> nr. <i>blarina</i> Townes	<i>Lymnaea humilis</i> Say
<i>C.</i> nr. <i>flavus</i> (Johannsen)	Planorbidae
<i>Pemphrochironomus vulneratus</i> Zetterstedt	<i>Cerastus parvus</i> (Say)
<i>Dicrotendipes modestus</i> (Say)	<i>Helisoma triolvis</i> (Say)
<i>D. normus</i> (Staeger)	Ancylida
<i>Endochironomus</i> nr. <i>tendens</i> (Fabricius)	<i>Ferrissia rivularis</i> (Say)
<i>Glyptotendipes</i> sp.	<i>F. sarda</i> (Say)
<i>Harnischia</i> nr. <i>spachensis</i> Townes	Mesogastropoda
<i>Parachironomus</i> sp.	Pleuroceridae
<i>Paracladopelma</i> sp.	<i>Coniothis virginica</i> (Lacini)
<i>Paratendipes</i> sp.	<i>Spirodon carinata</i> (Lorquiere)
<i>Phaenopsectra</i> (<i>Sergentia</i>) nr. <i>obediens</i> (Johannsen)	Valvatidae
<i>Phaenopsectra</i> (<i>Tribellog</i>) sp.	<i>Valvata tricarinata</i> (Say)
<i>Polypedilum ralla</i> group sp.	Hydrobiidae
<i>P. helveticum</i> (Coquillett)	<i>Arnicola limosa</i> (Say)
<i>P.</i> nr. <i>illinoense</i> (Malloch)	<i>Sithonia tentaculata</i> (Linnaeus)
<i>P. scalae</i> (Schränk)	Viviparidae
<i>Stenochironomus</i> sp.	<i>Campeloma decisa</i> (Say)
<i>Scaptotendipes</i> sp.	Pelecypoda
<i>Rheotanytarsus</i> nr. <i>exiguus</i> (Johannsen)	Eulamellibranchia
<i>Tanytarsus</i> nr. <i>huckleyi</i> Sublette	Unionidae
<i>Tanytarsus</i> spp.	<i>Anodonta cataraeta</i> (Say)
Ceratopogonidae	<i>Lilipia complanata</i> (Lightfoot)
<i>Palpomyia</i> group sp.	<i>Lamprellia</i> sp.
Simuliidae	Heterodonta
<i>Simulium</i> (<i>Palloria</i>) <i>vittatum</i> Zetterstedt	Sphaeriidae
Tabanidae	<i>Pisidium</i> sp.
<i>Chrysops</i> sp.	<i>Sphaerium</i> sp.
Empididae	
<i>Hemerodromia</i> sp.	
Dolichopodidae	
Ephydriidae	
Muscidae	
<i>Fannia</i> sp.	
<i>Lispe</i> sp.	

POOR ORIGINAL

1565 159

Table 3.2-1

Numbers and milligrams of biomass, = (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQ1-1A1, April 1976. Hashes indicate species not present or no weight measurement made.

Date	6 April				20 April			
Time	0935				0940			
Air Temp. (C)	13.5				25.5			
Water Temp. (C)	8.0				20.0			
Dissolved Oxygen (ppm)	9.8				8.4			
pH	7.6				8.0			
Secchi Disc (cm)	43.2				127.0			
River Stage (ft)	6.6				4.7			
Substrate	Sand, Coal, Mud, Detritus				Sand, Coal, Detritus, Mud			
Replicate	A	B	C	D	A	B	C	D
Nemertinea	-	1(-)	1(-)	2(-)	-	1(-)	-	-
Nematoda	-	-	1(-)	1(-)	1(-)	2(-)	1(-)	1(-)
Enchytraeidae	1(-)	-	1(-)	1(-)	-	-	-	-
<u>Nais breetscheri</u>	1(-)	1(-)	-	3(-)	-	-	-	-
<u>N. glinguis</u>	-	-	-	-	3(-)	1(-)	4(-)	-
<u>N. variabilis</u>	-	-	1(-)	1(-)	-	-	-	1(-)
<u>Branchiura sowerbyi</u>	-	-	-	-	1(0.6)	1(1.0)	-	1(3.5)
<u>Limnodrilus clapparedianus</u>	2(-)	-	3(-)	1(-)	10(5.9)	10(5.1)	-	-
<u>L. hoffmeisteri</u>	-	1(-)	2(-)	4(-)	-	-	12(7.0)	32(14.6)
<u>L. udekemianus</u>	-	-	1(-)	-	-	-	-	-
<u>Limnodrilus</u> spp.	4(-)	10(-)	6(-)	17(-)	19(11.2)	29(14.9)	38(22.1)	29(17.8)
<u>Peloscoides multisetosus</u>	1(-)	1(-)	-	1(-)	1(-)	-	-	-
<u>Tubificoides tubificoides</u>	-	-	-	-	-	1(-)	-	-
Imm. tub./cap. chaetae*	-	1(-)	1(-)	2(-)	-	-	3(-)	8(3.6)
Lumbriculidae	-	-	-	1(-)	-	-	1(-)	-
Megadrile	-	-	-	-	-	-	1(-)	1(-)
<u>Oecetis</u> sp.	1(-)	-	-	-	-	-	-	-
<u>Psychoda</u> sp. pupa	-	-	-	1(0.7)	-	-	-	-
<u>Procladius</u> sp.	-	-	1(-)	-	-	-	-	-
<u>Cricotopus</u> spp.	-	-	1(-)	-	-	-	-	-
<u>Cricotopus</u> spp. pupa	-	-	2(-)	-	-	-	-	-
<u>Parametriocnemus</u> sp.	-	-	1(-)	-	-	-	-	-
<u>Chironomus decorus</u>	-	-	-	-	1(-)	-	1(-)	-
<u>Phaenopsectra</u> nr. <u>obediens</u>	-	-	-	-	1(-)	-	-	4(-)
<u>Polypodium fallax</u> group sp.	-	-	-	-	-	-	-	1(-)
<u>Ferrissia</u> spp.	1(-)	-	-	-	-	-	-	-
<u>Contobasis virginica</u>	4(17.2)	-	6(82.3)	1(9.1)	7(103.2)	-	3(8.6)	-
<u>Pisidium</u> sp.	-	17(-)	16(-)	12(-)	5(-)	17(-)	4(-)	-
<u>Sphaerium</u> sp.	-	4(5.0)	12(-)	3(-)	-	2(-)	1(-)	3(-)

* Immature tubificid with capilliform chaetae.

1565 160

Table 3.2-1

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-1A1, May 1976. Dashes indicate species not present or no weight measurement made.

Date	4 May				18 May			
Time	1009				1015			
Air Temp. (C)	NA				19.0			
Water Temp. (C)	11.8				20.0			
Dissolved Oxygen (ppm)	9.0				8.0			
pH	7.8				NA			
Secchi Disc (cm)	66.0				61.9			
River Stage (ft)	4.9				5.6			
Substrate	Mud, Sand, Coal, Some Detritus				Mud, Sand, Coal, Some Detritus			
Sampl. Date	A	B	C	D	A	B	C	D
Nemertinea	-	-	-	-	-	-	1(0.5)	1(-)
Nematoda	12(1.4)	7(0.5)	7(0.3)	7(0.7)	4(-)	4(-)	2(-)	-
<i>Pleurotella repens</i>	PA	-	-	P	-	-	-	-
<i>Chaetogaster diaphanus</i>	-	-	-	-	2(-)	-	-	5(-)
<i>Leis brepseri</i>	12(0.3)	-	-	-	-	-	-	-
<i>S. elongus</i>	115(2.7)	153(2.7)	231(5.4)	138(3.2)	685(16.6)	58(1.2)	205(5.2)	324(8.4)
<i>S. var.abilis</i>	-	24(0.4)	24(0.6)	10(0.2)	59(1.4)	-	11(0.3)	-
<i>Chironomus serpentina</i>	-	-	-	-	30(0.7)	-	-	-
<i>Paranais trici</i>	-	35(0.6)	-	-	3(-)	-	-	15(0.4)
<i>Branchiura sowerbi</i>	1(-)	-	1(0.6)	-	-	-	-	-
<i>Chironomus tentationi</i>	-	-	-	-	-	-	8(2.3)	-
<i>Chironomus claretedelanus</i>	-	-	-	-	20(2.6)	-	-	-
<i>S. hoffmeisteri</i>	32(7.7)	54(2.1)	5(0.7)	10(1.5)	41(5.3)	11(2.2)	21(4.2)	65(13.3)
<i>S. adriaticus</i>	-	14(2.1)	-	-	-	-	-	-
<i>Chironomus</i> spp.	128(30.9)	82(12.3)	128(18.0)	121(18.6)	183(23.7)	69(13.9)	127(25.1)	43(9.2)
<i>Polyscolex multisetosus</i>	2(4.0)	-	-	2(0.7)	-	-	4(0.9)	3(1.2)
<i>Limn. subifex</i>	5(0.9)	13(2.6)	8(2.4)	-	44(9.2)	-	8(2.3)	17(3.1)
Imm. tub./cap. chaetae**	18(3.3)	21(4.2)	32(9.8)	-	52(10.8)	13(-)	31(8.9)	17(3.1)
<i>Magadride</i>	2(-)	1(-)	-	-	-	-	-	-
<i>Polychaeta lineata</i>	-	1(1.9)	-	-	-	-	-	-
<i>Polychaeta</i> sp.	-	-	-	-	-	-	1(-)	-
<i>Hydracarina</i>	1(-)	1(-)	-	-	-	-	-	-
<i>Stellus communis</i>	-	-	-	-	-	-	1(0.2)	-
<i>Limn. fasciatus</i>	-	1(-)	1(-)	-	4(0.5)	3(0.4)	4(1.1)	4(1.1)
<i>Limn. psyllus</i> sp.	1(-)	3(1.0)	1(0.8)	4(2.2)	-	-	-	-
<i>Limn. psyllus phalerata</i>	-	-	-	1(1.1)	-	-	-	-
<i>Limn. psyllus</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Limn. psyllus</i> sp.	1(-)	-	-	-	-	-	-	-
<i>Limn. psyllus</i> sp.	-	-	-	-	3(2.7)	2(-)	2(-)	2(-)
<i>Limn. psyllus</i> sp.	-	2(-)	-	-	-	-	-	-
<i>Limn. psyllus</i> spp.	-	5(0.1)	1(-)	3(0.3)	-	1(-)	-	-
<i>Limn. psyllus</i> spp. pupa	-	1(-)	2(-)	-	-	-	-	-
<i>Limn. psyllus</i> sp. A	1(-)	-	-	-	-	-	-	-
<i>Orthocladus (Orthocladus) sp.</i>	-	-	-	3(0.3)	-	-	-	-
<i>Orthocladus (Orthocladus) sp.</i>	1(-)	9(0.2)	18(0.5)	9(0.8)	-	2(-)	3(-)	-
<i>Orthocladus (Orthocladus) sp. pupa</i>	-	-	-	2(-)	-	1(-)	-	-
<i>Orthocladus</i> sp.	-	4(0.1)	1(-)	-	-	-	-	1(-)
<i>Limn. decorus</i>	-	-	2(-)	-	3(-)	5(-)	2(-)	2(-)
<i>S. decorus</i> pupa	1(-)	-	-	-	-	-	-	-
<i>Limn. decorus</i> sp.	-	1(-)	-	-	-	-	-	-
<i>Limn. decorus</i> nr. <i>obediens</i>	-	-	-	-	4(0.8)	1(-)	3(-)	2(-)
<i>Limn. decorus fallax</i> pupa	-	1(-)	-	-	-	-	-	-
<i>S. nr. illinoense</i>	3(-)	-	12(0.3)	-	-	-	-	-
<i>Tanytarsini</i> pupa	-	1(-)	-	-	-	-	-	-
<i>Tanytarsus</i> sp.	-	1(-)	-	-	-	1(-)	-	-
<i>Limn. group</i> sp.	-	-	-	-	-	-	1(-)	-
<i>Limn. sp.</i>	-	-	1(-)	-	-	-	-	-
<i>Limn. virginica</i>	2(9.9)	-	-	-	1(20.3)	-	-	1(30.7)
<i>Limn. sp.</i>	1(-)	-	2(-)	-	7(5.0)	-	2(1.0)	2(2.6)
<i>Limn. sp.</i>	-	2(-)	1(-)	-	-	-	1(-)	3(5.2)

NA = Not Available.

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

Table 3.2-3

Numbers and milligrams of biomass, μ (mg), of benthic organisms per Ponar grab (525 cm²) at Station TM-AQ1-1A1, June 1976. Dashes indicate species not present or no weight measurement data.

Date	1 June				15 June			
Time	0627				0751			
Air Temp. (C)	22.0				22.3			
Water Temp. (C)	19.0				22.5			
Dissolved Oxygen (ppm)	9.2				7.9			
pH	8.0				7.8			
Seichi Disc (cm)	76.2				76.2			
River Stage (ft)	5.1				4.1			
Substrate	Mud, Sand, Coal, Detritus				Mud, Sand, Coal, Detritus			
Sp. Code	A	B	C	D	A	B	C	D
<i>Neotoma</i>	1(-)	13(1.9)	22(1.9)	19(1.6)	-	3(-)	-	-
<i>Planorbella repens</i>	0*	-	-	-	-	-	-	-
<i>Arctonasis formosa</i>	1(-)	-	-	10(0.1)	-	-	-	-
<i>Mais brevis</i>	-	-	1(-)	13(0.1)	-	-	1(-)	-
<i>S. elongata</i>	2(-)	-	2(-)	14(0.2)	-	-	-	-
<i>S. variabilis</i>	1(-)	-	-	-	-	-	-	-
<i>Paranais trica</i>	1(-)	-	1(-)	13(0.1)	-	-	-	-
<i>Branchinella sowerbyi</i>	-	-	1(2.3)	-	2(-)	-	-	-
<i>Hydrilus teretifolius</i>	-	-	-	-	1(-)	1(-)	1(-)	-
<i>Limnocalanus macrurus</i>	-	149(28.4)	-	-	30(13.7)	12(3.9)	10(3.1)	-
<i>L. hoffmeisteri</i>	59(18.3)	256(48.8)	124(25.5)	116(16.8)	20(9.2)	59(19.1)	29(11.6)	36(6.6)
<i>L. vancouverianus</i>	10(3.1)	-	-	-	-	-	-	-
<i>Limnocalanus</i> spp.	69(21.4)	106(20.2)	162(33.3)	125(18.2)	39(17.8)	47(15.2)	40(11.5)	60(11.0)
<i>Polyscolus multisetosus</i>	-	2(0.4)	1(-)	-	-	-	-	-
<i>Tubificoides tubificus</i>	3(0.3)	6(0.9)	2(-)	13(2.3)	1(-)	3(-)	1(-)	1(-)
imm. tub./cap. chaetae**	11(1.2)	6(0.9)	6(-)	7(1.2)	8(-)	6(-)	-	3(-)
<i>Hydracarina</i>	1(-)	-	-	-	-	-	-	-
<i>Gammarus fasciatus</i>	1(0.4)	1(0.2)	-	-	1(1.1)	5(0.5)	13(1.3)	2(0.3)
<i>Ephemera serrata</i> group sp.	-	-	-	1(-)	-	-	-	-
<i>Cheumatopsyche</i> sp.	-	1(0.3)	1(0.1)	-	-	-	-	-
<i>Procladius</i> sp.	2(-)	4(0.2)	1(-)	1(-)	-	-	-	-
<i>Psectrocladius</i> sp.	-	-	-	1(-)	-	-	-	-
<i>Triclotopus sylvestris</i> group sp.	-	-	-	2(-)	-	-	-	-
<i>Triclotopus</i> spp.	-	4(0.2)	8(0.5)	1(-)	-	-	-	-
<i>Triclotopus</i> spp. pupa	-	-	1(-)	-	-	-	-	-
<i>Orthocladus</i> (Orthocladus) sp. A	-	-	-	1(-)	-	-	-	-
<i>Orthocladus</i> (Orthocladus) sp.	1(-)	-	5(-)	1(-)	-	-	-	-
<i>Psectrocladius</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Chironomus decorus</i>	2(-)	-	3(1.0)	9(0.2)	148(33.1)	71(16.5)	19(19.1)	109(20.7)
<i>C. decorus</i> pupae	-	-	-	-	-	1(-)	1(-)	-
<i>Cryptochironomus</i> nr. <i>blarina</i>	-	-	-	-	-	-	2(0.1)	-
<i>C. nr. fulvus</i>	-	-	2(0.7)	-	17(3.1)	15(2.6)	17(1.8)	17(4.1)
<i>Glyptotendipes</i> sp.	-	2(0.1)	-	-	-	-	-	-
<i>Glyptotendipes</i> sp.	-	1(0.1)	-	-	-	-	-	-
<i>Paratendipes</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Phaenopsectra</i> nr. <i>obedient</i>	1(-)	2(0.1)	3(0.1)	6(1.2)	10(-)	3(-)	21(3.2)	2(0.2)
<i>Polypedilum</i> nr. <i>scalaeum</i>	-	-	-	-	1(-)	5(-)	-	2(0.2)
<i>Microchironomus</i> sp.	-	-	-	-	-	-	5(0.6)	-
<i>Tanytarsini</i>	-	-	-	1(-)	-	-	-	-
<i>Rheotanytarsus</i> sp.	-	-	-	-	-	1(-)	-	-
<i>Tanytarsus</i> sp.	-	-	-	-	3(-)	2(-)	7(-)	-
<i>Palpomyia</i> group sp.	-	-	-	-	-	1(-)	-	-
<i>Empididae</i>	-	1(-)	-	-	-	-	-	-
<i>Heterodromia</i> sp. pupa	-	-	-	1(-)	-	-	-	-
<i>Gonobasis virginica</i>	1(59.0)	1(6.6)	1(55.0)	2(12.2)	1(38.7)	1(79.7)	4(171.8)	2(43.0)
<i>Pisidium</i> sp.	2(-)	-	1(-)	2(-)	4(3.4)	6(-)	4(-)	-
<i>Sphaerium</i> sp.	1(-)	1(-)	2(1.0)	1(-)	-	5(-)	8(-)	-

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

Fig 3.2-4

Numbers and milligrams of biomass, μ (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-1A1, July 1976. Dashes indicate species not present or no weight measurement made.

Date	6 July				20 July			
Time	0810				0805			
Air Temp. (C)	23.5				23.0			
Water Temp. (C)	24.0				24.0			
Dissolved Oxygen (ppm)	NA				7.4			
pH	8.3				8.0			
Seschi Disc. (cm)	58.4				45.7			
River Stage (ft)	4.7				4.3			
Substrate	Mud, Sand, Coal, Detritus				Mud, Sand, Coal, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	1(-)	-	1(3.6)	-	-	-	-
Nematoda	3(-)	2(-)	-	-	-	1(-)	1(-)	-
<i>Aulophorus furcatus</i>	-	1(-)	-	-	-	-	-	-
<i>Branchiura sowerbyi</i>	-	-	-	-	1(-)	2(2.6)	4(26.4)	3(1.5)
<i>Limnodrilus clapanetianus</i>	21(8.5)	9(3.6)	9(3.0)	49(27.8)	-	11(2.2)	19(5.5)	-
<i>L. hoffmeisteri</i>	31(12.6)	10(4.0)	44(14.6)	29(13.5)	50(18.6)	66(13.4)	47(13.6)	40(22.2)
<i>L. udekemianus</i>	-	-	-	-	-	-	-	10(5.7)
<i>Limnodrilus</i> spp.	135(54.9)	66(26.7)	70(23.3)	118(54.8)	91(33.3)	67(13.6)	76(22.0)	41(23.4)
<i>Pelosciolex multisetosus</i>	-	-	-	2(-)	-	-	-	-
<i>Tubifex tubifex</i>	-	1(-)	1(-)	-	1(-)	-	-	-
Imm. tub./cap. chaetae*	-	1(-)	3(-)	1(-)	2(-)	-	3(-)	-
<i>Sparganophilus eiseni</i>	-	-	-	-	-	-	1(-)	-
Hydracarina	-	-	-	-	-	1(-)	-	-
<i>Gammarus fasciatus</i>	-	4(0.6)	1(0.1)	1(3.5)	1(0.2)	1(-)	2(0.2)	1(-)
<i>Hydropsyche bifida</i> group sp.	-	1(0.3)	-	-	-	-	-	-
<i>H. phalerata</i>	-	1(0.4)	-	3(4.3)	-	-	-	-
<i>Promeseta</i> sp.	-	-	-	1(0.5)	-	-	-	-
<i>Stenelmis decorata</i>	-	2(1.2)	-	-	-	-	-	-
<i>Procladius</i> sp.	4(2.3)	3(-)	-	3(2.8)	-	1(-)	1(-)	-
<i>Ablabesmyia</i> sp.	-	-	-	2(1.9)	-	-	-	-
<i>Cricotopus</i> spp.	-	-	-	-	-	1(-)	-	-
<i>Chironomus decorus</i>	4(2.3)	9(0.4)	2(-)	2(1.9)	3(1.6)	5(-)	7(1.2)	1(-)
<i>Cryptochironomus</i> nr. <i>fulvus</i>	4(2.3)	2(-)	-	5(3.8)	12(5.3)	6(1.8)	5(-)	5(-)
<i>Harnischia</i> nr. <i>amachaerus</i>	-	-	-	-	-	1(-)	-	-
<i>Paratadoneima</i> sp.	1(0.6)	4(0.1)	1(-)	4(6.4)	-	1(-)	1(-)	-
<i>Phaenopsectra</i> nr. <i>obediens</i>	-	14(0.5)	-	-	-	-	-	-
<i>Polypedilum</i> nr. <i>illinoense</i>	-	-	-	2(1.9)	-	-	-	-
<i>P. scalanum</i>	2(1.1)	-	4(-)	8(0.7)	-	2(-)	1(-)	-
<i>Goniobasis virginica</i>	2(29.4)	3(117.5)	1(16.5)	4(99.2)	2(70.7)	1(63.0)	-	3(87.6)
<i>Pleidiu</i> sp.	1(-)	7(-)	2(-)	1(-)	2(-)	2(-)	5(-)	3(-)
<i>Sphaerium</i> sp.	3(6.9)	2(8.5)	-	-	1(1.6)	2(1.1)	1(0.7)	-

NA = Not Available.

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 163

POOR ORIGINAL

Table 3.2-5

Numbers and milligrams of biomass, μ (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQ1-LA1, August 1976. Dashes indicate species not present or no weight measurement made.

Date	3 August				17 August			
Time	0838				0820			
Air Temp. (C)	21.7				19.0			
Water Temp. (C)	22.0				19.0			
Dissolved Oxygen (ppm)	7.7				8.3			
pH	8.4				8.0			
Secchi Disc (cm)	50.3				15.2			
River Stage (ft)	4.2				4.5			
Substrate	Coal, Sand, Mud, Detritus				Coal, Sand, Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
Nematoda	-	1(-)	1(-)	1(-)	-	-	1(-)	-
<i>Branchiura sewerhi</i>	1(-)	-	-	2(9.2)	-	1(0.2)	-	1(1.2)
<i>Ilvodrillus campietoni</i>	-	-	-	-	-	-	1(-)	-
<i>Limnodrilus clapparedianus</i>	11(2.9)	11(2.3)	-	20(4.6)	-	10(3.0)	21(5.7)	31(8.2)*
<i>L. hoffmeisteri</i>	45(11.8)	11(2.3)	20(2.5)	61(13.9)	33(10.5)	141(42.2)	42(16.8)	92(24.4)
<i>L. udekemianus</i>	-	21(4.3)	-	10(2.3)	22(7.0)	-	-	-
<i>Limnodrilus</i> spp.	101(26.5)	87(17.9)	70(19.6)	122(27.8)	32(10.2)	60(18.0)	41(11.1)	134(35.6)
<i>Pelosciolex multisetosus</i>	-	-	-	-	-	1(0.3)	-	-
imm. tub./cap. chaetae*	-	1(-)	-	-	-	-	1(-)	3(-)
<i>Sparganophilus eiseni</i>	1(-)	-	-	-	-	-	-	-
<i>Crangonyx</i> sp.	1(-)	-	-	-	-	-	-	-
<i>Gammarus fasciatus</i>	1(-)	3(0.4)	-	2(0.4)	2(5.0)	-	-	1(0.4)
<i>Hexagenia</i> sp.	-	-	-	-	-	-	-	1(-)
<i>Caenis</i> sp.	-	-	-	-	-	-	-	1(-)
<i>Procladius</i> sp.	-	4(0.2)	1(-)	2(-)	9(1.6)	2(-)	2(-)	5(1.0)
<i>Chironomus decorus</i>	1(-)	-	-	1(-)	-	1(-)	-	3(-)
<i>Cryptochironomus</i> nr. <i>blarina</i>	-	-	-	9(0.8)	-	-	-	-
<i>C. nr. fulvus</i>	19(7.3)	32(4.0)	17(1.3)	19(1.7)	3(-)	1(-)	5(-)	10(2.1)
<i>Cryptochironomus</i> spp. pupa	-	-	-	2(-)	-	-	-	-
<i>Paratadonella</i> sp.	2(-)	-	-	19(1.2)	-	-	1(-)	1(-)
<i>Phaenopsectra</i> nr. <i>chediens</i>	-	-	1(-)	-	-	-	-	-
<i>Polypedilum scalaenum</i>	35(1.0)	48(1.9)	32(2.7)	28(1.4)	-	-	-	10(0.8)
<i>P. scalaenum</i> pupa	-	1(-)	-	1(-)	-	-	-	-
<i>Physa</i> sp.	-	-	-	3(-)	-	-	-	-
<i>Goniobasis virginica</i>	2(76.3)	1(31.4)	2(62.6)	2(123.7)	1(69.0)	2(126.2)	6(268.5)	8(421.2)
Calonidae immature	1(-)	-	-	-	-	-	-	-
<i>Pisidium</i> sp.	4(-)	7(-)	2(-)	9(3.0)	3(-)	3(-)	1(-)	2(-)
<i>Sphaerium</i> sp.	2(-)	8(15.7)	-	6(5.2)	-	-	-	-

* Immature tubificid with capilliform chaetae.

1565 164

Table 3.2-c

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TH-A01-1A1, September 1976. Dashes indicate species not present or no weight measurement made.

Date	7 September				21 September			
Time	0817				0830			
Air Temp. (C)	14.5				18.0			
Water Temp. (C)	19.0				20.0			
Dissolved Oxygen (ppm)	7.8				7.3			
pH	7.9				8.0			
Secchi Disc (cm)	50.8				50.8			
River Stage (ft)	3.5				4.1			
Substrate	Sand, Coal, Detritus				Sand, Coal, Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
<i>Turbellaria</i>	-	-	1(0.1)	-	1(-)	-	-	-
<i>Nemertinea</i>	-	1(-)	3(0.1)	3(0.6)	3(1.2)	3(-)	2(0.7)	4(-)
<i>Nematoda</i>	1(-)	2(-)	-	-	-	-	-	-
<i>Arctonasis lumbi</i>	-	-	-	-	-	-	1(-)	-
<i>Branchiura sowerbyi</i>	-	2(0.2)	2(1.7)	1(1.6)	1(0.7)	-	4(9.7)	1(-)
<i>Ilyodrilus templetoni</i>	-	-	-	-	2(-)	-	-	1(-)
<i>Limnodrilus clavigerianus</i>	-	-	11(2.7)	-	21(3.7)	-	10(2.6)	-
<i>L. hoffmeisteri</i>	30(9.5)	21(6.1)	-	42(15.7)	21(3.7)	40(7.6)	20(5.1)	9(2.1)
<i>Limnodrilus</i> spp.	39(12.4)	55(24.6)	44(10.7)	-	154(27.0)	49(9.3)	51(13.1)	57(13.2)
<i>Pelosclex multisetosus</i>	1(-)	-	-	-	-	-	1(-)	-
<i>Tubifex tubifex</i>	-	-	-	1(-)	-	-	-	-
imm. tub./cap. chaetae*	-	1(-)	4(-)	-	1(-)	1(-)	10(2.6)	-
<i>Gammarus fasciatus</i>	-	-	4(0.8)	-	1(0.5)	5(3.5)	10(1.8)	1(-)
<i>Hexagenia limbata</i>	-	-	1(0.9)	-	-	-	-	-
<i>Hexagenia</i> spp.	-	-	-	-	1(-)	-	-	-
<i>Gomphus spinicrus</i>	-	-	-	-	-	1(-)	-	-
<i>Macronia illinoensis</i>	-	-	1(2.3)	-	-	-	-	-
<i>Oecetis</i> sp.	-	-	1(-)	-	-	-	-	-
<i>Dubiraphia</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Procladius</i> sp.	-	-	-	-	1(-)	-	-	2(-)
<i>Ablabesmus</i> sp.	-	-	1(-)	-	1(-)	-	1(-)	-
<i>Cricotopus</i> spp.	-	-	-	-	-	-	1(-)	-
<i>Chironomus decorus</i>	-	-	1(-)	-	142(38.9)	66(19.5)	45(16.6)	26(7.7)
<i>C. decorus</i> pupa	-	-	-	-	2(-)	1(-)	3(-)	-
<i>Cryptochironomus</i> nr. <i>fulvus</i>	2(-)	-	1(-)	1(-)	-	1(-)	1(-)	-
<i>Cryptochironomus</i> spp. pupa	-	-	-	-	-	1(-)	-	-
<i>Demicyrtichironomus vulneratus</i>	-	-	-	-	-	-	-	1(-)
<i>Paracladocera</i> sp.	-	-	-	-	-	2(-)	1(-)	3(-)
<i>Polypedilum</i> nr. <i>illinoense</i>	-	-	-	-	-	-	1(-)	-
<i>P. scalaeum</i>	8(1.0)	-	1(-)	4(-)	-	-	-	-
<i>Ferrissia</i> spp.	-	-	-	-	2(-)	3(-)	-	-
<i>Goniobasis virginica</i>	3(134.2)	1(30.9)	5(287.7)	-	5(214.2)	9(459.4)	3(164.3)	2(83.3)
<i>Pisidium</i> sp.	29(-)	15(-)	27(-)	7(-)	16(-)	16(-)	21(-)	10(-)
<i>Sphaerium</i> sp.	8(7.6)	2(2.6)	-	-	-	-	5(-)	7(-)

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

Table 3.2-7

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TN-A12-141, October 1976. Dashes indicate species not present or no weight measurement made.

Date	5 October				19 October			
Time	0832				0830			
Air Temp. (C)	14.0				2.5			
Water Temp. (C)	15.0				9.0			
Dissolved Oxygen (ppm)	9.2				11.6			
pH	8.0				7.5			
Secchi Disc (cm)	30.5				27.9			
River Stage (ft)	4.8				5.3			
Substrate	Mud, Coal, Sand, Detritus				Sand, Coal, Mud			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	-	1(-)	1(-)	-	-
Nemertinea	-	-	1(-)	-	1(-)	3(-)	6(0.9)	1(-)
Nematoda	1(-)	5(0.4)	3(-)	4(-)	2(-)	1(-)	1(-)	1(-)
<i>Manayunkia speciosa</i>	-	-	-	-	-	-	-	1(-)
<i>Branchiura sowerbyi</i>	-	-	-	-	1(-)	-	-	-
<i>Limnodrilus claparotianus</i>	-	12(6.1)	-	-	-	-	10(2.5)	-
<i>L. hoffmeisteri</i>	20(7.9)	23(11.6)	21(2.8)	-	-	10(1.7)	21(5.3)	22(6.3)
<i>L. udekemianae</i>	-	12(6.1)	-	10(3.2)	10(1.6)	-	-	22(6.3)
<i>Limnodrilus</i> spp.	88(34.9)	92(36.6)	196(25.7)	88(28.0)	143(23.1)	164(28.3)	105(26.2)	21(6.1)
Imm. tub./cap. chaetae*	-	1(-)	-	-	1(-)	-	1(-)	-
<i>Gammarus fasciatus</i>	-	1(-)	1(0.4)	1(-)	-	1(-)	2(-)	-
<i>Hexagenia limbata</i>	1(3.7)	1(10.3)	-	1(3.7)	-	-	-	-
<i>Procladius</i> sp.	-	-	-	-	-	1(-)	-	-
<i>Chironomus</i> sp.	-	-	-	3(0.2)	-	-	-	-
<i>Hydropsyche bifida</i> group sp.	-	-	-	1(-)	-	-	-	-
<i>H. phalerata</i>	-	-	-	1(-)	-	-	-	-
<i>Dubiraphia</i> sp.	-	1(-)	-	-	-	-	-	-
<i>Procladius</i> sp.	-	-	-	2(-)	-	-	-	-
<i>Eukiefferiella</i> sp. A	-	1(-)	-	-	-	-	-	-
<i>Cricotopus</i> spp.	4(-)	2(-)	-	2(-)	-	-	-	-
<i>Orthocladus</i> (<i>Orthocladus</i>) sp.	1(-)	-	-	2(-)	-	-	-	-
<i>Psectrocladius</i> sp.	1(-)	-	-	1(-)	-	-	-	-
<i>Chironomus tentans</i>	32(12.5)	95(31.4)	106(32.5)	88(24.9)	-	1(-)	-	1(-)
<i>Cryptochironomus</i> or <i>blarina</i>	-	-	-	-	-	-	1(-)	-
<i>Picrotendipes modestus</i>	1(-)	2(-)	1(-)	7(-)	-	-	-	-
<i>Paratendipes</i> sp.	-	-	1(-)	2(-)	-	-	-	-
<i>Rhyacotendipes</i> sp.	1(-)	2(-)	-	-	-	-	-	-
<i>Tanytarsus</i> sp.	-	1(-)	-	1(-)	-	-	-	-
<i>Physa</i> sp.	-	-	-	-	6(-)	12(-)	15(-)	4(-)
<i>Lymnaea</i> sp.	-	-	-	-	3(-)	4(-)	2(-)	-
<i>Cyranus carnosus</i>	-	-	-	-	-	7(-)	8(-)	4(-)
<i>Ferrissia</i> spp.	1(-)	-	-	-	65(14.8)	267(-)	134(-)	135(-)
<i>Goniodontia virginica</i>	2(73.1)	1(208.8)	2(74.3)	2(123.8)	-	3(70.0)	1(66.7)	-
<i>Spirodon carinata</i>	-	-	-	-	-	1(-)	-	-
<i>Pisidium</i> sp.	8(-)	9(-)	16(-)	5(-)	28(-)	32(-)	39(-)	13(-)
<i>Sphaerium</i> sp.	2(-)	4(-)	3(-)	1(-)	52(-)	80(59.2)	68(35.3)	46(-)

* Immature tubificid with capilliform chaetae.

1565 166

Table 3.2-8

Numbers and milligrams of biomass, μ (mg), of benthic organisms per Ponar grab (529 cm²) at Station TD-AQ1-1A2, April 1976. Dashes indicate species not present or no weight measurement made.

Date	6 April				20 April			
Time	0950				0950			
Air Temp. (C)	16.5				24.0			
Water Temp. (C)	8.0				21.0			
Dissolved Oxygen (ppm)	9.7				8.1			
pH	8.4				7.9			
Secchi Disc (cm)	38.1				114.3			
River Stage (ft)	6.6				4.7			
Substrate	Sand, Coal, Mud, Detritus				Mud, Sand, Coal, Detritus			
Replicate	A	B	C	D	A	B	C	D
Amatoda	-	1(-)	1(-)	-	2(-)	1(-)	3(-)	1(-)
<i>Nais breesebergi</i>	-	-	-	-	-	1(-)	1(-)	-
<i>N. elingensis</i>	-	-	5(-)	-	6(-)	-	1(-)	2(-)
<i>N. variabilis</i>	-	-	-	-	-	1(-)	3(-)	-
<i>Branchinella sowerbyi</i>	-	-	-	-	-	-	1(-)	-
<i>Limnodrilus clapparedianus</i>	-	-	-	-	-	3(1.2)	-	-
<i>L. hoffmeisteri</i>	-	22(7.3)	3(-)	-	21(5.3)	3(1.2)	-	-
<i>L. melleokianus</i>	-	32(10.6)	1(-)	-	-	-	0(2.4)	-
<i>Limnodrilus</i> spp.	29(9.0)	-	11(-)	28(7.5)	-	12(3.9)	2(0.6)	11(5.2)
<i>Pelosclex multisetosus</i>	-	-	-	-	1(-)	1(-)	-	-
<i>Tubificoides tubificoides</i>	-	-	-	-	-	2(-)	-	-
Imm. tub. cap. chaetae*	9(2.8)	-	3(-)	-	-	1(-)	10(2.6)	13(6.2)
Megadrile	1(-)	-	-	-	-	-	-	-
<i>Gammarus fasciatus</i>	-	-	-	-	8(15.0)	-	-	-
<i>Cryptochironomus</i> sp. fulvus	-	-	-	1(-)	-	-	-	-
<i>Hyalella</i> sp. ar. obediens	-	-	-	-	-	-	1(-)	-
<i>Corbicula virginica</i>	7(137.2)	3(55.9)	20(549.7)	5(100.0)	2(38.0)	5(233.7)	7(180.7)	3(115.0)
<i>Pisidium</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Sphaerium</i> sp.	-	-	-	1(-)	-	-	-	-

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 167

POOR ORIGINAL

Table 3.2-9

Numbers and milligrams of biomass, = (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-1A2, May 1976. Dashes indicate species not present or no weight measurement made.

Date	4 May				18 May			
Time	0959				1028			
Air Temp. (C)	NA				19.0			
Water Temp. (C)	11.8				19.8			
Dissolved Oxygen (ppm)	8.9				7.0			
pH	7.8				NA			
Secchi Disc (cm)	71.1				45.7			
River Stage (ft)	4.9				4.6			
Substrate	Mud, Sand, Coal, Detritus				Mud, Sand, Some Coal, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	1(-)	-	-	-	-	-
Nematoda	1(-)	-	2(-)	3(-)	2(-)	-	-	1(-)
<i>Aulophorus furcatus</i>	-	-	-	-	1(-)	-	-	-
<i>Chaetogaster diaphanus</i>	-	-	-	-	75(-)	63(1.7)	88(-)	-
<i>Nais bretscheri</i>	-	-	-	-	8(0.2)	-	-	-
<i>N. elliptica</i>	36(0.5)	27(-)	3(-)	39(0.6)	7(0.2)	6(-)	-	64(1.2)
<i>N. variabilis</i>	9(0.1)	7(-)	-	13(0.2)	-	-	19(-)	48(0.9)
<i>Nais</i> sp.	-	3(-)	1(-)	-	-	-	-	-
<i>Ophidonais serpentina</i>	-	-	-	-	11(0.2)	2(-)	-	-
<i>Paranais frici</i>	-	-	-	-	3(0.1)	-	-	-
<i>Slavina appendiculata</i>	-	-	-	-	3(0.1)	1(-)	6(-)	-
<i>Aulodrilus plurisetus</i>	-	-	-	-	-	1(-)	-	-
<i>Branchiura sowerbyi</i>	-	1(-)	-	-	-	-	-	-
<i>Ilvodrillus templetoni</i>	-	-	-	-	-	-	1(-)	-
<i>Limnodrilus hoffmeisteri</i>	-	53(15.8)	1(-)	10(3.0)	11(1.1)	54(25.1)	2(-)	15(1.7)
<i>Limnodrilus</i> spp.	39(8.6)	43(12.8)	4(-)	19(5.8)	99(10.0)	-	8(-)	262(30.3)
<i>Peloscolex multisetosus</i>	-	-	-	-	6(-)	-	1(-)	2(-)
<i>Tubifex tubifex</i>	1(-)	-	-	-	2(-)	10(-)	-	2(-)
Imm. tub./cap. chaetae*	1(-)	-	-	-	2(-)	1(-)	2(-)	1(-)
Megadrile	-	-	-	1(-)	-	-	-	-
<i>Gammarus fasciatus</i>	-	-	-	-	1(-)	-	2(-)	-
<i>Cheumatopsyche</i> sp.	-	-	-	1(-)	-	-	-	-
<i>Stenelmis</i> sp.	-	-	-	-	-	-	-	1(-)
<i>Procladius</i> sp.	-	-	-	-	14(7.3)	8(1.7)	6(-)	18(5.1)
<i>Procladius</i> sp. pupa	-	-	-	-	-	-	-	1(-)
<i>Psectrotanytus</i> sp.	-	-	-	-	-	2(0.4)	-	-
<i>Cricotopus</i> spp.	-	-	-	-	1(-)	2(0.4)	-	-
<i>Cricotopus</i> spp. pupa	-	-	-	-	-	-	1(-)	-
<i>Orthocladus</i> (<i>Orthocladus</i>) sp.	-	-	-	5(-)	-	-	-	-
<i>Psectrocladius</i> sp.	4(0.4)	-	-	2(-)	-	-	-	-
<i>Chironomus decorus</i>	7(0.7)	1(-)	-	9(0.9)	45(19.8)	16(5.4)	17(5.5)	38(42.4)
<i>C. decorus</i> pupa	-	-	-	1(-)	1(1.5)	-	-	3(4.9)
<i>Phaenopsectra</i> nr. <i>obediens</i>	-	-	-	-	-	2(0.4)	-	-
<i>Physa</i> sp.	-	-	-	-	-	-	1(-)	-
<i>Goniobasis virginica</i>	2(3.9)	2(46.6)	5(206.8)	4(117.6)	2(134.6)	1(11.7)	7(256.1)	-
<i>Pisidium</i> sp.	-	1(-)	1(-)	-	-	-	-	-
<i>Sphaerium</i> sp.	1(-)	-	-	-	-	-	-	-

NA = Not Available.

* Immature tubificid with capilliform chaetae.

1565 168

POOR ORIGINAL

Table 3.2-10

Numbers and milligrams of biomass, = (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQ1-1A2, June 1976. Dashes indicate species not present or no weight measurement made.

Date	1 June				15 June			
Time	0837				0805			
Air Temp. (C)	21.5				24.0			
Water Temp. (C)	19.0				23.0			
Dissolved Oxygen (ppm)	9.2				7.2			
pH	8.3				8.0			
Secchi Disc (cm)	76.2				45.7			
River Stage (ft)	5.1				4.1			
Substrate	Coal, Sand, Mud, Detritus				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
Nematoda	8(-)	9(-)	1(-)	3(-)	6(-)	3(-)	1(-)	9(0.4)
<i>Acarteanaia lomondi</i>	-	-	-	-	-	1(-)	-	-
<i>Nais elinguis</i>	2(-)	-	-	2(-)	-	-	-	-
<i>N. variabilis</i>	2(-)	-	-	3(-)	-	-	-	-
<i>Alvodrillus templetoni</i>	1(-)	-	-	-	-	-	-	-
<i>Limnodrilus clonaredeiianus</i>	-	31(6.0)	-	-	30(7.5)	-	-	10(1.4)
<i>L. hoffmeisteri</i>	11(2.3)	153(29.4)	46(5.6)	66(16.6)	30(7.5)	39(16.5)	80(18.1)	30(4.2)
<i>Limnodrilus</i> spp.	55(11.6)	185(35.6)	57(6.9)	98(24.7)	61(15.3)	39(16.5)	159(35.9)	49(6.9)
<i>Pelosciolex multisetosus</i>	1(-)	1(-)	-	2(-)	2(-)	5(-)	2(0.2)	4(-)
<i>Tubifex tubifex</i>	3(-)	5(-)	3(-)	4(-)	-	2(0.2)	-	2(-)
imm. tub./cap. chaetae*	4(-)	2(-)	-	4(-)	2(-)	17(1.7)	2(-)	5(-)
<i>Gammarus fasciatus</i>	-	-	-	-	-	-	1(-)	-
<i>Hydropsyche</i> sp.	1(-)	-	-	-	-	-	-	-
<i>Chironomid</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Stenelmis</i> spp.	2(1.3)	-	-	1(1.0)	-	-	-	1(0.4)
<i>Procladius</i> sp.	10(2.7)	3(-)	4(-)	3(-)	6(2.2)	31(5.1)	7(1.6)	5(1.7)
<i>Psectrocanvpus</i> sp.	-	-	1(-)	-	-	-	-	-
<i>Chironomus decorus</i>	9(7.8)	14(25.2)	2(-)	18(6.1)	374(144.2)	251(85.4)	285(119.7)	220(98.6)
<i>C. decorus</i> pupa	-	2(-)	-	1(-)	2(1.3)	1(1.4)	-	-
<i>Chaenopsectra</i> nr. <i>obediens</i>	1(-)	-	-	-	6(1.1)	10(0.7)	1(-)	-
<i>Polypodilum</i> nr. <i>illinoense</i>	-	-	-	-	-	2(0.1)	-	-
<i>Palpomyia</i> group sp. pupa	-	-	-	-	-	1(-)	-	-
<i>Conibasis virginica</i>	-	2(67.5)	2(44.4)	3(76.6)	1(65.2)	3(88.7)	6(243.2)	1(38.1)
<i>Pisidium</i> sp.	1(-)	-	-	-	-	-	-	-

* Immature tubificid with capilliform chaetae.

1565 169

Table 3.2-11

Numbers and milligrams of biomass, \pm (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQ1-LA2, July 1976. Dashes indicate species not present or no weight measurement made.

Date	6 July				20 July			
Time	0825				0817			
Air Temp. (C)	23.0				22.0			
Water Temp. (C)	24.0				24.0			
Dissolved Oxygen (ppm)	NA				7.3			
pH	8.0				8.2			
Secchi Disc (cm)	58.4				30.5			
River Stage (ft)	4.7				4.3			
Substrate	Mud, Sand, Coal, Detritus				Mud, Detritus, Sand			
Replicate	A	B	C	D	A	B	C	D
<i>Nematoda</i>	10(0.7)	3(-)	2(-)	-	-	-	3(0.1)	3(-)
<i>Branchiura sowerbyi</i>	-	-	-	1(-)	-	-	-	-
<i>Hydrilus templetoni</i>	1(-)	-	-	-	-	1(-)	-	-
<i>Limnodrilus claperedenianus</i>	-	10(2.0)	9(1.3)	-	-	-	-	-
<i>L. hoffmeisteri</i>	21(3.8)	10(2.0)	27(4.0)	55(7.0)	32(5.5)	30(6.7)	57(5.7)	67(7.9)
<i>Limnodrilus</i> spp.	82(14.7)	174(35.0)	153(19.7)	86(11.2)	138(23.7)	179(40.1)	288(28.8)	270(31.8)
<i>Pelosclex multiserpens</i>	3(-)	-	-	1(-)	-	-	-	1(-)
<i>Tubifex tubifex</i>	-	2(0.2)	-	-	-	-	-	-
Imm. tub./cap. chaetae*	6(-)	10(0.8)	5(-)	1(-)	1(-)	7(-)	3(-)	4(-)
<i>Helobdella elongata</i>	-	-	-	-	-	1(0.8)	-	-
<i>Optiservus</i> sp.	-	-	1(0.6)	-	-	-	-	-
<i>Procladius</i> sp.	7(-)	4(-)	1(-)	1(-)	17(6.8)	19(1.9)	13(2.3)	5(-)
<i>Abiasesmyia</i> sp.	-	-	-	-	4(2.0)	1(-)	-	1(-)
<i>Chironomus decorus</i>	115(8.2)	136(10.3)	138(11.5)	114(13.2)	20(13.4)	17(14.1)	42(20.8)	24(13.9)
<i>C. decorus</i> pupa	-	1(1.1)	-	-	1(-)	-	1(-)	-
<i>Paracladopelma</i> sp.	-	-	2(-)	-	-	-	-	-
<i>Polypedilum halterale</i>	-	-	-	-	2(1.0)	-	-	-
<i>P. m. illinoense</i>	-	-	-	-	-	2(-)	-	-
<i>Palpomyia</i> group sp.	-	-	-	-	1(-)	-	1(-)	-
<i>Goniodontia virginica</i>	-	2(101.6)	1(46.9)	-	-	-	-	-
<i>Pisidium</i> sp.	-	-	-	-	-	-	-	-
<i>Sphaerium</i> sp.	-	2(1.7)	-	-	-	2(-)	-	1(-)

NA = Not Available.

* Immature tubificid with capilliform chaetae.

1565 170

Table 3.2-12

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (519 cm²) at station TM-AQI-1A2, August 1976. Dashes indicate species not present or no weight measurement made.

Date	3 August				17 August			
Time	0848				0842			
Air Temp. (C)	21.0				20.0			
Water Temp. (C)	21.0				18.0			
Dissolved Oxygen (ppm)	7.7				7.9			
pH	8.0				8.0			
Secchi Disc (cm)	45.7				15.2			
River Stage (ft)	4.1				4.5			
Substrate	Coal, Sand, Mud, Detritus				Mud, Detritus, Some Sand, Coal			
Replicate	A	B	C	D	A	B	C	D
Nematoda	-	-	2(-)	3(-)	2(-)	1(-)	4(0.6)	4(-)
<i>Limnodrilus cervix</i>	10(2.0)	-	-	-	-	-	-	-
<i>L. hoffmeisteri</i>	31(6.1)	31(5.4)	42(7.4)	46(9.2)	32(6.3)	115(20.6)	40(6.4)	57(11.2)
<i>Limnodrilus</i> spp.	52(10.2)	216(37.8)	218(38.2)	104(20.9)	84(16.5)	94(16.8)	100(16.1)	182(35.9)
<i>Peloscoides multisetus</i>	-	2(0.4)	-	-	-	-	-	-
<i>Tubifex tubifex</i>	-	-	-	-	1(-)	-	-	-
imm. tub./cap. chaetae*	3(-)	3(-)	3(-)	3(-)	2(-)	-	12(0.9)	9(1.1)
<i>Helobdella</i> spp.	-	-	-	-	-	1(-)	-	-
<i>Orconectes</i> spp.	-	-	-	-	-	-	-	1(-)
<i>Promelasia</i> sp.	-	-	-	1(-)	-	-	-	-
<i>Stenelmis</i> spp.	-	-	-	-	-	-	-	-
<i>Procladius</i> sp.	6(0.5)	25(2.9)	11(2.8)	12(1.6)	20(3.1)	1(0.4)	-	-
<i>Ablabesmyia</i> sp.	-	-	-	1(-)	-	30(6.0)	28(3.4)	26(4.8)
<i>Chironomus decorus</i>	69(20.3)	109(32.8)	159(29.5)	138(30.5)	47(19.5)	53(36.3)	79(38.7)	94(48.3)
<i>C. decorus</i> pupa	-	1(-)	3(-)	2(-)	2(-)	1(-)	3(-)	-
<i>Cryptochironomus</i> nr. <i>fulvus</i>	2(0.2)	2(-)	-	1(-)	-	-	-	-
<i>Paracladopelma</i> sp.	-	1(-)	3(0.4)	-	-	-	-	-
<i>Goniobasis virginica</i>	2(143.1)	3(198.3)	1(30.1)	-	-	3(3.6)	5(238.3)	3(118.6)
<i>Pisidium</i> sp.	-	-	-	-	-	-	2(-)	-

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 171

Table 3 - 13

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-1A2, September 1976. Dashes indicate species not present or no weight measurement made.

Date	7 September				21 September			
Time	0830				0840			
Air Temp. (C)	14.5				18.0			
Water Temp. (C)	19.0				19.0			
Dissolved Oxygen (ppm)	9.0				8.0			
pH	8.0				8.0			
Secchi Disc (cm)	33.0				30.5			
River Stage (ft)	3.5				4.1			
Substrate	Mud, Detritus, Coal				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
Nemertinea	-	-	-	-	-	1(-)	-	-
Nematoda	-	1(-)	1(-)	1(-)	-	-	-	1(-)
<i>Limnodrilus claparedelanus</i>	-	-	-	-	13(2.4)	-	-	17(2.1)
<i>L. hoffmeisteri</i>	81(9.1)	72(10.2)	34(5.2)	73(9.9)	51(9.3)	21(3.2)	144(20.6)	17(2.1)
<i>L. udekemianus</i>	-	14(2.0)	-	-	-	-	-	17(2.1)
<i>Limnodrilus</i> spp.	322(36.0)	273(38.8)	253(38.5)	381(51.9)	254(46.5)	230(35.5)	529(75.7)	367(44.6)
<i>Pelosciolex multisetosus</i>	-	-	-	-	1(-)	-	-	1(-)
Imm. tub./cap. chaetae*	3(-)	-	-	-	-	4(-)	7(-)	4(-)
<i>Helobdella elongata</i>	-	-	-	1(0.5)	-	-	-	-
<i>Helobdella</i> sp.	-	-	-	1(-)	-	-	-	-
<i>Hexagramia limbata</i>	-	-	-	1(3.0)	-	-	1(19.4)	1(6.5)
<i>Procladius</i> sp.	12(1.9)	25(3.4)	15(2.7)	22(3.0)	5(-)	12(3.9)	9(2.5)	9(2.1)
<i>Chironomus decorus</i>	68(18.8)	69(11.0)	41(8.1)	37(7.7)	27(8.1)	20(7.1)	28(8.4)	28(9.0)
<i>C. decorus</i> pupa	-	-	-	-	-	1(-)	1(-)	1(-)
<i>Cryptochironomus</i> nr. <i>fulvus</i>	-	3(-)	1(-)	2(-)	-	-	-	-
<i>Paratubificoides</i> sp.	-	-	1(-)	-	-	-	-	-
<i>Gonichasis virginica</i>	3(96.0)	2(142.3)	1(92.4)	2(58.9)	3(120.2)	5(215.4)	3(218.4)	2(211.7)
<i>Pisidium</i> sp.	4(-)	-	1(-)	-	1(-)	-	-	-

* Immature tubificid with capilliform chaetae.

1565 172

Table 1.2-14

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at station TM-AQ1-1A2, October 1970. Dashes indicate species not present or no weight measurement made.

Date	5 October				10 October			
Time	0850				0845			
Air Temp. (C)	13.0				3.0			
Water Temp. (C)	15.5				7.1			
Dissolved Oxygen (ppm)	9.4				11.5			
pH	8.0				7.1			
Secchi Disc (cm)	27.9				30.5			
River Stage (ft)	4.8				5.3			
Substrate	Mud, Detritus, Some Sand				Sand, Coal, Litter, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	-	-	-	-	1(-)
Nemertinea	-	-	-	-	2(-)	-	-	-
<u>Planatella repens</u>	-	-	-	-	-	-	-	-
<u>Limnospilus templeroni</u>	-	1(-)	-	-	-	-	-	-
<u>Limnospilus cervix</u>	-	-	-	10(1.8)	-	-	-	-
<u>L. clausenianus</u>	-	-	10(2.8)	-	-	-	-	-
<u>L. buffingtoni</u>	17(2.9)	20(6.9)	39(11.1)	41(8.5)	10(1.6)	-	10(6.7)	18(2.1)
<u>L. udekemianus</u>	-	-	-	-	-	-	10(3.1)	37(7.8)
<u>Limnospilus</u> spp.	390(67.1)	109(55.8)	173(50.6)	174(16.3)	59(9.5)	59(10.7)	179(60.2)	364(80.7)
<u>Pelocolex multisetosus</u>	-	-	-	1(-)	-	-	-	-
Imm. tub./cap. chaetae**	-	2(-)	2(-)	2(-)	-	-	-	-
<u>Gammarus fasciatus</u>	-	-	-	-	3(0.1)	16(5.4)	2(-)	26(4.9)
Collembola	-	-	-	-	-	-	-	1(-)
<u>Hexagenia limbata</u>	-	-	-	-	1(1.7)	-	-	2(26.9)
<u>Gomphus spiniceps</u>	-	-	-	-	-	-	1(50.1)	-
Corixidae	-	-	-	-	-	-	-	1(-)
<u>Hydropsyche phalerata</u>	-	-	-	-	-	1(0.9)	-	2(0.7)
<u>Procladius</u> sp.	-	-	-	-	-	-	-	1(-)
<u>Stenelmis</u> spp.	-	-	1(0.5)	-	-	-	-	-
<u>Chaoborus punctipennis</u>	1(0.3)	-	-	-	-	-	-	-
<u>Procladius</u> sp.	3(-)	3(-)	2(-)	8(1.1)	-	-	-	-
<u>Chironomus decorus</u>	10(3.2)	13(3.6)	14(4.6)	11(2.4)	1(-)	-	1(-)	1(-)
<u>Cryptochironomus</u> nr. <u>fulvus</u>	-	-	-	-	-	1(-)	-	1(-)
<u>Microtendipes modestus</u>	-	1(-)	-	-	-	-	-	-
<u>Palaeomonis</u> group sp.	-	1(-)	-	-	-	-	-	-
<u>Physa</u> sp.	-	-	-	-	-	1(-)	1(-)	-
<u>Ferrissia</u> spp.	-	-	-	-	8(-)	193(-)	7(-)	1(-)
<u>Conchosis virginica</u>	4(198.2)	3(146.1)	-	2(84.5)	-	1(-)	-	-
<u>Pisidium</u> sp.	3(-)	-	-	1(-)	-	7(-)	-	-
<u>Sphaerium</u> sp.	-	-	-	1(-)	5(-)	13(-)	1(-)	1(-)

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 173

Table 3.2-15

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A1, April 1976. Dashes indicate species not present or no weight measurement made.

Date	6 April				20 April			
Time	1045				1004			
Air Temp. (C)	19.0				24.5			
Water Temp. (C)	8.0				20.5			
Dissolved Oxygen (ppm)	10.3				7.9			
pH	8.4				7.7			
Secchi Disc (cm)	30.5				61.0			
River Stage (ft)	6.6				4.7			
Substrate	Mud, Detritus, Some Coal				Mud, Detritus, Coal			
Replicate	A	B	C	D	A	B	C	D
<i>Turbellaria</i>	-	-	-	-	1(0.3)	-	-	1(0.8)
<i>Nemertinea</i>	-	-	2(-)	-	-	-	-	-
<i>Urostella gracilis</i>	-	-	-	-	-	-	-	-
<i>Nereis brevifrons</i>	-	-	2(-)	-	1(-)	-	-	-
<i>Limnodrilus clavigerulus</i>	-	1(-)	-	-	12(4.7)	1(-)	-	11(8.7)
<i>L. hoffmeisteri</i>	3(2.0)	8(-)	3(-)	1(-)	12(4.7)	5(-)	91(51.7)	-
<i>L. udebenianus</i>	-	2(-)	2(-)	-	-	1(-)	-	-
<i>Limnodrilus</i> spp.	20(13.3)	6(-)	3(-)	4(-)	15(9.8)	5(-)	-	22(17.4)
<i>Polyscolus multisetosus</i>	-	-	-	-	-	4(1.6)	2(0.8)	-
Imm. tub./cap. chaetae**	-	-	-	1(-)	-	2(-)	-	-
<i>Hebichia elongata</i>	-	-	-	-	-	-	1(0.8)	-
<i>H. lineata</i>	-	-	-	-	1(-)	-	-	-
<i>Ercobdellidae</i>	-	1(-)	-	-	-	-	-	3(20.4)
<i>Moerobdella microstoma</i>	-	-	-	-	2(21.1)	-	-	-
<i>Orthocladus</i> (Orthocladus) sp.	-	-	-	-	1(-)	-	-	-
<i>Cryptochironomus</i> nr. <i>fulvus</i>	-	-	-	-	-	-	1(-)	-
<i>Phaenopsectra</i> nr. <i>obediens</i>	-	-	-	-	-	1(-)	-	-
<i>Goniobasis virginica</i>	-	-	-	-	10(357.0)	4(15.8)	-	3(103.9)
<i>Amnicola limosa</i>	-	-	-	-	-	2(-)	-	-
<i>Pisidium</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Sphaerium</i> sp.	-	-	-	-	3(-)	2(-)	-	-

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 174

Table 3.2-16

Numbers and milligrams of biomass, \pm (mg), of benthic organisms per Ponar grab (529 cm²) at Station EM-AQ1-11A1, May 1976. Dashes indicate species not present or no weight measurement made.

Date	4 May				18 May			
Time	1024				1141			
Air Temp. (C)	NA				19.0			
Water Temp. (C)	11.0				14.0			
Dissolved Oxygen (ppm)	10.0				7.2			
pH	7.7				NA			
Secchi Disc (cm)	63.5				20.5			
River Stage (ft)	4.9				4.6			
Substrate	Mud, Detritus, Some Coal				Mud, Detritus, Some Sand			
Replicates	A	B	C	D	A	B	C	D
Nemertinea	-	2(-)	-	2(-)	-	-	-	-
Nematoda	5(0.2)	-	2(-)	3(-)	1(-)	1(-)	2(-)	2(-)
<u>Chaetogaster diaphanus</u>	-	-	-	-	-	-	-	1(-)
<u>Nais bretscheri</u>	1(-)	2(-)	3(-)	3(-)	-	2(-)	1(-)	3(-)
<u>N. elinguis</u>	2(-)	5(-)	2(-)	5(-)	3(-)	3(-)	6(-)	3(-)
<u>N. variabilis</u>	-	-	-	1(-)	-	-	-	-
<u>Paranais frici</u>	-	1(-)	-	-	-	-	-	-
<u>Stirlingia appendiculata</u>	-	-	-	-	-	-	-	1(-)
<u>Limnodrilus hoffmeisteri</u>	11(1.0)	4(0.3)	-	-	12(16.2)	16(1.2)	45(3.2)	-
<u>L. uckerianus</u>	-	-	-	6(0.8)	-	-	-	32(3.8)
<u>Limnodrilus</u> spp.	95(8.2)	17(1.4)	42(3.0)	24(3.2)	255(93.9)	372(11.2)	703(81.4)	915(109.0)
<u>Peloscolex multisetus</u>	2(0.2)	-	-	1(-)	-	-	-	-
imm. tub./cap. chaetae*	-	-	-	1(-)	-	-	2(-)	-
<u>Helobdella elongata</u>	1(0.8)	-	-	1(1.2)	-	-	-	-
Erpobdellidae	-	2(1.3)	-	1(0.4)	4(-)	4(7.8)	-	-
<u>Mooreobdella microstoma</u>	-	-	1(-)	-	-	-	-	-
<u>Gammarus fasciatus</u>	7(0.4)	9(1.0)	8(0.9)	27(2.4)	1(-)	7(2.1)	6(0.8)	2(-)
<u>Stenelmis</u> spp.	-	1(0.1)	-	-	-	-	-	-
<u>Psychoda</u> sp. pupa	1(-)	-	-	-	-	-	-	-
<u>Telmatoctopus</u> sp. pupa	-	-	-	1(0.2)	-	-	-	-
<u>Procladius</u> sp.	-	-	-	-	3(-)	-	-	2(-)
<u>Cricotopus</u> spp.	1(-)	-	-	1(-)	-	-	-	1(-)
<u>Orthocladius</u> (<u>Orthocladius</u>) sp.	4(-)	-	-	-	-	-	-	-
<u>Chironomus decorus</u>	2(-)	2(-)	3(-)	4(-)	15(7.5)	20(8.3)	6(-)	14(8.0)
<u>C. decorus</u> pupa	-	-	-	-	-	1(-)	-	-
<u>Cryptochironomus</u> nr. <u>folius</u>	-	-	1(-)	-	-	-	-	-
<u>Phaenocarpa</u> nr. <u>ghediensis</u>	-	-	-	-	-	-	-	2(-)
<u>Polypedium</u> nr. <u>illinoense</u>	1(-)	-	-	-	-	-	-	2(-)
Empididae	-	-	1(-)	-	-	-	-	-
<u>Coninbasis virginica</u>	1(7.7)	-	2(123.2)	3(104.9)	-	-	2(102.4)	1(36.9)
<u>Pisidium</u> sp.	-	1(-)	2(-)	-	-	-	-	-
<u>Sphaerium</u> sp.	-	3(0.3)	2(2.5)	3(6.3)	-	-	-	-

NA = Not Available

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 175

Table 3.2-17

Numbers and milligrams of biomass, μ (mg), of benthic organisms per Ponar grab 529 cm² at Station TH-AQI-11A1, June 1976. dashes indicate species not present or no weight measurement made.

Date	1 June				15 June			
Time	0850				0820			
Air Temp. °C	22.5				23.0			
Water Temp. °C	19.5				23.0			
Dissolved Oxygen (ppm)	7.8				7.7			
pH	8.0				8.0			
Secchi Disc cm	63.5				35.6			
River Stage ft.	5.1				4.1			
Substrate	Mud, Detritus, Some Gravel				Mud, Detritus, Some Sand, Some Gravel			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	1(0.5)	-	1(-)	-	1(-)
Nemertinea	1(-)	1(0.5)	-	-	-	1(-)	-	-
Plumatella perversa	-	P	P	P	-	-	-	-
Nematoda	6(-)	6(-)	11(-)	7(-)	9(1.1)	8(-)	15(1.1)	23(2.5)
Arctonemoides longicollis	-	-	-	1(-)	-	-	-	-
Maia brevis	3(-)	7(-)	3(-)	1(-)	-	-	1(-)	1(-)
M. elonensis	5(-)	1(-)	-	1(-)	-	-	1(-)	-
M. variabilis	-	-	-	1(-)	-	-	-	-
Paranais trilineatus	-	1(-)	-	-	-	-	-	-
Stauronema appendiculata	1(-)	-	-	1(-)	-	-	-	-
Limnodrilus hoffmeisteri	-	13(3.1)	-	-	-	-	-	-
L. hoffmeisteri	112(17.5)	81(19.3)	68(10.6)	147(25.3)	112(16.0)	116(15.0)	185(30.3)	113(27.5)
Limnodrilus sp.	24(84.3)	229(54.3)	359(66.4)	159(27.3)	187(40.9)	559(72.3)	278(57.6)	321(88.9)
Pelocoryx malincheus	2(-)	-	3(0.7)	5(0.9)	-	-	1(-)	-
Tubificoides tubificus	2(-)	-	-	-	-	-	-	-
Imm. tub./cap. chaetae**	-	13(3.1)	2(-)	-	-	-	-	-
Helobdella elongata	1(0.9)	1(2.8)	1(1.1)	-	1(0.1)	15(6.0)	5(2.4)	8(0.9)
Helobdella sp.	-	-	-	2(3.6)	-	-	-	-
Erpobdellidae	1(0.6)	3(0.9)	4(18.8)	1(-)	2(1.1)	7(7.0)	6(25.7)	6(3.2)
Monocorbellia nigrostrata	-	-	-	2(24.2)	-	-	-	-
Gammarus fasciatus	9(9.7)	5(0.6)	8(4.1)	33(18.5)	17(5.7)	90(26.4)	61(14.2)	72(15.9)
Hexapenia lirata	-	-	-	-	-	1(-)	-	-
Ceraclea tarsipunctata	-	-	-	-	-	-	-	-
Gomphus spini sp.	-	-	1(123.0)	-	-	-	1(-)	-
Stenelmis sp.	1(0.5)	1(1.4)	-	1(1.4)	1(152.0)	-	-	-
Procladius sp.	-	-	1(-)	-	2(0.4)	3(3.3)	2(1.7)	1(0.8)
Abalatesmyia nr. philospharax	-	-	-	-	-	2(0.9)	5(-)	6(-)
Orthocladus setipalpis, sp.	-	1(-)	-	-	-	-	1(-)	-
Isotrochodonta sp.	-	-	-	1(-)	-	-	-	-
Chironomus decorus	2(-)	3(-)	3(-)	3(-)	10(2.3)	11(2.8)	35(6.0)	42(7.1)
C. decorus pupa	-	-	3(2.8)	-	-	-	-	-
Cryptochironomus nr. fulvus	-	1(-)	-	-	6(0.9)	9(0.5)	7(-)	3(-)
C. nr. fulvus pupa	-	-	-	-	-	1(-)	-	-
Glyptotendipes sp.	-	-	1(-)	-	-	-	-	-
Harnischia sp. pupa	-	-	-	-	-	-	-	-
Paratendipes sp.	-	1(-)	-	-	-	1(-)	-	-
Phaenopsectra nr. phediens	3(-)	-	7(-)	1(-)	5(0.8)	16(0.8)	5(-)	12(-)
P. nr. phediens pupa	-	-	-	1(-)	-	-	-	-
Polypedilum haterale	-	-	1(-)	-	2(0.1)	3(0.1)	3(-)	3(-)
Polypedilum sp.	-	1(-)	-	-	-	-	-	-
Rheotanytarsus sp.	-	-	-	-	-	-	-	-
Tanytarsus sp.	1(-)	3(-)	1(-)	2(-)	4(0.2)	2(0.2)	8(-)	2(-)
Tanytarsus sp. pupa	-	-	-	-	-	-	-	-
Simulium sp.	-	1(-)	-	-	-	-	-	1(-)
Physa sp.	-	-	-	-	-	-	-	-
Goniobasis virgatula	-	3(62.8)	1(36.9)	2(46.7)	1(57.6)	8(293.6)	9(516.6)	9(425.4)
Pflidium sp.	1(-)	1(-)	2(-)	1(-)	-	-	3(-)	1(-)
Sphaerium sp.	1(-)	9(16.3)	1(4.4)	4(13.8)	8(21.9)	9(18.0)	2(-)	7(-)

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

1565 176

POOR ORIGINAL

Table 3.2-18

Numbers and milligrams of biomass, \pm (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A1, July 1976. Dashes indicate species not present or no weight measurement made.

Date	6 July				20 July			
Time	0840				0835			
Air Temp. (C)	23.5				23.0			
Water Temp. (C)	24.5				24.5			
Dissolved Oxygen (ppm)	NA				8.2			
pH	8.0				8.5			
Secchi Disc (cm)	25.4				15.2			
River Stage (ft)	4.7				4.3			
Substrate	Mud, Sand, Coal, Detritus				Mud, Sand, Coal, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	1(-)	-	1(0.3)	-	-	-	2(0.4)	3(4.1)
Nematoda	12(0.7)	11(0.5)	10(0.1)	9(1.0)	-	6(-)	5(-)	2(-)
<i>Urematella gracilis</i>	-	pe	-	-	-	P	-	-
<i>Limnodrilus hoffmeisteri</i>	99(15.3)	10(1.8)	58(11.1)	6(14.8)	29(7.6)	68(10.6)	40(4.6)	77(14.4)
<i>L. udakemianus</i>	12(1.8)	-	-	10(2.5)	-	-	-	-
<i>Limnodrilus</i> spp.	197(30.4)	234(42.9)	138(26.5)	120(29.5)	138(36.2)	216(33.8)	159(18.2)	308(57.5)
<i>Peloscoides multisetosus</i>	1(-)	-	1(-)	-	-	1(-)	-	-
<i>Helobdella elongata</i>	5(2.6)	7(1.9)	16(6.4)	5(2.6)	4(2.2)	7(2.1)	-	9(11.8)
Erpobdellidae	3(9.6)	4(17.8)	14(30.1)	-	5(24.2)	7(26.8)	1(0.6)	2(34.2)
<i>Gammarus fasciatus</i>	1(0.1)	3(0.3)	12(5.9)	-	-	-	-	-
<i>Oecetis</i> sp.	1(-)	-	-	-	-	-	-	-
<i>Procladius</i> sp.	-	-	-	-	-	-	-	-
<i>Chironomus texensis</i>	-	-	-	1(-)	-	1(-)	-	5(1.3)
<i>Cryptochironomus</i> nr. <i>blarina</i>	-	1(-)	-	-	-	5(-)	7(1.0)	2(0.2)
<i>C. nr. tubicis</i>	7(1.0)	8(1.6)	5(0.6)	6(-)	-	-	-	2(0.4)
<i>Paratubificoides</i> sp.	2(-)	-	4(0.5)	-	-	3(-)	9(1.2)	17(3.3)
<i>Phaenopsectra</i> nr. <i>rhodiens</i>	-	-	-	1(-)	-	-	-	4(0.5)
<i>Polypodium balticum</i>	-	-	-	-	-	-	-	-
<i>P. scalatum</i>	-	-	-	-	-	-	2(0.3)	-
<i>Gonichasis virginica</i>	3(31.2)	4(196.5)	1(50.4)	-	-	1(-)	4(0.5)	-
<i>Pisidium</i> sp.	14(-)	-(-)	2(-)	2(-)	4(183.0)	3(147.7)	4(207.7)	4(172.7)
<i>Sphaerium</i> sp.	6(3.2)	1(-)	1(-)	2(57.0)	5(2.0)	3(2.6)	1(-)	1(-)

NA = Not Available.

* Quantization not possible; denotes organism present.

POOR ORIGINAL

1565 177

Table 3.2-19

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A1, August 1976. Dashes indicate species not present or no weight measurement made.

Date	3 August				17 August			
Time	0905				0900			
Air Temp. (C)	21.0				19.0			
Water Temp. (C)	21.0				19.0			
Dissolved Oxygen (ppm)	8.7				8.9			
pH	8.0				8.0			
Secchi Disc (cm)	30.5				15.2			
River Stage (ft)	4.1				4.5			
Substrate	Mud, Detritus, Some Sand				Mud, Detritus, Some Gravel, Coal			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	2(0.3)	15(4.5)	12(1.5)	8(2.2)	5(1.2)
Nematoda	14(0.7)	9(0.6)	12(1.0)	-	11(1.3)	9(0.8)	8(0.6)	9(0.5)
<i>Urnatella gracilis</i>	-	P*	-	-	-	-	-	P
<i>Limnodrilus hoffmeisteri</i>	122(33.8)	99(18.2)	40(13.2)	69(17.2)	138(29.7)	129(27.7)	71(9.6)	308(40.7)
<i>Limnodrilus</i> spp.	92(25.5)	211(38.9)	119(39.2)	147(36.7)	175(37.7)	192(41.5)	172(23.3)	351(46.4)
<i>Peloscolex multisetosus</i>	1(-)	-	-	-	1(-)	-	1(-)	-
<i>Tubifex tubifex</i>	1(-)	-	-	-	-	-	-	-
imm. tub./cap. chaetae**	-	1(-)	1(-)	1(-)	-	1(-)	-	-
<i>Helobdella elongata</i>	2(0.2)	1(0.6)	1(1.0)	4(2.8)	5(2.5)	-	1(0.5)	6(2.8)
<i>Helobdella</i> spp.	-	-	-	-	-	1(-)	-	-
Erpobdellidae	2(1.8)	1(3.8)	2(10.9)	4(48.9)	1(0.5)	2(4.0)	1(5.7)	2(12.2)
<i>Gammarus fasciatus</i>	-	-	1(1.0)	10(3.3)	77(6.4)	24(1.0)	8(0.6)	11(0.9)
<i>Oecetis</i> sp.	-	-	-	-	1(-)	-	-	-
<i>Procladius</i> sp.	-	3(-)	-	-	8(1.6)	10(2.0)	5(2.0)	1(-)
<i>Chironomus decorus</i>	1(-)	3(-)	1(-)	2(-)	-	-	1(-)	1(-)
<i>Cryptochironomus</i> nr. <i>fulvus</i>	7(-)	2(-)	3(-)	-	2(0.4)	-	3(-)	3(-)
<i>Cryptochironomus</i> spp. pupa	-	2(-)	-	-	-	-	-	-
<i>Demicryptochironomus vulneratus</i>	-	-	-	1(-)	-	-	-	-
<i>Paracisodopelma</i> sp.	1(-)	2(-)	1(-)	2(-)	-	-	-	-
<i>Polypedium scalanum</i>	1(-)	-	-	-	-	-	-	-
<i>Physa</i> sp.	-	1(-)	-	-	-	-	-	-
<i>Goniobasis virginica</i>	2(53.9)	5(220.1)	3(130.2)	4(211.3)	5(281.5)	6(149.2)	12(455.5)	8(473.3)
<i>Carpeloma decisa</i>	-	-	-	-	1(-)	-	-	-
<i>Pisidium</i> sp.	6(-)	63(-)	5(2.4)	10(2.5)	36(9.8)	25(3.9)	16(-)	22(9.9)
<i>Sphaerium</i> sp.	1(0.9)	-	-	2(1.7)	3(-)	-	-	-

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 178

Table 3.2-20

Numbers and milligrams of biomass, * (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQ1-11A1, September 1976. Dashes indicate species not present or no weight measurement made.

Date	7 September				21 September			
Time	0850				0855			
Air Temp. (C)	15.0				17.5			
Water Temp. (C)	19.5				19.0			
Dissolved Oxygen (ppm)	9.8				8.4			
pH	7.8				8.0			
Secchi Disc (cm)	27.9				30.5			
River Stage (ft)	3.5				4.1			
Substrate	Mud, Detritus, Some Coal, Gravel				Mud, Detritus, Some Coal, Gravel			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	2(0.7)	3(1.0)	11(3.5)	12(4.2)	5(2.2)	1(0.9)	1(0.7)	6(1.7)
Nematoda	5(0.5)	-	13(0.8)	1(-)	1(-)	4(-)	13(0.8)	8(0.9)
<i>Urnatella gracilis</i>	-	-	-	-	-	-	P*	-
<i>Limnodrilus hoffmeisteri</i>	58(10.4)	206(27.1)	132(23.1)	142(26.8)	145(13.8)	21(3.3)	10(1.1)	13(1.7)
<i>Limnodrilus</i> spp.	303(54.5)	356(46.8)	279(48.8)	371(70.1)	726(68.9)	157(24.5)	139(14.9)	306(42.6)
<i>Pelosciolex multisetosus</i>	-	-	-	-	1(-)	-	-	-
<i>Helobdella elongata</i>	-	-	1(1.4)	-	-	1(2.0)	1(0.8)	4(3.1)
<i>Helobdella</i> sp.	-	-	-	-	-	-	2(-)	-
Erpobdellidae	1(6.0)	2(18.1)	-	-	-	1(11.4)	-	2(3.5)
<i>Gammarus fasciatus</i>	-	2(1.1)	1(-)	2(0.2)	-	-	2(0.2)	-
<i>Corophus spiniceps</i>	-	-	-	-	-	1(-)	-	-
<i>Procladius</i> sp.	-	-	1(-)	-	3(-)	4(-)	2(-)	2(-)
<i>Chironomus decorus</i>	-	1(-)	-	-	8(0.4)	-	-	-
<i>C. decorus</i> pupa	-	-	1(-)	-	-	-	-	-
<i>Cryptochironomus</i> nr. <i>fulvus</i>	-	1(-)	1(-)	-	-	3(-)	2(-)	1(-)
<i>Paratubificoides</i> sp.	-	-	1(-)	-	-	-	-	1(-)
<i>Polypedilum halterale</i>	-	1(-)	1(-)	-	-	-	-	-
<i>Physa</i> sp.	-	-	-	-	-	1(-)	-	-
<i>Ferrissia</i> spp.	-	-	-	-	-	1(-)	-	-
<i>Goniobasis virginica</i>	-	1(4.1)	14(281.2)	-	6(193.9)	13(355.0)	15(315.5)	15(539.9)
<i>Annicola limosa</i>	-	-	-	-	1(-)	-	-	1(-)
<i>Campelema decisa</i>	-	-	-	-	1(-)	-	-	-
<i>Pisidium</i> sp.	-	20(4.4)	58(36.1)	-	22(-)	14(-)	13(-)	56(14.1)
<i>Sphaerium</i> sp.	-	-	4(4.4)	-	8(-)	1(-)	1(2.1)	1(-)

*Quantization not possible; denotes organism present.

POOR ORIGINAL

1565 179

Table 3.2-21

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A1, October 1976. Dashes indicate species not present or no weight measurement made.

Date	5 October				19 October			
Time	0914				0855			
Air Temp. (C)	13.0				3.0			
Water Temp. (C)	16.0				9.0			
Dissolved Oxygen (ppm)	11.0				12.4			
pH	7.9				7.6			
Secchi Disc (cm)	22.9				25.4			
River Stage (ft)	4.8				5.3			
Substrate	Mud, Sand, Detritus				Mud, Detritus, Some Sand			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	34(12.4)	3(1.5)	9(5.2)	7(2.8)	2(0.6)	1(-)	1(-)	3(1.0)
Nemertinea	-	-	-	-	1(-)	1(-)	3(-)	-
Nematoda	7(0.9)	1(-)	-	5(-)	13(0.5)	4(-)	5(-)	4(-)
<u>Plumatella repens</u>	P*	-	-	-	-	-	-	-
<u>Limnodrilus hoffmeisteri</u>	11(2.1)	29(3.9)	20(2.5)	57(6.4)	17(2.6)	65(10.8)	-	-
<u>Limnodrilus</u> spp.	204(38.2)	329(40.1)	185(23.5)	421(46.9)	486(73.1)	906(149.9)	471(61.3)	936(130.2)
<u>Pelosciolex multisetaeus</u>	-	-	-	-	-	-	1(-)	1(-)
<u>Helobdella elongata</u>	1(0.7)	-	1(0.8)	-	3(1.8)	-	1(1.6)	1(0.5)
<u>Helobdella</u> sp.	4(-)	5(-)	1(-)	-	-	-	1(-)	-
<u>Erpobdellidae</u>	-	2(46.8)	-	-	1(16.7)	-	-	-
<u>Mooreobdella microstoma</u>	-	-	-	-	-	1(29.5)	-	-
<u>Hydracarina</u>	-	-	-	-	-	-	1(-)	-
<u>Gammarus fasciatus</u>	13(6.0)	1(0.5)	-	-	23(8.1)	20(5.5)	10(1.5)	33(10.4)
<u>Hexagenia limbata</u>	-	-	-	-	2(40.0)	1(17.1)	-	-
<u>Hydropsyche phalerata</u>	1(0.6)	-	-	-	-	-	-	-
<u>Procladius</u> sp.	1(-)	-	-	-	-	-	-	-
<u>Cryptochironomus</u> nr. <u>fulvus</u>	1(-)	3(-)	-	4(-)	-	3(-)	-	1(-)
<u>Polypedium scalaenum</u> sp.	-	-	-	-	-	1(-)	-	-
<u>Physa</u> sp.	2(-)	-	-	-	-	-	-	-
<u>Confobasis virginica</u>	26(552.4)	10(290.9)	7(140.8)	6(345.3)	5(208.3)	1(71.0)	3(187.2)	-
<u>Annicola limosa</u>	1(-)	-	-	1(-)	-	-	-	1(-)
<u>Pisidium</u> sp.	7(-)	15(10.2)	20(9.2)	32(3.2)	11(-)	8(-)	6(-)	11(-)
<u>Sphaerium</u> sp.	-	2(-)	-	-	4(-)	1(-)	-	3(-)

* Quantization not possible; denotes organism present.

POOR ORIGINAL

1565 180

Table 3.2-22

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A2, April 1975. Dashes indicate species not present or no weight measurement made.

Date	6 April				20 April			
Time	1019				1026			
Air Temp. (C)	20.0				25.0			
Water Temp. (C)	8.0				21.0			
Dissolved Oxygen (ppm)	9.9				8.6			
pH	8.4				7.6			
Secchi Disc (cm)	33.0				76.2			
River Stage (ft)	6.6				4.7			
Substrate	Mud, Detritus, Some Coal				Mud, Coal, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	6(2.2)	1(0.7)	-	-	-
Uremertinea	-	2(0.4)	-	3(-)	-	-	-	-
Nematoda	-	-	-	-	1(-)	-	-	-
<u>Nais bretscheri</u>	-	-	-	-	2(-)	-	-	-
<u>N. variabilis</u>	2(-)	-	-	-	-	-	-	-
<u>Branchiura sowerbyi</u>	-	-	-	-	-	-	-	-
<u>Limnodrilus claparedeianus</u>	2(-)	-	-	3(1.4)	9(4.9)	-	1(0.8)	-
<u>L. hoffmeisteri</u>	7(-)	42(32.5)	4(-)	11(5.3)	19(10.4)	3(-)	4(-)	19(13.7)
<u>L. udekemianus</u>	-	-	2(-)	4(1.9)	10(5.5)	2(-)	-	-
<u>Limnodrilus spp.</u>	3(-)	31(24.0)	5(-)	18(8.6)	-	4(-)	5(-)	10(7.2)
<u>Pelosciolex multisetosus</u>	1(-)	-	-	-	-	-	-	-
<u>Mooreobdella microstoma</u>	-	-	-	-	-	-	-	-
<u>Gammarus fasciatus</u>	-	-	1(5.8)	-	1(-)	-	-	1(-)
<u>Cheumatopsyche sp.</u>	-	-	-	1(0.6)	-	-	-	1(-)
<u>Chironomus decorus</u>	-	-	-	1(-)	-	-	-	-
<u>Phaenopsectra nr. ohedians</u>	-	-	-	-	1(-)	-	-	-
<u>Goniobasis virginica</u>	-	-	-	-	10(204.8)	-	1(20.9)	6(90.1)
<u>Sphaerium sp.</u>	-	-	-	1(-)	-	-	-	-

POOR ORIGINAL

1565 181

Table 3.2-23

Numbers and milligrams of biomass, = (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A2, May 1976. Dashes indicate species not present or no weight measurement made.

Date	4 May				18 May			
Time	1040				1057			
Air Temp. (C)	NA				19.0			
Water Temp. (C)	11.8				19.8			
Dissolved Oxygen (ppm)	8.9				7.0			
pH	7.6				NA			
Secchi Disc (cm)	71.1				35.6			
River Stage (ft)	4.9				4.6			
Substrate	Mud, Detritus				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	1(0.4)	-	-	-	1(0.2)	-	-
Nemertinea	-	1(-)	-	-	1(-)	-	-	-
Nematoda	1(-)	-	2(-)	2(-)	1(-)	1(-)	-	-
Enchytraeidae	-	-	1(-)	-	-	-	-	-
Chaetocaster diaphanus	-	-	-	-	-	-	-	-
Nais bartscheri	-	-	-	-	-	-	4(-)	-
N. elinguis	2(-)	5(-)	-	2(-)	-	3(-)	1(-)	-
N. var.abilis	-	3(-)	2(-)	-	4(-)	3(-)	3(-)	-
Slavina appendiculata	2(-)	4(-)	-	1(-)	1(-)	-	2(-)	-
Branchiura sowerbyi	-	-	-	-	-	-	1(-)	-
Limnophilus clapparedeianus	-	-	1(4.4)	-	-	-	-	-
L. hoffmeisteri	-	-	-	-	-	-	28(5.7)	-
L. pfeifferianus	13(1.8)	48(6.9)	33(10.1)	-	10(3.6)	65(11.9)	98(11.8)	80(25.4)
Limnophilus spp.	13(1.8)	-	-	-	-	22(4.0)	47(8.5)	-
Pelocoxys multisetosus	99(12.5)	249(35.9)	89(27.3)	39(6.3)	113(40.1)	524(96.1)	196(39.7)	498(157.9)
Tubificax tubificax	-	4(1.2)	-	-	-	-	-	-
Imm. tub./cap. chaetae*	-	-	-	-	-	1(-)	1(-)	-
Helethella elongata	-	-	-	-	2(-)	-	1(-)	2(-)
Helethellidae	1(0.8)	-	-	-	-	-	-	1(0.7)
Monobdella microstoma	-	-	-	-	-	-	-	-
Orchodactylus fasciatus	3(-)	2(-)	1(0.1)	-	1(19.4)	-	-	-
Orthocladus sp.	-	2(-)	-	-	-	2(0.4)	2(0.5)	-
Orthocladus spp.	-	1(-)	-	-	3(-)	-	4(-)	2(-)
Orthocladus (Orthocladus) sp.	-	-	-	-	-	-	-	-
Chironomus decorus	3(-)	38(2.7)	6(-)	1(-)	-	-	-	-
C. decorus puna	-	-	-	-	4(-)	50(40.9)	21(14.4)	58(51.2)
Chironomus nr. fluvius	1(-)	-	-	-	-	1(-)	-	1(-)
Chironomus nr. obediens	-	-	-	1(-)	1(-)	-	-	-
Polypedilum halterale	1(-)	-	-	-	-	-	1(-)	-
Limnodynastes sp.	-	1(-)	-	-	2(-)	-	-	-
Conchobasis virginica	4(53.7)	1(16.3)	-	-	1(53.8)	-	4(142.4)	2(28.5)
Pisidium sp.	-	3(-)	3(4.0)	1(-)	-	1(-)	-	-
Sphaerium sp.	-	-	-	-	-	-	1(-)	-

NA = Not Available

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 182

Table 3.2-24

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A2, June 1976. Dashes indicate species not present or no weight measurement made.

Date	1 June				15 June			
Time	0900				0835			
Air Temp. (C)	22.0				23.0			
Water Temp. (C)	19.0				23.0			
Dissolved Oxygen (ppm)	9.2				7.0			
pH	7.8				7.8			
Secchi Disc (cm)	80.4				40.6			
River Stage (ft)	5.1				4.1			
Substrate	Mud, Detritus, Some Coal, Some Gravel				Mud, Detritus, Some Coal			
Replicates	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	-	1(0.8)	-	-	-
Nematoda	4(-)	-	6(-)	1(-)	-	2(-)	2(-)	1(-)
<i>Plumatella repens</i>	-	-	P*	-	-	-	-	-
<i>Arctonais limoni</i>	-	-	1(-)	-	-	-	-	-
<i>Nais breitscheri</i>	4(-)	-	5(-)	4(-)	-	-	-	-
<i>N. elinguis</i>	-	-	8(-)	2(-)	-	-	-	-
<i>N. variabilis</i>	-	-	3(-)	-	-	-	-	-
<i>Paranais frici</i>	-	-	3(-)	-	-	-	-	-
<i>Slavina appendiculata</i>	-	-	4(-)	-	-	-	-	-
<i>Ilvodrillus temietoni</i>	1(-)	-	1(-)	-	-	-	-	-
<i>Limnodrilus clavigerianus</i>	11(3.0)	-	77(18.2)	133(28.7)	12(5.5)	10(3.3)	-	19(6.0)
<i>L. hoffmeisteri</i>	42(11.4)	83(16.4)	115(27.2)	300(64.7)	37(17.0)	21(6.9)	64(41.7)	130(40.8)
<i>L. udekemianus</i>	-	-	-	-	-	-	-	18(5.6)
<i>Limnodrilus</i> spp.	189(51.4)	691(136.3)	269(63.6)	699(150.7)	261(120.1)	63(20.8)	52(33.9)	278(87.2)
<i>Peloscotlex ferox</i>	-	-	3(-)	1(-)	-	-	-	-
<i>P. multiseorus</i>	-	-	3(-)	-	-	2(0.3)	1(0.4)	1(0.3)
<i>Tubifex tubifex</i>	-	-	3(-)	-	-	-	-	-
Imm. tub./cap. chaetae**	-	3(-)	3(-)	-	-	-	-	-
Megadrile	-	2(-)	3(-)	1(-)	-	-	4(-)	-
<i>Helobdella elongata</i>	-	-	-	-	-	-	-	-
Erpobdellidae	1(0.3)	-	1(0.5)	-	3(4.6)	-	-	1(0.2)
<i>Gammarus fasciatus</i>	1(-)	1(-)	1(-)	4(0.9)	5(3.2)	2(1.6)	1(0.6)	2(3.7)
<i>Procladius</i> sp.	-	1(-)	3(-)	3(-)	-	3(-)	9(3.5)	10(0.9)
<i>Cricotopus</i> spp.	-	-	2(-)	-	-	-	2(-)	6(0.6)
<i>Orthocladus</i> (<i>Orthocladus</i>) sp.	1(-)	-	-	-	-	-	-	-
<i>Chironomus decorus</i>	20(22.2)	71(35.4)	17(8.3)	3(8.5)	329(81.8)	41(13.8)	17(4.1)	72(25.3)
<i>C. decorus</i> pupa	1(1.3)	2(2.1)	-	1(-)	-	1(-)	-	-
<i>Cryptochironomus</i> nr. <i>fulvus</i>	-	-	1(-)	1(-)	1(-)	-	-	-
<i>Phaenopsectra</i> nr. <i>obediens</i>	-	-	1(-)	-	-	-	4(-)	6(2.0)
<i>Polypedium halterale</i>	-	-	-	-	-	-	-	-
<i>Rheotanytarsus</i> sp.	-	-	-	-	-	1(-)	-	2(0.2)
<i>Tanytarsus</i> sp.	-	-	2(-)	-	-	-	1(-)	-
<i>Phrysa</i> sp.	-	-	-	-	-	-	-	2(0.2)
<i>Goniobasis virginica</i>	1(47.0)	-	5(232.9)	-	-	5(232.5)	4(217.8)	10(400.3)
<i>Campelema decisa</i>	-	-	1(-)	-	-	-	-	1(-)
<i>Pisidium</i> sp.	-	1(2.1)	3(-)	1(-)	2(-)	1(-)	-	-
<i>Sphaerium</i> sp.	-	2(2.4)	-	1(4.5)	-	1(-)	2(-)	1(-)

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 183

Table 3.2-25

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-11A2, July 1976. Dashes indicate species not present or no weight measurement made.

Date	6 July				20 July			
Time	0855				0850			
Air Temp. (C)	24.5				23.0			
Water Temp. (C)	24.5				24.5			
Dissolved Oxygen (ppm)	NA				7.3			
pH	8.0				8.3			
Secchi Disc (cm)	58.4				15.2			
River Stage (ft)	4.7				4.3			
Substrate	Mud, Detritus, Some Coal				Mud, Detritus, Some Gravel			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	-	-	-	2(0.2)	-
Nematoda	1(-)	-	1(-)	-	-	-	1(-)	-
<i>Limnodrilus claviger</i>	-	15(5.2)	-	-	-	-	20(3.7)	-
<i>L. hoffmeisteri</i>	68(22.1)	91(31.7)	85(18.7)	62(18.3)	118(20.2)	48(10.8)	20(3.7)	17(2.9)
<i>L. udekemianus</i>	-	-	-	-	20(3.4)	-	102(19.0)	34(5.7)
<i>Limnodrilus</i> spp.	270(87.6)	273(95.0)	556(122.2)	311(91.9)	295(50.6)	337(75.7)	345(64.2)	372(62.9)
<i>Peloscoides multisetosus</i>	3(4.0)	-	-	-	-	-	1(-)	-
imm. tub./cap. chaetae	-	-	-	-	1(-)	-	-	-
Erpobdellidae	-	-	-	-	1(6.5)	1(2.3)	2(120.4)	-
<i>Gammarus fasciatus</i>	-	-	1(-)	-	-	-	5(0.8)	-
<i>Procladius</i> sp.	2(-)	1(-)	3(-)	1(-)	13(3.3)	3(2.1)	4(-)	1(-)
<i>Chironomus decurvus</i>	24(8.6)	17(6.4)	38(16.3)	4(-)	2(0.2)	-	6(1.0)	3(-)
<i>Cryptochironomus</i> sp. <i>fulvus</i>	15(2.9)	5(2.3)	24(3.7)	2(-)	8(0.7)	-	3(-)	4(-)
<i>Paratubificoides</i> sp.	3(-)	-	-	-	2(0.2)	-	2(-)	-
<i>Polypodium palustrale</i>	-	1(-)	-	-	-	-	1(-)	5(-)
<i>P. haltere</i> e pupa	-	-	-	-	-	-	1(-)	-
<i>P. scalaenum</i>	1(-)	-	-	-	-	-	1(-)	-
<i>Tanytarsus</i> sp.	-	-	-	-	-	-	1(-)	-
<i>Goniodontia viridula</i>	4(166.2)	3(93.0)	3(92.0)	-	4(75.3)	1(43.1)	6(194.3)	1(68.2)
<i>Amnicola limosa</i>	-	-	-	-	-	-	1(-)	-
<i>Pisidium</i> sp.	2(-)	5(2.1)	4(-)	-	-	-	14(-)	1(-)
<i>Sphaerium</i> sp.	-	-	9(3.3)	-	1(-)	-	1(2.0)	4(2.6)

NA = Not Available.

* Immature tubificoid with capilliform chaetae.

POOR ORIGINAL

1565 184

Table 3.2-26

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQ1-11A2, August 1976. Dashes indicate species not present or no weight measurement made.

	3 August				17 August			
Date	0918				0915			
Time	21.0				19.0			
Air Temp. (C)	21.0				19.0			
Water Temp. (C)	7.8				7.8			
Dissolved Oxygen (ppm)	7.9				8.2			
pH	30.5				15.2			
Secchi Disc (cm)	4.1				4.5			
River Stage (ft)								
Substrate	Mud, Detritus				Mud, Detritus, Some Sand, Coal			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	4(1.3)	3(0.8)	10(2.7)	2(0.4)	2(0.3)	-	3(0.4)	5(1.5)
Nematoda	-	-	2(-)	3(-)	2(-)	-	6(-)	2(-)
<u>Urematella gracilis</u>	-	-	-	-	P*	-	-	-
<u>Limnodrilus hoffmeisteri</u>	70(21.4)	31(11.6)	81(20.1)	41(9.5)	58(15.9)	60(18.5)	123(33.0)	72(15.1)
<u>L. udekerianus</u>	10(3.1)	-	-	-	-	-	-	-
<u>Limnodrilus</u> spp.	141(43.1)	103(38.6)	153(38.0)	102(23.7)	59(16.1)	30(9.3)	246(66.0)	104(21.9)
<u>Pelosciolex pulchrisetosus</u>	-	-	2(-)	1(-)	-	-	1(-)	1(-)
imm. tub./cap. chaetae**	2(-)	-	-	-	-	-	2(-)	-
<u>Helobdella elongata</u>	2(2.0)	-	1(0.2)	-	4(2.4)	1(0.5)	1(0.3)	-
<u>H. lineata</u>	-	-	-	-	-	-	-	2(-)
Erpobdellidae	-	-	1(5.7)	-	-	1(0.4)	1(11.4)	1(2.5)
<u>Gammarus fasciatus</u>	2(2.1)	-	-	-	5(0.9)	5(4.0)	26(1.1)	11(3.1)
<u>Hexagenia limbata</u>	-	-	-	-	1(0.2)	-	-	-
<u>Procladius</u> sp.	-	3(0.8)	13(1.7)	2(0.4)	5(1.5)	12(2.3)	17(3.2)	11(2.4)
<u>Coelotanypus scapularis</u>	-	-	-	1(0.2)	-	-	-	-
<u>Chironomus decorus</u>	8(1.8)	7(1.1)	5(0.8)	3(0.3)	1(-)	1(-)	-	1(-)
<u>Cryptochironomus</u> nr. <u>blarina</u>	-	1(0.3)	-	-	-	-	-	-
<u>C. nr. fulvus</u>	12(4.5)	2(0.5)	5(-)	2(0.4)	-	-	1(-)	1(-)
<u>Paracladopelma</u> sp.	-	-	-	1(0.2)	-	-	-	-
<u>Polypedilum palterale</u>	2(0.8)	-	-	-	-	-	-	-
<u>Physo</u> sp.	-	-	-	-	-	1(-)	-	-
<u>Goniobasis virginica</u>	2(150.9)	3(156.9)	5(178.5)	1(103.8)	6(248.6)	6(342.0)	3(116.6)	11(511.3)
<u>Amnicola limosa</u>	-	-	1(-)	-	-	-	1(-)	-
<u>Camelona decisa</u>	-	-	-	-	1(-)	-	-	-
<u>Pisidium</u> sp.	6(4.4)	9(-)	5(3.3)	4(-)	2(-)	1(-)	7(-)	5(-)
<u>Sphaerium</u> sp.	1(1.0)	2(-)	1(1.2)	1(-)	-	-	-	1(-)

* Quantization not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 185

Table 3.2-27

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station ~~TX~~-AQ1-11A2, September 1976. Dashes indicate species not present or no weight measurement made.

Date	7 September				21 September			
Time	0904				0906			
Air Temp. (°C)	15.0				17.0			
Water Temp. (°C)	19.0				19.0			
Dissolved Oxygen (ppm)	9.0				7.6			
pH	7.9				8.1			
Secchi Disc (cm)	30.5				30.5			
River Stage (ft)	3.5				4.1			
Substrate	Mud, Detritus, Some Sand				Mud, Detritus, Some Sand, Coal			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	23(9.2)	30(10.3)	7(2.0)	-	6(2.1)	-	3(1.2)	-
Nemertinea	-	-	-	-	1(-)	-	-	-
Nematoda	4(-)	2(-)	-	1(-)	2(-)	-	3(-)	1(-)
<u>Ureatella gracilis</u>	-	-	-	-	-	P*	-	-
<u>Limnodrilus hoffmeisteri</u>	110(19.5)	219(32.8)	20(3.4)	176(9.7)	20(3.5)	61(10.8)	51(9.5)	41(7.9)
<u>L. udekemianus</u>	-	-	-	59(3.2)	10(1.8)	-	-	21(4.2)
<u>Limnodrilus</u> spp.	141(25.0)	379(56.9)	556(94.9)	646(35.5)	219(38.4)	173(30.7)	251(44.4)	62(12.0)
<u>Poloscolex multisetosus</u>	-	1(-)	-	-	6(0.4)	-	-	2(-)
<u>Helobdella elongata</u>	-	1(0.8)	-	-	1(0.5)	-	1(0.5)	1(1.6)
<u>Helobdella</u> sp.	1(-)	-	-	-	8(-)	6(-)	-	-
<u>Placobdella montifera</u>	1(-)	-	-	-	-	-	-	-
Erpobdellidae	-	3(0.9)	1(30.2)	-	-	-	-	-
<u>Gammarus fasciatus</u>	11(7.7)	13(3.0)	-	-	52(18.2)	35(15.6)	1(1.2)	1(1.7)
<u>Procladius</u> sp.	7(1.4)	7(2.1)	-	-	2(0.3)	3(-)	1(1.1)	1(-)
<u>Coelotanypus scapularis</u>	-	-	-	-	2(0.3)	-	-	-
<u>Chironomus decorus</u>	1(-)	4(0.4)	-	-	1(-)	-	3(-)	-
<u>Cryptochironomus</u> nr. <u>fulvus</u>	1(-)	3(-)	11(2.9)	1(-)	3(-)	3(-)	-	-
<u>Paratadiopelma</u> sp.	-	-	-	-	2(-)	-	1(-)	-
<u>Polypedium scalanum</u>	-	-	2(0.5)	1(-)	-	-	-	-
<u>Physa</u> sp.	-	-	-	-	-	3(-)	-	-
<u>Ferrissia</u> spp.	-	-	-	1(-)	-	1(-)	-	-
<u>Goniobasis virginica</u>	10(460.4)	8(257.1)	-	3(67.1)	23(301.8)	28(1660.1)	11(554.5)	5(293.4)
<u>Ammicola limosa</u>	1(-)	4(-)	-	-	1(-)	7(-)	1(-)	1(-)
<u>Campeloma decisa</u>	-	-	-	1(-)	-	2(-)	-	-
<u>Pisidium</u> sp.	15(-)	40(20.7)	63(7.1)	9(-)	17(-)	47(-)	5(-)	1(-)
<u>Sphaerium</u> sp.	4(-)	5(-)	-	-	3(-)	13(-)	-	-

* Quantization not possible; denotes organism present.

POOR ORIGINAL

1565 186

Table 3.2-28

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQ1-11A2, October 1976. Dashes indicate species not present or no weight measurement made.

Date	5 October				19 October			
Time	0937				0909			
Air Temp. (C)	13.5				4.0			
Water Temp. (C)	15.5				9.0			
Dissolved Oxygen (ppm)	9.4				11.8			
pH	8.0				7.6			
Secchi Disc (cm)	27.9				33.0			
River Stage (ft)	4.8				5.3			
Substrate	Mud, Detritus				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
<i>Turbellaria</i>	6(2.8)	2(1.0)	4(3.0)	-	2(1.3)	5(1.6)	3(0.1)	1(1.0)
<i>Nemertinea</i>	-	-	2(-)	-	-	-	1(-)	1(-)
<i>Nematoda</i>	1(-)	1(-)	4(-)	2(-)	-	2(-)	-	2(-)
<i>Limnodrilus hoffmeisteri</i>	35(4.4)	30(5.6)	-	10(1.6)	-	-	-	58(7.2)
<i>L. udekemianus</i>	-	-	-	-	10(2.3)	32(6.9)	-	-
<i>Limnodrilus</i> spp.	259(32.6)	209(39.2)	228(39.6)	200(31.7)	220(49.7)	365(78.3)	422(88.2)	521(64.9)
<i>Pelosciolex multisetosus</i>	-	1(-)	-	-	-	1(-)	-	-
<i>Helobdella elongata</i>	-	-	-	-	1(0.7)	-	2(0.6)	1(1.0)
<i>Helobdella</i> sp.	6(-)	-	2(-)	-	-	4(-)	4(-)	-
<i>Erpobdellidae</i>	1(20.3)	-	-	-	-	-	-	-
<i>Gammarus fasciatus</i>	2(1.0)	-	-	-	3(0.1)	15(10.4)	22(4.5)	6(3.7)
<i>Hydropsyche phalerata</i>	-	-	-	-	-	-	1(1.3)	1(0.7)
<i>Procladius</i> sp.	1(-)	-	-	-	-	-	-	-
<i>Ceclorhynchus scapularis</i>	-	-	-	1(-)	-	-	1(-)	-
<i>Orthocladius (Orthocladius)</i> sp.	-	1(-)	-	-	-	-	-	-
<i>Chironomus decorus</i>	-	1(-)	-	-	-	1(-)	1(-)	2(-)
<i>Cryptochironomus</i> nr. <i>fulvus</i>	4(-)	5(-)	3(-)	3(-)	-	1(-)	5(-)	2(-)
<i>Phyca</i> sp.	-	-	-	-	-	-	1(-)	-
<i>Ferrissia</i> spp.	-	1(-)	-	-	-	-	-	-
<i>Goniobasis virginica</i>	11(498.3)	12(-)	5(197.2)	1(4.5)	3(56.3)	19(908.0)	8(386.2)	8(372.3)
<i>Limicola limosa</i>	3(-)	5(-)	-	-	-	2(-)	-	1(-)
<i>Pisidium</i> sp.	33(-)	28(-)	12(-)	9(-)	5(-)	22(3.4)	-	15(-)
<i>Sphaerium</i> sp.	-	-	1(-)	-	1(-)	-	-	-

POOR ORIGINAL

1565 187

Table 3.2-29

Numbers and milligrams of biomass, " (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-981, April 1976. Dashes indicate species not present or no weight measurement made.

Date	6 April				20 April			
Time	1036				1044			
Air Temp. (C)	19.5				25.0			
Water Temp. (C)	8.0				21.0			
Dissolved Oxygen (ppm)	9.7				8.2			
pH	8.4				7.4			
Secchi Dis. (cm)	45.7				94.0			
River Stage (ft)	6.6				4.7			
Substrate	Mud, Detritus, Some Coal				Mud, Coal, Detritus			
Replicate	A	B	C	D	A	B	C	D
Nematoda	-	1(-)	1(-)	-	-	-	-	-
Enchytraeidae	1(-)	-	-	-	-	-	-	-
<u>Nais variabilis</u>	-	-	1(-)	-	-	1(-)	-	-
<u>Limnodrilus claredeianus</u>	1(-)	-	-	-	-	-	-	-
<u>L. hoffmeisteri</u>	11(-)	10(4.4)	21(11.1)	20(15.4)	20(17.8)	39(12.2)	56(29.9)	38(17.9)
<u>L. udekemianus</u>	3(-)	-	-	-	10(8.9)	-	7(3.7)	-
<u>Limnodrilus</u> spp.	10(-)	19(8.4)	32(16.9)	39(30.1)	10(8.9)	29(9.1)	56(29.9)	10(4.7)
<u>Pelecocox multisetosus</u>	1(-)	-	-	-	-	-	-	1(-)
<u>Tubifex tubifex</u>	-	-	-	-	-	-	1(-)	-
Imm. tub./cap. chaetae*	1(-)	-	-	-	-	-	1(-)	-
<u>Optioservus</u> sp.	-	-	-	-	-	-	-	1(-)
<u>Procladius</u> sp.	-	-	1(-)	-	-	-	-	-
<u>Phaenopsectra</u> sp.	-	-	-	-	1(-)	-	-	-
<u>Tanytarsus</u> sp.	-	-	-	-	-	-	1(-)	-
<u>Goniobasis virginica</u>	-	-	-	-	-	-	1(68.5)	-
<u>Pisidium</u> sp.	-	-	1(-)	-	-	-	1(-)	2(-)

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 188

Table 3.2-30

Numbers and milligrams of biomass, \pm (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-981, May 1976. Dashes indicate species not present or no weight measurement made.

Date	4 May				18 May			
Time	1102				1012			
Air Temp. (C)	9.0				19.0			
Water Temp. (C)	11.8				19.8			
Dissolved Oxygen (ppm)	8.6				6.8			
pH	7.6				NA			
Secchi Disc (cm)	71.1				35.6			
River Stage (ft)	4.9				4.6			
Substrate	Mud, Detritus				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
<i>Nematoda</i>	-	-	-	-	-	-	1(-)	-
<i>Manayunkia speciosa</i>	-	-	-	1(-)	-	-	-	-
<i>Chaetopaster diaphanus</i>	1(-)	-	-	-	9(-)	-	-	-
<i>Nais bierschleri</i>	49(0.9)	8(0.1)	1(-)	-	2(-)	-	-	-
<i>N. elinguis</i>	29(0.6)	8(0.1)	1(-)	25(1.1)	1(-)	-	-	-
<i>N. variabilis</i>	49(0.9)	23(0.3)	2(-)	-	-	-	-	1(-)
<i>Nais</i> sp.	-	16(0.2)	-	-	-	1(-)	-	-
<i>Flavina appendiculata</i>	-	-	1(-)	-	-	-	-	-
<i>Limnodrilus templetoni</i>	10(3.2)	-	-	-	-	10(6.3)	-	1(-)
<i>Limnodrilus hoffmeisteri</i>	29(9.3)	33(16.4)	16(6.5)	46(-)	80(35.2)	54(40.0)	41(59.8)	40(20.9)
<i>N. udekemianus</i>	-	-	-	-	-	-	-	10(5.2)
<i>Limnodrilus</i> spp.	48(15.4)	33(16.4)	39(15.8)	34(-)	101(44.5)	32(20.0)	21(30.6)	61(31.9)
<i>Poloscolex multiserpens</i>	-	-	-	-	-	-	-	1(-)
<i>Tubifex rubifex</i>	10(3.2)	1(-)	8(3.2)	-	-	-	-	2(-)
Imm. tub./cap. chaetae*	19(5.1)	-	8(3.2)	1(-)	5(-)	21(13.1)	-	4(-)
<i>Helobdella lineata</i>	-	1(-)	-	-	-	-	-	-
<i>Gerrhus fasciatus</i>	9(1.3)	3(-)	1(-)	1(2.7)	3(0.6)	1(0.1)	-	-
<i>Chaoborus punctipennis</i>	1(0.3)	-	-	-	-	-	-	-
<i>Procladius</i> sp.	-	-	6(0.2)	-	4(-)	7(-)	-	5(-)
<i>Ericotopus</i> spp.	-	2(-)	-	-	-	-	-	-
<i>Orthocladus (Orthocladus)</i> sp.	-	-	-	1(-)	-	-	-	-
<i>Chironomus decorus</i>	76(19.5)	32(9.8)	77(8.8)	33(10.3)	309(111.5)	286(194.5)	90(82.8)	325(212.1)
<i>C. decorus</i> pupa	-	-	-	-	5(4.2)	5(3.3)	2(-)	10(6.5)
<i>Paratendipes</i> sp.	3(0.2)	-	-	-	-	-	-	-
<i>Goniodontia virginica</i>	1(42.1)	1(46.8)	-	-	-	-	-	-
<i>Pisidium</i> sp.	-	-	-	1(2.1)	-	-	-	-

NA = Not Available.

* Immature tubificid with capilliform chaetae.

POOR ORIGINAL

1565 189

Table 3.2-31

Numbers and milligrams of biomass, \pm (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-981, June 1976. Dashes indicate species not present or no weight measurement made.

Date	1 June				15 June			
Time	0917				0850			
Air Temp. (C)	21.5				23.0			
Water Temp. (C)	19.0				23.0			
Dissolved Oxygen (ppm)	9.0				8.0			
pH	7.9				7.5			
Secchi Disc (cm)	94.0				55.9			
River Stage (ft)	5.1				4.1			
Substrate	Mud, Detritus, Coal				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	-	-	-	-	-	-	-	-
Nematoda	6(-)	1(-)	1(-)	5(-)	4(-)	1(-)	-	-
Enchytraeidae	-	1(-)	-	-	-	4(-)	3(-)	-
<u>Arctonais lomondi</u>	4(-)	-	8(-)	2(-)	-	-	-	-
<u>Nais bretscheri</u>	-	-	-	-	-	3(-)	-	-
<u>N. alinguis</u>	1(-)	-	-	-	-	1(-)	-	-
<u>N. variabilis</u>	1(-)	1(-)	-	-	-	-	-	-
<u>Paranais friei</u>	-	1(-)	-	28(0.6)	-	2(-)	-	-
<u>Branchiura sowerbyi</u>	-	-	-	-	-	-	-	-
<u>Hydrodillus templetoni</u>	-	-	-	-	-	-	1(-)	1(-)
<u>Hydrodillus hoffmeisteri</u>	20(5.9)	92(32.7)	174(40.9)	50(13.2)	588(67.0)	13-(26.8)	286(39.5)	154(28.3)
<u>Hydrodillus</u> spp.	111(32.7)	58(20.6)	289(68.0)	99(26.1)	932(106.2)	345(68.9)	246(36.9)	201(27.6)
<u>Peloscoides multisetosus</u>	-	-	-	1(0.7)	1(-)	-	-	-
<u>Tubifex tubifex</u>	1(-)	-	1(-)	1(-)	-	-	-	-
Imm. tub./cap. chaetae*	6(-)	4(-)	2(-)	5(-)	2(-)	5(0.3)	1(-)	3(-)
<u>Gammarus fasciatus</u>	3(1.4)	2(0.9)	1(-)	4(1.0)	3(1.8)	21(1.3)	7(-)	3(-)
<u>Caprellid</u> sp.	1(0.5)	-	-	-	-	6(2.5)	6(6.9)	2(3.3)
<u>Stenelmis</u> spp.	1(0.7)	-	1(0.5)	-	1(0.4)	-	-	-
<u>Procladius</u> sp.	10(-)	4(-)	-	1(-)	45(10.9)	2(1.9)	2(2.0)	-
<u>Chironomus decorus</u>	106(114.4)	84(92.5)	196(224.6)	144(140.9)	409(216.7)	56(6.8)	52(5.8)	37(21.5)
<u>C. decorus</u> pupa	8(6.9)	2(-)	15(14.1)	4(1.2)	326(151.2)	512(178.5)	232(116.5)	5(-)
<u>Cryptochironomus</u> nr. <u>fulvus</u>	-	-	1(-)	-	2(1.3)	3(3.6)	9(8.2)	-
<u>Polypedilum nalterale</u>	-	-	-	-	2(0.1)	-	-	-
<u>Tanytarsus</u> sp.	-	-	-	-	3(0.2)	1(-)	-	1(-)
<u>Capitobasis virginica</u>	1(60.4)	-	-	1(54.8)	2(0.1)	1(-)	-	-
<u>Pisidium</u> sp.	-	-	-	-	-	2(87.2)	2(99.7)	2(86.3)
<u>Sphaerium</u> sp.	1(-)	-	-	-	2(-)	4(-)	1(-)	-
					4(-)	-	-	1(-)

* Immature tubificid with capilliform chaetae.

1565 190

POOR ORIGINAL

Table 3.2-32

Numbers and milligrams of biomass, \pm (sg), of benthic organisms per Ponar grab (524 cm²) at station IM-AQ1-451, July 1976. Dashes indicate species not present or no weight measurement made.

Date	6 July				7 July			
Time	0910				0910			
Air Temp. (C)	24.5				24.0			
Water Temp. (C)	24.5				24.5			
Dissolved Oxygen (ppm)	NA				6.9			
pH	7.9				8.3			
Secchi Disc (cm)	61.0				30.5			
River Stage (ft)	4.7				4.3			
Substrate	Mud, Detritus				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	1(-)	2(0.2)	-	-	6(1.0)	1(-)	4(0.7)	-
Nematoda	1(-)	-	1(-)	-	-	1(-)	-	1(-)
<i>Plumatella repens</i>	-	-	-	-	-	-	Pos*	-
<i>Branchiura sowerbyi</i>	-	1(10.5)	-	-	-	-	-	-
<i>Hydrinus complanatus</i>	-	1(-)	-	1(-)	-	-	-	-
<i>Limnodynastes clausenianus</i>	-	-	-	-	-	15(2.7)	-	-
<i>L. hoffmanni</i>	65(17.6)	86(25.0)	90(20.2)	85(30.2)	36(6.0)	92(16.5)	31(11.6)	11(5.1)
<i>L. wickhami</i>	-	-	-	-	-	-	10(3.7)	21(11.6)
<i>Limnodynastes</i> spp.	98(25.3)	327(95.0)	231(51.8)	207(61.4)	265(44.6)	245(43.9)	143(53.3)	85(27.1)
<i>Polydora multisetosa</i>	2(0.1)	1(-)	1(-)	1(-)	1(-)	1(-)	-	-
Imm. tub./cap. chaetae***	1(-)	7(-)	2(-)	2(-)	-	-	2(-)	-
Megadrile	-	-	-	-	-	1(-)	-	1(-)
<i>Helobdella elongata</i>	1(0.5)	-	-	-	-	-	1(0.2)	-
<i>Ceratonereis fasciatus</i>	-	-	-	-	1(0.2)	-	-	-
<i>Hexapoda loricata</i>	-	-	-	-	1(5.8)	-	-	-
<i>Procladius</i> sp.	7(-)	5(-)	2(-)	5(-)	12(3.0)	14(2.7)	14(2.8)	3(-)
<i>Chironomus tentans</i>	5(-)	17(1.7)	8(3.5)	3(-)	33(5.5)	65(2.5)	49(5.4)	8(1.1)
<i>Cryptochironomus</i> nr. <i>fulvus</i>	2(-)	-	3(-)	-	5(0.8)	3(-)	-	-
<i>Paratubificoides</i> sp.	1(-)	-	-	-	-	-	-	-
<i>Polynoidium halterale</i>	-	-	-	-	-	1(-)	-	-
<i>Lanvaterus</i> sp.	-	1(-)	-	-	-	-	-	-
<i>Contobasis virginica</i>	3(37.1)	-	-	4(231.9)	2(98.4)	2(101.8)	4(181.4)	-
<i>Ariccola limosa</i>	-	-	-	1(-)	-	-	-	-
<i>Platidius</i> sp.	9(-)	8(-)	3(-)	4(-)	-	6(-)	1(-)	-
<i>Sphaerium</i> sp.	11(-)	-	1(-)	2(4.7)	-	-	1(1.9)	-

NA = Not Available

* Non-quantitative sample.

** Quantitation not possible; denotes organism present.

*** Immature tubificid with capilliform chaetae.

1565 191

POOR ORIGINAL

Table 3.2-13

Numbers and milligrams of biomass, \pm (mg), of benthic organisms per Ponar grab (529 cm²) at Station TM-AQI-981, August 1976. Dashes indicate species not present or no weight measurement made.

Date	3 August				17 August			
Time	0938				0935			
Air temp. (°C)	21.0				19.0			
Water temp. (°C)	21.2				18.5			
Dissolved oxygen (ppm)	8.4				7.7			
pH	8.9				8.0			
Secchi disc (cm)	33.0				20.3			
River stage (ft)	4.1				4.5			
Substrate	Mud, Detritus				Mud, Detritus, Some Coal			
Replicate	A	B	C	D	A	B	C	D
<i>Forbellaria</i>	-	36(0.4)	-	-	4(0.9)	-	2(0.7)	4(0.9)
<i>Vermetidae</i>	-	-	-	-	1(-)	6(-)	1(-)	1(-)
<i>Branchiura</i> <i>suberthol</i>	-	-	-	-	1(-)	-	-	-
<i>Alpheidae</i> <i>holmquisti</i>	-	-	-	-	-	-	-	-
<i>L. alcockianus</i>	39(15.8)	29(8.0)	18(5.1)	20(7.8)	34(4.9)	45(14.8)	87(18.7)	58(6.3)
<i>Alpheidae</i> spp.	-	-	-	-	11(1.6)	-	-	-
<i>Polysyllus</i> <i>multisetosus</i>	117(47.5)	79(21.9)	105(29.9)	80(31.2)	235(34.0)	89(29.2)	186(40.0)	306(33.1)
Imm. tub./sp. chaetae*	1(-)	-	1(-)	1(-)	1(-)	1(-)	1(-)	1(-)
<i>Hexadrille</i>	-	-	-	-	1(-)	-	1(-)	1(-)
<i>Helicella</i> <i>lineata</i>	-	-	-	-	-	1(-)	-	-
<i>Corbicula</i> <i>lineolata</i>	-	1(1.1)	-	-	-	-	-	1(-)
<i>Chironomus</i> <i>monticorensis</i>	-	-	-	-	-	-	-	1(0.3)
<i>Procladius</i> sp.	5(-)	14(4.2)	13(3.7)	10(1.9)	14(2.2)	27(5.3)	15(3.3)	13(4.1)
<i>Chironomus</i> <i>lexington</i>	37(14.7)	19(6.3)	27(14.9)	19(6.8)	29(6.5)	8(3.9)	14(3.3)	21(5.5)
<i>L. dentatus</i> pupa	-	1(-)	-	-	-	-	-	-
<i>Procladius</i> <i>monticorensis</i> nr. <i>fulvus</i>	8(2.4)	6(3.0)	4(4.0)	4(-)	5(-)	1(-)	3(-)	4(3.2)
<i>Procladius</i> spp. pupa	-	-	-	-	1(-)	-	-	-
<i>Polysyllus</i> <i>halgerae</i>	2(-)	-	-	-	2(-)	-	-	-
<i>Polysyllus</i> group sp.	-	-	-	-	1(-)	-	-	2(-)
<i>Gonistacus</i> <i>virginica</i>	-	-	-	-	-	-	-	-
<i>Valvata</i> <i>tricarinata</i>	3(185.0)	7(322.4)	3(131.8)	4(252.6)	2(112.1)	3(173.8)	2(145.3)	5(355.8)
<i>Amphipoda</i> <i>lineata</i>	-	5(-)	-	-	-	-	-	1(-)
<i>Pisidium</i> sp.	11(-)	8(-)	4(-)	11(-)	21(-)	14(-)	4(-)	18(-)
<i>Phaenicia</i> sp.	-	28(-)	29(-)	4(-)	6(-)	1(-)	1(-)	-

* Immature tubificid with capilliform chaetae.

1565 192

Table 3.2-34

Numbers and milligrams of biomass, - (mg), of benthic organisms per Ponar grab (529 cm²) at station 100A, 1-12, September 1976. Dashes indicate species not present or no weight measurement made.

Date	7 September				21 September			
Time	0925				0925			
Air Temp. (C)	15.5				16.5			
Water Temp. (C)	19.0				19.5			
Dissolved Oxygen (ppm)	8.6				7.9			
pH	7.9				8.1			
Secchi Disc (cm)	43.2				30.5			
River Stage (ft)	3.5				4.1			
Substrate	Mud, Detritus				Mud, Detritus			
Replicate	A	B	C	D	A	B	C	D
Turbellaria	8(1.9)	4(0.8)	9(3.0)	1(-)	-	7(3.0)*	6(3.3)	1(-)
Nematoda	-	-	-	-	-	-	1(-)	-
<i>Monacella gracilis</i>	-	-	-	-	-	-	P*	-
<i>Alloporus perplectus</i>	-	-	-	-	-	2(-)	13(1.8)	-
<i>Limnolabus hoffmeisteri</i>	89(16.5)	99(18.1)	36(9.1)	21(6.9)	172(33.2)	104(22.6)	163(25.7)	161(35.1)
<i>A. plicatulus</i>	-	-	-	-	-	15(3.3)	-	-
<i>Limnolabus</i> spp.	155(28.7)	315(57.6)	192(48.7)	150(49.1)	122(23.6)	254(55.2)	290(45.7)	354(78.2)
<i>Polysolen multisetosus</i>	-	-	-	1(-)	-	1(-)	5(-)	3(-)
Imm. tub./cap. chaetae**	1(-)	-	4(-)	1(-)	-	5(-)	7(0.4)	1(-)
<i>Helobdella elongata</i>	-	1(1.2)	-	-	-	-	-	-
<i>Helobdella</i> sp.	-	-	-	1(-)	-	-	-	-
<i>Limnolabus fasciatus</i>	-	-	-	-	-	-	1(0.7)	-
<i>Limnolabus limbae</i>	1(6.5)	-	-	-	-	-	-	1(6.1)
<i>Limnolabus</i> sp.	-	-	1(-)	-	-	-	-	-
<i>Procladius</i> sp.	6(0.4)	2(-)	3(2.1)	12(2.4)	-	3(-)	2(-)	3(0.4)
<i>Asclerodermus scapularis</i>	-	1(-)	-	-	-	-	-	-
<i>Ablabesmyia</i> sp.	2(0.1)	-	-	-	-	-	-	2(0.3)
<i>Chironomus decorus</i>	3(2.7)	-	2(1.4)	-	-	27(2.1)	70(6.5)	12(1.4)
<i>Cryptochironomus</i> nr. <i>folius</i>	-	-	3(2.1)	6(5.0)	-	4(-)	3(-)	6(0.2)
<i>Polyneura halperae</i>	9(0.4)	-	-	-	1(-)	-	-	-
<i>P. scalanum</i> pupa	-	-	-	-	-	1(-)	-	-
<i>Gerris parvus</i>	-	-	-	-	1(-)	-	1(-)	-
<i>Gerris</i> sp.	-	-	-	-	-	-	-	-
<i>Conchosis virginica</i>	3(148.7)	7(324.4)	2(100.3)	2(101.8)	2(108.0)	3(185.6)	7(350.1)	6(267.5)
<i>Malva tricornata</i>	1(-)	1(-)	1(-)	-	-	-	-	2(-)
<i>Amphibia lineata</i>	1(-)	-	3(-)	1(-)	-	1(-)	-	2(-)
<i>Pisidium</i> sp.	2(-)	15(12.0)	6(1.0)	20(-)	4(-)	16(-)	17(-)	18(-)
<i>Sphaerium</i> sp.	1(5.8)	-	-	10(-)	2(-)	8(-)	4(-)	-

* Quantitation not possible; denotes organism present.

** Immature tubificid with capilliform chaetae.

Table 3.2-35

Depth and milligrams of biomass, \pm mg, of benthic organisms per Ponar grab (529 cm²) at station TM-AQ1-931, October 1976. Dashes indicate species not present or no weight measurement made.

Date	5 October				19 October			
Time	1700				0925			
Air temp. (C)	13.5				4.0			
Water temp. (C)	15.5				9.0			
Dissolved oxygen (ppm)	8				11.6			
pH	8.1				7.4			
Secchi disc (cm)	27.9				35.6			
River stage (ft)	4.8				5.3			
Substrate	Mud, detritus				Mud, detritus			
Replicate	A	B	C	D	A	B	C	D
<i>Turbellaria</i>	3(2.5)	3(2.5)	5(4.3)	2(1.7)	-	1(0.3)	-	-
<i>Nematoda</i>	-	-	2(-)	-	1(-)	2(0.6)	-	1(-)
<i>Branchiura owerthofi</i>	-	-	-	-	-	1(-)	-	-
<i>Lysanillus tenuicollis</i>	8(0.7)	1(-)	2(-)	-	-	-	-	1(-)
<i>Lysanillus lativestris</i>	135(26.3)	84(21.4)	116(30.4)	56(12.2)	45(13.0)	-	50(17.8)	70(26.9)
<i>L. subsericans</i>	-	1(0.4)	-	-	1(0.3)	21(7.7)	-	35(13.4)
<i>Lysanillus</i> spp.	422(86.1)	103(112.0)	367(96.2)	389(85.0)	113(30.1)	233(85.6)	149(53.1)	314(120.5)
<i>Polyscolus multisetosus</i>	4(-)	3(1.2)	-	-	-	1(-)	-	7(1.1)
Imm. tub. cap. chaetae*	18(2.0)	-(-)	3(-)	1(-)	-	-	1(-)	2(-)
<i>Helobdella elongata</i>	-	1(1.9)	-	-	-	-	-	1(0.3)
<i>Gammarus fasciatus</i>	-	-	-	-	-	1(-)	-	-
<i>Desmoulinia limbata</i>	-	1(4.9)	-	1(8.6)	-	1(11.3)	1(0.6)	2(24.4)
<i>Desmoulinia</i> sp.	-	-	-	-	-	1(0.4)	-	1(-)
<i>Procladius</i> sp.	3(2.7)	2(-)	4(-)	2(-)	-	1(-)	-	-
<i>Ablabesmus</i> sp.	-	1(-)	-	-	-	-	-	-
<i>Orthocladinae</i>	-	-	-	-	-	-	-	1(-)
<i>Chironomus decorus</i>	47(13.4)	11(2.6)	18(3.5)	22(6.1)	-	1(-)	-	2(-)
<i>Cryptochironomus</i> nr. <i>fulvus</i>	4(0.8)	13(2.2)	2(-)	10(1.5)	-	1(-)	1(-)	2(-)
<i>Chironomus</i> sp.	3(154.5)	3(152.5)	3(241.2)	3(251.7)	1(46.0)	-	-	-
<i>Amphipoda litorea</i>	3(-)	1(-)	2(-)	-	-	-	-	-
<i>Pisidium</i> sp.	7(3.4)	15(-)	12(-)	15(3.0)	-	-	6(-)	3(-)
<i>Caprellid</i> sp.	8(-)	4(-)	-	-	-	22(-)	6(-)	-

* Immature tubificid with capilliform chaetae.

Table 9.2-10

Numbers and percent composition of benthic organisms collected at the macroinvertebrate stations, April through October 1970. Dashes indicate species not present.

	No. of individuals		No. of species		No. of individuals		No. of species		No. of individuals		No. of species	
	N	%	N	%	N	%	N	%	N	%	N	%
Turbellaria	6	0.04	2	0.01	154	0.70	133	0.68	78	0.38	374	0.64
Nemertinea	18	0.18	3	0.02	14	0.08	12	0.06	4	0.02	67	0.12
Nematoda	143	1.45	48	0.65	111	1.43	67	0.88	42	0.54	667	0.74
<i>Nematode species</i>	1	0.01	-	-	-	-	-	-	1	0.01	2	0.01
Intertracidae	3	0.03	-	-	-	-	-	-	1	0.01	2	0.01
<i>Aspidogaster laticollis</i>	52	0.54	1	0.01	1	0.00	1	0.01	17	0.09	32	0.08
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	-	-	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	1	0.01	-	-	1	0.01	24	0.07
<i>Aspidogaster laticollis</i>	3	0.03	1	0.01	1	0.01	-	-	1	0.01	185	0.51
<i>Aspidogaster laticollis</i>	12	0.12	1	0.01	34	0.15	1	0.01	1	0.01	158	0.44
<i>Aspidogaster laticollis</i>	143	1.45	1	0.01	17	0.12	1	0.01	1	0.01	158	0.44
<i>Aspidogaster laticollis</i>	132	1.37	1	0.01	2	0.01	1	0.01	1	0.01	158	0.44
<i>Aspidogaster laticollis</i>	-	-	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	30	0.32	1	0.01	-	-	-	-	1	0.01	23	0.06
<i>Aspidogaster laticollis</i>	46	0.50	1	0.01	2	0.01	1	0.01	1	0.01	19	0.05
<i>Aspidogaster laticollis</i>	-	-	1	0.01	1	0.01	-	-	1	0.01	1	0.01
<i>Aspidogaster laticollis</i>	-	-	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	35	0.38	3	0.02	-	-	-	-	3	0.03	45	0.12
<i>Aspidogaster laticollis</i>	48	0.51	1	0.01	-	-	-	-	1	0.01	152	0.42
<i>Aspidogaster laticollis</i>	-	-	1	0.01	-	-	-	-	-	-	20	0.05
<i>Aspidogaster laticollis</i>	-	-	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	511	5.41	133	1.87	18	0.17	114	1.73	18	0.23	1039	1.17
<i>Aspidogaster laticollis</i>	455	4.82	437	65.04	1450	56.16	184	2.81	1447	64.76	6516	72.16
<i>Aspidogaster laticollis</i>	142	1.48	120	0.74	85	0.24	411	2.1	374	0.87	412	1.01
<i>Aspidogaster laticollis</i>	-	-	-	-	-	-	1	0.01	-	-	1	0.01
<i>Aspidogaster laticollis</i>	23	0.17	32	0.25	31	0.15	12	0.15	1	0.01	184	0.19
<i>Aspidogaster laticollis</i>	374	3.78	17	1.23	32	0.11	13	0.11	1	0.01	714	0.24
<i>Aspidogaster laticollis</i>	-	-	-	-	-	-	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	2	0.02
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	12	0.1
<i>Aspidogaster laticollis</i>	5	0.05	2	0.01	-	-	2	0.02	2	0.02	144	0.16
<i>Aspidogaster laticollis</i>	-	-	2	0.01	123	0.55	14	0.15	5	0.05	6	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	1	0.00	2	0.02	2	0.02	6	0.01
<i>Aspidogaster laticollis</i>	1	0.01	2	0.01	18	0.07	31	0.18	1	0.01	51	0.05
<i>Aspidogaster laticollis</i>	-	-	-	-	-	-	1	0.01	-	-	1	0.01
<i>Aspidogaster laticollis</i>	-	-	-	-	105	0.47	22	0.11	-	-	127	0.14
<i>Aspidogaster laticollis</i>	-	-	-	-	6	0.03	2	0.01	-	-	8	0.1
<i>Aspidogaster laticollis</i>	-	-	-	-	1	0.00	-	-	-	-	5	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	54	0.34	626	2.82	262	1.28	51	0.25	104	0.45	104	0.45
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	4	0.03	6	0.04	2	0.02	1	0.01	1	0.01	24	0.03
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	3	0.01	-	-	-	-	5	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	14	0.14	1	0.01	-	-	1	0.01	2	0.01	14	0.02
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	4	0.04	3	0.02	1	0.00	2	0.01	-	-	12	0.01
<i>Aspidogaster laticollis</i>	6	0.06	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	-	-	1	0.01	-	-	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	1	0.00	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	2	0.01	-	-	2	0.01	-	-	1	0.01	5	0.01
<i>Aspidogaster laticollis</i>	2	0.01	2	0.01	-	-	-	-	-	-	4	0.01
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	-	-	-	-	3	0.02	5	0.01
<i>Aspidogaster laticollis</i>	1	0.01	1	0.01	-	-	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	2	0.01	7	0.05	10	0.04	-	-	6	0.03	24	0.03
<i>Aspidogaster laticollis</i>	1	0.01	-	-	1	0.00	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	1	0.01	-	-	1	0.00	-	-	-	-	1	0.01
<i>Aspidogaster laticollis</i>	-	-	1	0.01	-	-	-	-	2	0.01	3	0.01
<i>Aspidogaster laticollis</i>	60	0.6	476	3.12	69	0.31	145	0.74	119	2.25	1199	1.32
<i>Aspidogaster laticollis</i>	1	0.01	3	0.02	-	-	-	-	-	-	4	0.00
<i>Aspidogaster laticollis</i>	-	-	-	-	-	-	5	0.03	1	0.01	6	0.01
<i>Aspidogaster laticollis</i>	-	-	-	-	1	0.00	-	-	-	-	1	0.00
<i>Aspidogaster laticollis</i>	5	0.04	7	0.05	-	-	-	-	5	0.03	17	0.02
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	2	0.01
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	1	0.01	1	0.01
<i>Aspidogaster laticollis</i>	-	-	-	-	-	-	-	-	-	-	2	0.00
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	51	0.06
<i>Aspidogaster laticollis</i>	41	0.50	4	0.03	1	0.01	3	0.02	2	0.02	3	0.00
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	3	0.00
<i>Aspidogaster laticollis</i>	3	0.03	-	-	-	-	-	-	-	-	3	0.00
<i>Aspidogaster laticollis</i>	3	0.03	5	0.04	6	0.03	3	0.02	1	0.01	70	0.08
<i>Aspidogaster laticollis</i>	35	0.44	-	-	-	-	-	-	-	-	1	0.00
<i>Aspidogaster laticollis</i>	4	0.01	-	-	-	-	-	-	-	-	18	0.02
<i>Aspidogaster laticollis</i>	9	0.07	6	0.04	3	0.01	-	-	-	-	104	0.03
<i>Aspidogaster laticollis</i>	1028	3.15	204	20.07	213	0.96	896	4.58	1044	19.50	1044	10.03
<i>Aspidogaster laticollis</i>	13	0.02	-	-	2	0.01	1	0.01	-	-	16	0.02
<i>Aspidogaster laticollis</i>	232	1.43	14	0.09	126	0.57	145	0.74	111	0.56	618	0.68
<i>Aspidogaster laticollis</i>	1	0.01	-	-	1	0.00	-	-	-	-	2	0.00
<i>Aspidogaster laticollis</i>	11	0.04	1	0.01	-	-	-	-	-	-	12	0.01
<i>Aspidogaster laticollis</i>	2	0.01	-	-	-	-	-	-	-	-	2	0.00
<i>Aspidogaster laticollis</i>	2	0.01	-	-	1	0.00	-	-	-	-	2	0.00
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.00
<i>Aspidogaster laticollis</i>	1	0.01	-	-	-	-	-	-	-	-	1	0.00

Table 1.2. continued.

	TM A11-111	TM A11-11A	TM A11-11B	TM A11-11C	TM A11-11D	TM A11-11E	Total
<i>Leptocryptus</i> sp.	-	-	1	1	-	-	2
<i>Leptocryptus</i> sp.	46	0.34	2	0.05	14	0.48	64
<i>Leptocryptus</i> sp.	-	-	-	-	-	-	-
<i>Leptocryptus</i> sp.	78	0.57	21	0.14	44	0.24	143
<i>Leptocryptus</i> sp.	-	-	-	-	-	-	-
<i>Leptocryptus</i> sp.	-	-	-	-	-	-	-
<i>Leptocryptus</i> sp.	18	0.13	4	0.03	17	0.12	57
<i>Leptocryptus</i> sp.	210	1.54	-	-	2	0.01	212
<i>Leptocryptus</i> sp.	5	0.04	-	-	-	-	5
<i>Leptocryptus</i> sp.	-	-	-	-	-	-	-
<i>Leptocryptus</i> sp.	4	0.03	-	-	1	0.01	5
<i>Leptocryptus</i> sp.	14	0.1	-	-	1	0.01	15
<i>Leptocryptus</i> sp.	2	0.02	4	0.03	-	-	6
<i>Leptocryptus</i> sp.	1	0.01	-	-	-	-	1
<i>Leptocryptus</i> sp.	1	0.01	-	-	-	-	1
<i>Leptocryptus</i> sp.	1	0.01	-	-	-	-	1
<i>Leptocryptus</i> sp.	4	0.03	3	0.02	5	0.03	12
<i>Leptocryptus</i> sp.	9	0.07	-	-	-	-	9
<i>Leptocryptus</i> sp.	14	0.1	-	-	-	-	14
<i>Leptocryptus</i> sp.	604	4.48	109	0.77	2	0.01	715
<i>Leptocryptus</i> sp.	117	0.88	144	0.94	145	1.07	410
<i>Leptocryptus</i> sp.	1	0.01	-	-	-	-	1
<i>Leptocryptus</i> sp.	-	-	-	-	-	-	-
<i>Leptocryptus</i> sp.	-	-	-	-	-	-	-
<i>Leptocryptus</i> sp.	-	-	-	-	-	-	-
<i>Leptocryptus</i> sp.	1	0.01	-	-	-	-	1
<i>Leptocryptus</i> sp.	119	0.89	24	0.17	101	0.74	244
<i>Leptocryptus</i> sp.	154	1.15	25	0.18	118	0.87	317
Total	1314	9.72	244	1.72	245	1.78	1840

Table 1.3.17

Estimates of density (number m⁻²) of the dominant taxa (all of the listed species) at the nearshore station, April through October 1976. Asterisks indicate species not present.

TM A11-11A	TM A11-11B	TM A11-11C	TM A11-11D	TM A11-11E	TM A11-11F	TM A11-11G
<i>Leptocryptus</i>	14	4.12	43	-	-	-
<i>Leptocryptus</i>	61	47	474	274	247	-
<i>Leptocryptus</i>	503	244.7	1207	2414	2425	1744
<i>Leptocryptus</i>	2	33	24	24	24	24
<i>Leptocryptus</i>	-	40	54	24	24	24
<i>Leptocryptus</i>	2	21	14	24	24	24
<i>Leptocryptus</i>	5	35	858	24	24	24
<i>Leptocryptus</i>	144	33	45	54	73	133
<i>Leptocryptus</i>	33	54	4	-	-	-
<i>Leptocryptus</i>	7	-	144	45	-	-
<i>Leptocryptus</i>	130	144.5	2736	1441	1412	7330
<i>Leptocryptus</i>	44	-	-	-	-	73
<i>Leptocryptus</i>	14	7	2	-	-	111
<i>Leptocryptus</i>	-	111	143	144	142	244
<i>Leptocryptus</i>	-	333	2736	1444	1444	744
<i>Leptocryptus</i>	2	5	2	7	5	14
<i>Leptocryptus</i>	-	-	14	-	-	-
<i>Leptocryptus</i>	491	4032	8632	4410	5744	7444
<i>Leptocryptus</i>	12	40	52	52	-	-
<i>Leptocryptus</i>	-	144	121	38	17	134
<i>Leptocryptus</i>	-	14	40	17	24	2
<i>Leptocryptus</i>	-	144	245	14	14	-
<i>Leptocryptus</i>	2	7	24	40	40	240
<i>Leptocryptus</i>	-	35	24	-	-	-
<i>Leptocryptus</i>	33	44	819	83	-	-
<i>Leptocryptus</i>	437	1044	7744	7722	3483	7444
<i>Leptocryptus</i>	43	142	43	344	24	4042
<i>Leptocryptus</i>	7	24	74	14	114	94
<i>Leptocryptus</i>	-	24	43	44	144	113
<i>Leptocryptus</i>	2	404	1373	222	81	2
<i>Leptocryptus</i>	-	14	14	61	92	12
<i>Leptocryptus</i>	-	35	24	-	-	243
<i>Leptocryptus</i>	2	-	-	35	-	-
<i>Leptocryptus</i>	492	1447	7722	5000	4444	7441
<i>Leptocryptus</i>	47	24	-	73	24	204
<i>Leptocryptus</i>	-	43	64	2	5	2
<i>Leptocryptus</i>	2	52	444	144	242	24
<i>Leptocryptus</i>	-	244	444	444	444	244
<i>Leptocryptus</i>	4	2	17	73	115	14

Table 3.2-38

Estimates of biomass (mg/m²) of the dominant taxa by weight at the macroinvertebrate stations, April through October 1976. Dashes indicate species not present; * indicates organism present but weight not determined.

	Apr	May	Jun	Jul	Aug	Sep	Oct
TM-AQI-1A1							
<u>Limnodynastes bilineatus</u>	107.0	461.5	720.2	866.1	687.4	378.3	614.4
<u>Chironomus tentans</u>	*	*	214.1	17.5	*	145.4	284.4
<u>Gammarus viridulus</u>	520.0	143.9	1101.1	1144.4	1484.9	1144.9	154.2
TM-AQI-1A2							
<u>Limnodynastes bilineatus</u>	111.1	248.4	544.2	545.1	626.2	112.8	141.2
<u>Chironomus tentans</u>	*	141.4	1177.1	242.2	605.4	144.4	17.3
<u>Gammarus viridulus</u>	846.8	1635.1	1412.8	144.4	1704.1	1744.4	114.4
TM-AQI-11A1							
<u>Limnodynastes bilineatus</u>	233.7	864.4	1444.9	419.3	1132.6	1125.7	1447.7
<u>Chironomus tentans</u>	*	54.2	44.4	1.9	*	1.0	*
<u>Gammarus viridulus</u>	1126.4	895.8	3401.7	1965.0	4714.1	1442.4	4243.6
TM-AQI-11A2							
<u>Limnodynastes bilineatus</u>	240.3	1170.4	2104.3	1414.4	444.7	1025.0	1023.2
<u>Chironomus tentans</u>	*	258.0	444.2	78.8	4.5	1.0	*
<u>Gammarus viridulus</u>	746.2	767.2	2671.3	1730.0	4273.6	4714.4	5724.4
TM-AQI-981							
<u>Limnodynastes bilineatus</u>	512.1	857.0	1540.4	1314.4	421.4	1404.1	2075.9
<u>Chironomus tentans</u>	*	1275.3	3422.4	601.7	145.3	14.0	60.5
<u>Gammarus viridulus</u>	161.4	210.1	454.3	1517.3	1444.4	1744.6	2124.5

Table 3.2-39

Diversity (D) values at the macroinvertebrate stations, April through October 1976.

Station Number	Apr	May	Jun	Jul	Aug	Sep	Oct
TM-AQI-1A1	2.75	2.24	2.44	1.83	2.06	2.40	2.64
TM-AQI-1A2	2.46	2.74	1.76	1.37	1.53	0.42	1.32
TM-AQI-11A1	1.78	0.69	1.44	1.20	1.31	0.79	0.73
TM-AQI-11A2	1.76	1.11	1.37	0.93	1.34	1.30	0.44
TM-AQI-981	0.66	1.48	1.48	1.13	1.66	1.08	0.88

Table 3.2-40

Indices of percent similarity of species composition (PSc) between the macroinvertebrate stations sampled April through October 1974, 1975, and 1976. Prefix TM-AQI- deleted for table.

1974		85	1A2	68			1975
	79	77	11A1	64	87		
	84	88	83	11A2	65	93	93
87	74	86	83	981	64	74	85
11A2	11A1	1A2	1A1		1A1	1A2	11A1
77	76	91	61	981			
	89	75	60	11A2			
		70	57	11A1			
1976			64	1A2			

Table 3.2-41

Analysis of variance, randomized block design for number of
taxa, April through October 1976.

		SS	df	MS	F
Total	52	2711.48800	51		
Taxa	4	614.41010	153.70710	10.7414*	
Date	13	855.87100	48.14392	4.7432*	
Error	52	740.20700	14.2425		

* Denotes $\alpha = 0.05$.

Table 3.2-42

Comparison of mean number of taxa at the Macmillanstrata
station using Student-Newman-Keuls multiple range test, April
through October 1976. Mean number of taxa underscored
are not significantly different ($\alpha = 0.05$).

Taxa	1A2	1A1	1A2	1A1	1A1
Mean No. of taxa	14	17	19	20	23

Table 3.2-43

Comparison of mean number of taxa by sampling date (April through October 1976) using Student-Newman-Keuls
multiple range test.

	1A2	1A1	1A2	1A1	1A2	1A1	1A2	1A1	1A2	1A1	1A2	1A1	1A2	1A1	1A2	1A1	1A2	1A1
6 Apr	-*	-	-**	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
10 Apr	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1 May	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
18 May	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1 Jun	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
15 Jun	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
6 Jul	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
20 Jul	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
3 Aug	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
17 Aug	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
7 Sep	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
21 Sep	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
5 Oct	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

* Dash indicates not significant.
** Significant at 0.05.

Table 3.2-44

150

Three-factor analysis of variance for densities of *Limnodrilus hoffmeisteri* at the macroinvertebrate stations, May through October 1974 through 1976.

Source	df	SS	MS	F
Total	719	262.70432		
Year	2	24.55151	12.27576	14.24*
Date	11	13.44256	1.22205	14.34*
Station	4	63.52646	15.88161	179.44*
Year x Date	22	31.77747	1.44443	16.74*
Year x Station	8	26.16405	3.27051	37.12*
Date x Station	44	17.37276	0.39484	4.51*
Year x Date x Station	88	44.91197	0.50923	5.72*
Error	540	40.91209	0.07574	

* Significant at 0.05.

Table 3.2-45

Comparison of logarithmic mean density of *Limnodrilus hoffmeisteri* by year (May through October 1974-1976) using Student-Newman-Keuls multirange test ($\alpha = 0.05$).

Year	1974	1975	1976
Logarithmic Mean Density	3.12415	3.35345	3.57244

Table 3.2-46

Comparison of logarithmic mean density of *Limnodrilus hoffmeisteri* by sampling date (May through October 1974-1976) using Student-Newman-Keuls multirange test.

	I May	I Sep	II Sep	I Sep	II Aug	I Aug	II Jul	I Jul	II Jun	I Jun	II May
I May	***	-	***	x	x	-	x	x	x	x	-
II May	x	x	x	x	-	-	x	x	x	x	
I Jun	x	x	-	-	x	x	-	-	-	-	
II Jun	x	x	-	-	-	x	-	-	-	-	
I Jul	x	x	-	-	-	x	-	-	-	-	
II Jul	x	x	-	-	x	x	-	-	-	-	
I Aug	x	x	x	x	-	-	-	-	-	-	
II Aug	x	x	-	-	-	-	-	-	-	-	
I Sep	x	x	-	-	-	-	-	-	-	-	
II Sep	x	x	-	-	-	-	-	-	-	-	
I Oct	-	-	-	-	-	-	-	-	-	-	

* Signifies first (I) or second (II) sampling date.

** Dash indicates not significant.

*** Significant at 0.05.

Table 3.2-47

Comparison of logarithmic mean density of *Limnodrilus hoffmeisteri* at the macroinvertebrate stations using Student-Newman-Keuls multirange test, May through October 1974-1976. Mean densities underscored are not significantly different ($\alpha = 0.05$).

Station	1A1	1A2	9B1	11A2	11A1
Logarithmic Mean Density	2.60672	3.36002	3.42609	<u>3.59059</u>	<u>3.61244</u>

POOR ORIGINAL

Table 3.3-1

Summary of data collected at the sediment-water interface for 1975, 1976, and 1978.

	1975				1976				1978				Total 1975-1978
	10/25	10/26	10/27	10/28	10/29	10/30	10/31	11/1	11/2	11/3	11/4	11/5	
% of rain collected	23	55	82	75	58	75	3	50	16	92	34	18	71
% of streamflow collected	106	100	1018	216	1275	1572	800	1398	2270	843	2335	1690	1583
% of streamflow	1.08	1.05	5.97	2.00	5.06	51.0	2.92	11.4	25.0	2.80	5.00	2.70	6.4
Acoustic index (m)	1.50	2.00	1.00	1.00	1.00	1.00	0.08	1.00	1.00	1.00	1.00	1.00	1.00
Flow rate (m ³ /sec)	6.50	85.50	47.40	65.00	76.00	65.00	84.27	86.16	70.16	82.07	82.07	78.28	74.08
Channel width (m)	11.95	1.73	7.55	16.45	11.00	20.97	5.95	6.10	18.00	10.94	7.94	18.16	10.50
Flow rate (m ³ /sec)	0.54	116.9	500.1	565.3	600.2	600.2	1006.9	2262.0	1047.9	1008.5	1007.2	805.8	105.6
Channel width (m)	4.00	5.2	95.2	112.5	217.3	106.9	42.0	18.7	144.3	100.2	117.6	200.1	20.4
Channel width (m)													

1565 200




Figure 3.1-1. Location of benthic macroinvertebrate stations in the vicinity of Toms.





Figure 3.2-1. Percent composition of macroinvertebrate taxa, based on the total number of individuals collected at the macroinvertebrate stations, April through October 1970.



POOR ORIGINAL

155




Figure 3.3-2. Diversity (D) values at macroinvertebrate stations April through October 1974 (---), 1975 (—), and 1976 (-.-.-). "Status" refers to water quality.

1565 204

4.0 AMBIENT WATER QUALITY

The ETS, Appendix B, Section 4.2.2 requires that certain water quality parameters be analyzed semimonthly from April through October in conjunction with macroinvertebrate studies. Results of the 1976 program are given below.

4.1 METHODS

Surface water samples for chemical analyses were collected semimonthly April through October 1976 at each macroinvertebrate station mentioned in Section 3.0 (Table 3.1-1, Figure 3.1-1). Samples were preserved in nalgene bottles and sterilized ground glass bottles (fecal coliform) in the field and transported, in an insulated plastic cooler at approximately 4 C, to the laboratory for analysis. Water quality parameters and methods of determination are given in Table 4.1-1. Dissolved oxygen determinations were made in the field with a YSI Model 54 Oxygen Analyzer.

Geometric means of fecal coliform densities were calculated for all possible combinations of five consecutive samples at each station. These were compared to the limit (200 fecal coliform colonies per 100 ml) established in the "Water Quality Criteria" for the Commonwealth of Pennsylvania (Pennsylvania Department of Environmental Resources 1971).

4.2 RESULTS

4.2.1 PHYSICOCHEMICAL PARAMETERS

Except for dissolved oxygen and total suspended solids, the pattern of monthly distribution of the other parameters was similar (Table 4.2-1).

Mean values (except dissolved oxygen and suspended solids) were high in September and low in April and May. The mean concentrations of dissolved oxygen were high in October and April and low in July. Suspended solids were high in August and low April through June.

Differences in the concentrations of each parameter between the stations were examined using the data in Tables 4.2-2 through 4.2-8. The monthly mean concentrations of most parameters at Stations 1A1, 1A2, 11A2, and 9B1 generally differed from those at Station 11A1 (TMINS Discharge). Mean values for conductivity, dissolved oxygen, alkalinity, nitrate nitrogen, total phosphates, soluble orthophosphates, sulfates, chlorides, and suspended solids were generally higher at Station 11A1. Differences were generally small. Mean values for other parameters at 11A1 were similar to those found at the other stations.

Mean concentrations of the parameters at Station 1A2 (upstream from the Discharge) and Station 11A2 (downstream from the Discharge) were similar. Any changes in water quality appeared to be limited to the immediate area of the Discharge. Similar conditions were observed in 1974 and 1975 (Potter and Associates 1975, 1976).

Inspection of the mean values of the water quality parameters for 1974 through 1976 indicated that no detectable differences resulted from the operation of TMINS, Unit 1. The requirements of the ETS, Appendix B, Section 4.2.2 have been fulfilled; monitoring will continue as prescribed in the ETS for Unit 2.

4.2.2 FECAL COLIFORM

Mean monthly fecal coliform densities were lowest in April (213 colonies per 100 ml); they increased throughout the summer and peaked (10,768 colonies per 100 ml) in September (Table 4.2-1). Densities declined during October. Highest overall mean density was found at Station 1A1 (upstream from the Discharge), which had the highest density on 8 of the 14 sample dates (Tables 4.2-2 through 4.2-8). The lowest overall mean density was found at the Discharge (Station 11A1), which had the lowest density on 5 of the 14 sample dates. No consistent trends among Stations 1A2, 11A2, and 9B1 were noted.

The lowest geometric mean for sets of five consecutive samples at all stations was recorded for the period 6 April through 2 June (Table 4.2-9). The highest geometric mean was recorded at all stations for the periods 20 July through 21 September and 3 August through 5 October. The lowest overall geometric mean occurred at Station 11A1 for the period 6 April through 2 June.

For all stations the geometric mean of fecal coliform densities exceeded the limit established for the Commonwealth of Pennsylvania.

Fecal coliform densities for 1974 through 1976 revealed similar trends. The densities were generally higher at Station 1A1 and lower at Station 11A1. There appeared to be no detectible effect on the growth of bacteria as a result of the operation of TMINS, Unit 1. The ETS requirements for fecal coliform have been completed. Additional collections will be as prescribed in the ETS for Unit 2.

1565 207

Anonymous. 1975. Standard methods for the examination of water and waste water. 14th edition. American Public Health Association, Inc., New York, New York. 1193 pp.

Anonymous. 1974. Methods for chemical analysis of water and wastes. Environmental Protection Agency, Cincinnati, Ohio. 312 pp.

Orion, Research Inc. 1967. Instruction manual halide ions. Cambridge, Massachusetts. 20 pp.

Pennsylvania Department of Environmental Resources. 1971. Water quality criteria, Chapter 93. Rules and regulations, Title 25. Article II, Water resources. 98 pp.

Potter, W.A. and Associates. 1975. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Annual Report for 1974. Ichthyological Associates, Inc. 468 pp.

Potter, W.A. and Associates. 1976. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Annual Report for 1975. Ichthyological Associates, Inc. 395 pp.

POOR ORIGINAL

1565 208

POOR ORIGINAL

Table 4.1-1

Parameter, method used, and reference for analysis of water taken in the vicinity of THINS in 1976.

Parameter	Method	Reference
pH	Glass electrode	APHA (1975)
Conductivity	Line operated conductivity meter @ 20 C	APHA (1975)
Dissolved Oxygen	YSI Model 54-Oxygen Analyzer	
Total alkalinity	Potentiometric titration	APHA (1975)
Total phosphate	Persulfate digestion; single reagent	EPA (1974)
Orthophosphate	Single reagent	EPA (1974)
Ammonia nitrogen	Specific ion electrode	EPA (1974)
Nitrate nitrogen	U.V. spectrophotometric	APHA (1975)
Nitrite nitrogen	Colorimetric	APHA (1975)
Sulfate	Turbidmetric	EPA (1974)
Chloride	Specific ion electrode	Orion Inc. (1967)
Suspended solids	Non-filterable residue	APHA (1975)
Fecal coliform	Membrane filter (MF)	APHA (1975) PA DFR (1971)

1565 209

POOR ORIGINAL

Table 4.2-1

Monthly mean, minimum, and maximum concentrations of selected water quality parameters taken in the vicinity of DMNS in 1976. Values are expressed in ppm except for conductivity, pH, and fecal coliform.

Month		Jan	Feb	Mar	Apr	May	Jun	Jul
Parameter								
Conductivity (umhos/cm @ 25°C)	Mean	206	203	256	254	246	324	333
	Min.	145	186	209	208	186	244	179
	Max.	279	230	416	346	329	446	323
pH	Mean	-	-	-	-	-	-	-
	Min.	7.24	7.44	7.66	7.64	7.50	7.66	7.26
	Max.	8.06	7.98	8.14	7.94	8.17	8.28	7.51
Dissolved Oxygen	Mean	9.1	8.4	8.4	7.4	8.1	8.3	10.1
	Min.	7.7	6.8	7.0	6.9	7.7	7.3	8.1
	Max.	10.3	10.0	9.8	8.2	8.9	9.8	12.0
Total Alkalinity	Mean	34.9	40.8	49.4	45.7	57.7	76.1	32.8
	Min.	22.4	33.2	36.4	39.2	46.4	54.8	28.8
	Max.	52.8	52.0	69.6	55.0	76.2	100.2	38.4
Ammonia nitrogen	Mean	0.07	0.08	0.06	0.06	0.05	0.12	0.12
	Min.	0.05	0.03	0.05	0.05	0.04	0.08	0.07
	Max.	0.12	0.09	0.09	0.07	0.06	0.18	0.15
Nitrate nitrogen	Mean	0.89	0.82	1.02	1.35	1.95	3.02	1.80
	Min.	0.59	0.52	0.48	0.94	0.18	0.41	0.90
	Max.	1.26	1.25	2.19	2.07	4.49	6.14	2.75
Nitrite nitrogen	Mean	0.02	0.03	0.02	0.02	0.03	0.03	0.02
	Min.	0.01	0.01	0.01	0.01	0.01	0.02	0.02
	Max.	0.03	0.06	0.03	0.02	0.02	0.04	0.02
Phosphate, Total	Mean	0.33	0.38	0.42	0.43	0.66	0.67	0.48
	Min.	0.24	0.28	0.26	0.23	0.43	0.52	0.32
	Max.	0.40	0.53	0.64	0.90	0.97	0.85	0.78
Phosphate, Soluble ortho	Mean	0.08	0.11	0.04	0.10	0.17	0.20	0.19
	Min.	0.03	0.03	0.01	0.04	0.01	0.02	0.04
	Max.	0.15	0.23	0.13	0.22	0.28	0.32	0.23
Sulfate	Mean	54.2	50.5	70.2	68.5	55.2	79.5	64.9
	Min.	36.1	36.9	54.1	55.6	24.3	54.6	45.0
	Max.	82.2	64.0	129.8	113.6	96.4	116.6	102.6
Chloride	Mean	11.5	9.7	14.2	12.4	12.6	17.2	10.3
	Min.	8.2	8.9	10.6	11.3	7.4	14.2	9.1
	Max.	13.5	10.5	23.0	17.4	18.1	22.7	13.8
Suspended Solids (Non-filterable)	Mean	20.6	19.3	20.6	38.7	53.4	30.5	34.9
	Min.	5.2	12.4	13.8	15.8	20.8	20.0	24.1
	Max.	49.4	25.2	28.8	142.0	92.8	45.2	57.2
Fecal Coliform (Colonies/100 ml)	Mean	223	1146	3038	3197	4771	10768	2404
	Min.	82	396	1417	1600	1367	2333	200
	Max.	170	2714	4500	7600	17000	44000	5224

1565 210

Table 4.2-2

Summary of selected physicochemical parameters taken on 6 and 20 April 1976 in the vicinity of THINS. Values are expressed in ppm except for conductivity, pH, and total coliform.

Location	Date	Conductivity ($\mu\text{mhos/cm}$) at 20°C	pH	Dissolved Oxygen	Total Alkalinity as CaCO_3	$\text{NH}_3\text{-N}$	$\text{Mg}^{2+}\text{-N}$	$\text{Mn}^{2+}\text{-N}$	Pb (Total)	Pb (ortho)	SO_4	Cl	Suspended Solids ($\mu\text{mhos/cm}$) (filterable)	Total Coliform (colonies/ 100 ml)
TH-AQ1-1A1	6 Apr	149	7.24	9.8	22.4	0.06	0.09	0.01	0.40	0.03	37.9	8.2	54.2	113
TH-AQ1-1A2		145	7.27	9.7	23.6	0.05	0.72	0.01	0.16	0.03	36.1	8.5	41.2	82
TH-AQ1-11A1		145	7.40	10.3	24.4	0.08	0.75	0.02	0.16	0.03	36.1	8.5	47.4	85
TH-AQ1-11A2		145	7.34	9.9	24.0	0.06	0.74	0.01	0.32	0.03	37.0	8.2	35.6	83
TH-AQ1-9B1		145	7.32	9.7	24.8	0.08	0.75	0.01	0.22	0.03	36.1	8.5	24.4	84
TH-AQ1-1A1	20 Apr	270	8.06	8.4	35.2	0.06	0.59	0.01	0.24	0.08	82.2	12.0	5.2	135
TH-AQ1-1A2		264	7.98	8.1	31.0	0.07	1.26	0.01	0.15	0.15	68.6	12.8	15.2	570
TH-AQ1-11A1		268	8.00	7.9	31.6	0.04	0.82	0.02	0.28	0.10	76.8	13.5	7.6	400
TH-AQ1-11A2		262	8.02	8.6	50.0	0.10	1.16	0.01	0.36	0.15	68.0	12.8	13.2	350
TH-AQ1-9B1		265	7.98	8.2	51.5	0.12	1.25	0.01	0.34	0.15	63.7	12.4	7.8	240
TH-AQ1-1A1	Apr	209	-	9.1	28.8	0.06	0.70	0.01	0.32	0.05	60.0	10.1	19.7	124
TH-AQ1-1A2		202	-	8.9	37.3	0.06	0.94	0.02	0.16	0.09	52.4	10.6	26.2	326
TH-AQ1-11A1		207	-	9.1	33.0	0.06	0.79	0.02	0.32	0.06	56.4	11.0	28.5	253
TH-AQ1-11A2		204	-	9.3	37.6	0.08	0.95	0.02	0.34	0.09	52.5	10.5	24.4	212
TH-AQ1-9B1		205	-	9.0	34.3	0.10	1.00	0.01	0.11	0.19	49.9	10.5	17.1	162

MEAN VALUES FOR APRIL 1976

Table 4.2-3

Summary of selected physicochemical parameters taken on 4 and 18 May 1976 in the vicinity of THINS. Values are expressed in ppm except for conductivity, pH, and total coliform.

Location	Date	Conductivity ($\mu\text{mhos/cm}$) at 20°C	pH	Dissolved Oxygen	Total Alkalinity as CaCO_3	$\text{NH}_3\text{-N}$	$\text{Mg}^{2+}\text{-N}$	$\text{Mn}^{2+}\text{-N}$	Pb (Total)	Pb (ortho)	SO_4	Cl	Suspended Solids ($\mu\text{mhos/cm}$) (filterable)	Total Coliform (colonies/ 100 ml)
TH-AQ1-1A1	4 May	190	7.44	9.0	31.6	0.09	0.76	0.01	0.36	0.14	47.5	10.1	21.6	626
TH-AQ1-1A2		186	7.54	8.9	33.4	0.07	0.84	0.01	0.33	0.08	58.7	10.5	21.6	946
TH-AQ1-11A1		216	7.66	10.0	33.4	0.09	0.78	0.02	0.28	0.16	62.2	10.5	15.8	410
TH-AQ1-11A2		172	7.52	8.9	33.2	0.09	0.76	0.02	0.32	0.07	52.3	10.3	21.0	640
TH-AQ1-9B1		190	7.50	8.6	33.2	0.09	0.78	0.02	0.29	0.07	48.7	10.1	12.4	500
TH-AQ1-1A1	18 May	232	7.98	8.0	43.4	0.03	0.52	0.02	0.14	0.03	64.0	9.2	16.0	274
TH-AQ1-1A2		204	7.52	7.0	27.8	0.08	0.64	0.04	0.47	0.16	44.0	9.9	21.2	1200
TH-AQ1-11A1		201	7.86	8.2	52.0	0.08	1.25	0.06	0.53	0.23	36.9	9.4	25.2	1950
TH-AQ1-11A2		209	7.56	7.0	50.8	0.07	0.86	0.04	0.55	0.11	50.6	8.9	14.2	1247
TH-AQ1-9B1		214	7.58	6.8	50.8	0.14	1.07	0.05	0.59	0.19	40.6	9.4	20.4	1425
TH-AQ1-1A1	May	211	-	8.5	38.5	0.06	0.64	0.02	0.35	0.06	55.8	9.7	18.9	1480
TH-AQ1-1A2		195	-	8.0	40.6	0.18	0.72	0.03	0.40	0.12	46.4	9.7	21.4	1440
TH-AQ1-11A1		208	-	9.1	42.7	0.08	1.02	0.04	0.50	0.18	28.5	9.8	23.4	710
TH-AQ1-11A2		200	-	8.6	40.0	0.08	0.81	0.03	0.48	0.10	51.4	9.4	23.1	1204
TH-AQ1-9B1		202	-	7.7	34.0	0.08	0.72	0.04	0.44	0.11	39.6	9.4	16.1	1478

MEAN VALUES FOR MAY 1976

Table 4.2-4

Summary of selected physicochemical parameters taken on 1 and 15 June 1976 in the vicinity of 28°N. Values are expressed in ppm except for conductivity, pH, and fecal coliform.

Location	Date	Conductivity (micromhos/cm at 20°C)	pH	Dissolved oxygen	Total Alkalinity as CaCO ₃	SiO ₂ -N	NO ₃ -N	NO ₂ -N	PO ₄ -P (Total)	PO ₄ -P (Ortho)	% Cl	% suspended solids	Total coliform
1 Jun													
TS-AQ1-1A1	216	7.70	7.70	7.2	36.4	0.04	0.76	0.01	0.28	0.03	59.1	1.8	100
TS-AQ1-1A2	215	7.68	7.68	9.2	36.5	0.07	0.78	0.01	0.30	0.03	59.1	1.8	100
TS-AQ1-11A1	212	7.68	7.68	9.8	36.4	0.06	0.81	0.01	0.29	0.03	59.1	1.8	100
TS-AQ1-11A2	214	7.67	7.67	9.2	36.4	0.06	0.78	0.01	0.28	0.03	59.1	1.8	100
TS-AQ1-9B1	209	7.66	7.66	9.0	36.2	0.06	0.80	0.01	0.28	0.03	59.1	1.8	100
15 Jun													
TS-AQ1-1A1	256	8.15	8.15	7.9	43.4	0.06	0.48	0.02	0.41	0.03	59.1	1.8	100
TS-AQ1-1A2	276	7.98	7.98	7.2	65.4	0.06	1.22	0.02	0.56	0.03	59.1	1.8	100
TS-AQ1-11A1	416	8.13	8.13	7.7	69.6	0.06	2.19	0.03	0.64	0.13	129.8	2.0	117
TS-AQ1-11A2	275	8.03	8.03	7.0	66.8	0.05	1.16	0.02	0.58	0.03	68.5	1.8	100
TS-AQ1-9B1	283	7.93	7.93	8.0	64.4	0.05	1.18	0.02	0.54	0.02	68.5	1.8	100
MEAN VALUES FOR 1976													
TS-AQ1-1A1	231	-	-	8.6	41.4	0.08	0.62	0.02	0.45	0.02	67.1	1.7	100
TS-AQ1-1A2	246	-	-	8.2	50.9	0.06	1.00	0.02	0.43	0.03	67.0	1.8	100
TS-AQ1-11A1	314	-	-	8.6	53.0	0.06	1.50	0.02	0.47	0.08	94.5	1.8	100
TS-AQ1-11A2	254	-	-	8.1	51.6	0.06	0.97	0.02	0.43	0.03	63.7	1.8	100
TS-AQ1-9B1	246	-	-	8.5	50.3	0.06	0.94	0.02	0.40	0.03	63.7	1.8	100

Table 4.2-5

Summary of selected physicochemical parameters taken on 6 and 20 July 1976 in the vicinity of 28°N. Values are expressed in ppm except for conductivity, pH, and fecal coliform.

Location	Date	Conductivity (micromhos/cm at 20°C)	pH	Dissolved oxygen	Total Alkalinity as CaCO ₃	SiO ₂ -N	NO ₃ -N	NO ₂ -N	PO ₄ -P (Total)	PO ₄ -P (Ortho)	% Cl	% suspended solids	Total coliform
6 Jul													
TS-AQ1-1A1	211	7.78	7.78	NA	39.4	0.06	1.04	0.01	0.23	0.04	59.9	11.1	100
TS-AQ1-1A2	216	7.78	7.78	NA	41.6	0.06	1.02	0.01	0.24	0.04	62.2	11.1	100
TS-AQ1-11A1	340	7.94	7.94	NA	42.8	0.07	1.77	0.01	0.41	0.08	111.6	17.0	1967
TS-AQ1-11A2	216	7.75	7.75	NA	41.2	0.06	1.06	0.01	0.26	0.04	62.2	11.9	100
TS-AQ1-9B1	216	7.74	7.74	NA	40.8	0.07	1.04	0.01	0.26	0.04	59.9	11.3	100
20 Jul													
TS-AQ1-1A1	258	7.64	7.64	7.6	34.2	0.07	0.94	0.01	0.31	0.10	67.4	12.8	100
TS-AQ1-1A2	212	7.69	7.69	7.3	67.2	0.05	1.35	0.02	0.54	0.15	61.6	13.3	100
TS-AQ1-11A1	212	7.91	7.91	8.2	55.0	0.05	2.02	0.02	0.90	0.22	78.2	17.4	4000
TS-AQ1-11A2	259	7.71	7.71	7.3	51.4	0.06	1.67	0.02	0.62	0.18	64.5	16.4	4450
TS-AQ1-9B1	243	7.66	7.66	6.9	48.4	0.06	1.54	0.02	0.56	0.16	55.6	13.3	4450
MEAN VALUES FOR JULY 1976													
TS-AQ1-1A1	230	-	-	7.4	39.3	0.06	0.99	0.01	0.28	0.07	63.6	12.0	1783
TS-AQ1-1A2	235	-	-	7.3	44.4	0.06	1.18	0.02	0.34	0.09	61.9	12.3	2953
TS-AQ1-11A1	316	-	-	8.2	48.9	0.06	1.92	0.02	0.66	0.15	95.8	17.2	3250
TS-AQ1-11A2	264	-	-	7.3	46.3	0.06	1.37	0.02	0.54	0.11	61.4	13.1	3183
TS-AQ1-9B1	250	-	-	6.9	49.6	0.06	1.29	0.02	0.51	0.10	57.7	12.1	2787

NA = Not Available.

Table 4.2-6

Summary of selected physicochemical parameters taken on 3 and 17 August 1976 in the vicinity of TINS. Values are expressed in ppm except for conductivity, pH, and total alkalinity.

Location	Date	Conductivity (micromhos) at 20 C	pH	Dissolved Oxygen	Total Alkalinity as CaCO ₃	NO ₃ -N	NO ₂ -N	NO ₂ -N (Total)	PO ₄ (ortho)	SO ₄	Cl	Suspended Solids Can- filterable	Total coliform colonies/ 100 ml
TN-AQ1-1A1	3 Aug	308	8.17	7.7	52.2	0.05	0.18	0.01	0.43	96.4	16.0	29.8	1700
TN-AQ1-1A2		260	7.85	7.7	73.2	0.06	1.06	0.02	0.54	45.5	15.2	25.0	1390
TN-AQ1-11A1		329	8.06	8.7	76.2	0.05	1.95	0.02	0.24	74.4	18.1	45.6	1367
TN-AQ1-11A2		283	7.87	7.8	73.2	0.04	1.60	0.02	0.19	55.1	16.3	37.0	2480
TN-AQ1-901		275	7.88	8.4	68.0	0.05	0.99	0.02	0.50	74.4	15.6	29.2	1920
TN-AQ1-1A1	17 Aug	168	7.53	8.3	46.4	0.06	2.24	0.02	0.81	24.3	8.0	92.8	3800
TN-AQ1-1A2		166	7.50	7.9	48.6	0.04	2.32	0.02	0.71	24.3	7.4	69.6	2800
TN-AQ1-11A1		324	7.94	8.9	46.6	0.04	4.49	0.02	0.97	89.7	13.6	82.6	5380
TN-AQ1-11A2		166	7.54	7.8	46.4	0.04	2.22	0.02	0.71	11.4	7.6	69.2	5900
TN-AQ1-901		185	7.54	7.7	46.4	0.04	2.48	0.02	0.65	17.6	8.0	57.6	5750
MEAN VALUES FOR AUGUST 1976													
TN-AQ1-1A1	Aug	238	-	8.0	49.3	0.06	1.21	0.02	0.62	60.4	12.0	56.8	10400
TN-AQ1-1A2		213	-	7.8	60.9	0.05	1.69	0.02	0.62	34.4	11.3	46.8	2976
TN-AQ1-11A1		326	-	8.8	61.4	0.04	3.22	0.02	0.82	82.0	15.9	64.6	3334
TN-AQ1-11A2		225	-	7.8	59.8	0.04	1.91	0.02	0.65	43.2	12.0	53.1	4170
TN-AQ1-901		230	-	8.1	57.2	0.04	1.71	0.02	0.58	56.0	11.8	43.5	3835

Table 4.2-7

Summary of selected physicochemical parameters taken on 7 and 21 September 1976 in the vicinity of TINS. Values are expressed in ppm except for conductivity, pH, and total alkalinity.

Location	Date	Conductivity (micromhos) at 20 C	pH	Dissolved Oxygen	Total Alkalinity as CaCO ₃	NO ₃ -N	NO ₂ -N	NO ₂ -N (Total)	PO ₄ (ortho)	SO ₄	Cl	Suspended Solids Can- filterable	Total coliform colonies/ 100 ml
TN-AQ1-1A1	7 Sep	335	8.12	7.8	68.0	0.18	0.41	0.02	0.66	95.0	17.2	40.0	2700
TN-AQ1-1A2		336	8.25	9.0	99.2	0.12	3.97	0.05	0.74	58.4	12.2	30.7	2700
TN-AQ1-11A1		456	8.26	9.8	100.2	0.08	6.14	0.03	0.88	116.6	22.7	25.2	4200
TN-AQ1-11A2		375	8.24	9.0	99.4	0.11	4.40	0.03	0.78	86.3	18.4	33.2	4500
TN-AQ1-901		350	8.28	8.6	79.4	0.15	1.92	0.03	0.65	90.7	17.9	20.2	5821
TN-AQ1-1A1	21 Sep	306	7.68	7.3	54.8	0.14	1.10	0.02	0.52	94.2	16.7	20.4	1875
TN-AQ1-1A2		244	7.67	8.0	67.6	0.13	2.72	0.03	0.63	52.6	14.2	32.4	3074
TN-AQ1-11A1		318	7.90	8.4	65.6	0.09	3.90	0.03	0.75	82.0	17.7	37.2	2033
TN-AQ1-11A2		257	7.68	7.6	65.2	0.12	2.81	0.03	0.59	50.6	14.4	30.8	4025
TN-AQ1-901		286	7.66	7.9	64.4	0.12	2.79	0.03	0.60	62.2	14.7	29.4	4000
MEAN VALUES FOR SEPTEMBER 1976													
TN-AQ1-1A1	Sep	326	-	7.6	61.4	0.16	0.26	0.02	0.57	93.6	17.2	20.2	1037
TN-AQ1-1A2		290	-	8.5	81.4	0.12	3.34	0.04	0.68	44.5	15.7	34.4	3166
TN-AQ1-11A1		362	-	9.1	82.9	0.08	5.02	0.03	0.80	80.0	20.2	41.2	3099
TN-AQ1-11A2		316	-	8.3	80.3	0.12	1.61	0.03	0.60	71.4	16.4	32.4	3099
TN-AQ1-901		308	-	8.2	71.4	0.14	2.30	0.03	0.63	76.5	16.8	27.8	4025

Table 4.2-M

Record of selected physiological parameters taken on 5 and 19 October 1976. In the vicinity of 1917, values are expressed in the except for conductivity, pH, and Ca^{2+} (mM).

Run and Location	Date	Conductivity cmhos/cm at 25°C	pH	Dissolved Oxygen	Total Alkalinity meq/l	Na ⁺ %	Mg ²⁺ %	Ca ²⁺ %	Fe ³⁺ %	NO ₃ ⁻ %	SiO ₂ %	Unidentified %	Unidentified mg/l	Total Calcium mg/l
T7-A01-1A1	5 OCT	1.06	7.35	9.2	32.0	0.09	2.30	0.02	0.50	0.12	51.8	0.3	4.3	100.0
T7-A01-1A2	179	1.79	7.38	9.6	38.2	0.08	2.02	0.02	0.51	0.12	45.0	0.1	46.8	45.0
T7-A01-11A1	311	2.50	7.50	11.0	36.2	0.08	2.02	0.02	0.51	0.12	45.0	0.1	46.8	45.0
T7-A01-11A2	192	1.92	7.38	9.2	35.8	0.07	2.14	0.02	0.51	0.13	49.6	0.6	49.3	49.3
T7-A01-901	192	1.92	7.38	9.8	35.6	0.08	2.04	0.02	0.47	0.13	51.8	0.3	46.8	46.8
T7-A01-1A1	207	2.07	7.36	11.6	28.8	0.12	0.01	0.01	0.36	0.02	20.6	9.2	40.0	53.8
T7-A01-1A2	205	1.91	7.26	11.8	29.6	0.15	0.03	0.01	0.42	0.02	18.9	9.2	33.4	53.8
T7-A01-11A1	225	2.25	7.28	12.4	29.6	0.15	1.08	0.01	0.49	0.02	25.2	10.3	40.8	56.7
T7-A01-11A2	206	2.06	7.28	11.8	29.6	0.12	0.92	0.01	0.45	0.02	68.4	9.1	46.2	30.8
T7-A01-901	204	2.04	7.33	11.6	29.6	0.15	0.92	0.01	0.32	0.06	66.0	9.3	24.0	28.8
20-AN VALUES FOR OCTOBER 1976														
T7-A01-1A1	201	-	-	10.4	30.4	0.12	1.00	0.02	0.44	0.06	61.2	9.6	40.0	28.73
T7-A01-1A2	192	-	-	10.6	34.0	0.12	1.42	0.02	0.46	0.09	56.7	9.8	50.1	29.1
T7-A01-11A1	269	-	-	11.7	33.0	0.12	1.92	0.02	0.42	0.15	88.9	12.1	59.0	28.08
T7-A01-11A2	200	-	-	10.6	32.7	0.11	1.53	0.02	0.49	0.10	59.0	9.8	40.2	24.6
T7-A01-901	194	-	-	10.4	32.5	0.12	1.00	0.02	0.40	0.06	58.9	10.0	27.9	17.50

Control colony count.

Table 4.2-9

Combinations of five consecutive samples for determination of the geometric mean of fecal coliform density at ambient water quality stations, April through October 1976.

Station	1A1	1A2	11A1	11A2	9B2		1A1	1A2	11A1	11A2	9B1
6 Apr	113	82	85	82	84	15 Jun	4500	3150	1417	3450	2188
20 Apr	135	570	400	340	240	6 Jul	1600	1967	2050	1917	1975
4 May	646	396	410	640	530	20 Jul	1967	4000	4450	4450	7600
19 May	2714	1700	1050	1767	1625	3 Aug	17000	1390	1367	2480	1920
2 Jun	2364	3500	3050	3652	3100	17 Aug	3800	2800	5300	5900	5750
Geometric mean	576	643	537	651	557	Geometric mean	3911	2494	2479	3364	3250
20 Apr	135	570	400	340	240	6 Jul	1600	1967	2050	1917	1975
4 May	646	396	410	640	530	20 Jul	1967	4000	4450	4450	7600
19 May	2714	1700	1050	1767	1625	3 Aug	17000	1390	1367	2480	1920
2 Jun	2364	3500	3050	3652	3100	17 Aug	3800	2800	5300	5900	5750
15 Jun	4500	3150	1417	3450	2188	7 Sep	44000	7250	4200	9500	5821
Geometric mean	1203	1334	943	1371	1070	Geometric mean	6171	2946	3081	4119	3953
4 May	646	396	410	640	530	20 Jul	1967	4000	4450	4450	7600
19 May	2714	1700	1050	1767	1625	3 Aug	17000	1390	1367	2480	1920
2 Jun	2364	3500	3050	3652	3100	17 Aug	3800	2800	5300	5900	5750
15 Jun	4500	3150	1417	3450	2188	7 Sep	44000	7250	4200	9500	5821
6 Jul	1600	1967	2050	1917	1975	21 Sep	16875	3083	2333	8625	6000
Geometric mean	1972	1710	1307	1938	1631	Geometric mean	9884	3223	3162	5565	4936
19 May	2714	1700	1050	1767	1625	3 Aug	17000	1390	1367	2480	1920
2 Jun	2364	3500	3050	3652	3100	17 Aug	3800	2800	5300	5900	5750
15 Jun	4500	3150	1417	3450	2188	7 Sep	44000	7250	4200	9500	5821
6 Jul	1600	1967	2050	1917	1975	21 Sep	16875	3083	2333	8625	6000
20 Jul	1967	4000	4450	4450	7600	5 Oct	5214	4550	4450	4393	3300
Geometric mean	2464	2715	2106	2856	2778	Geometric mean	12012	3308	3162	5550	4178
2 Jun	2364	3500	3050	3652	3100	17 Aug	3800	2800	5300	5900	5750
15 Jun	4500	3150	1417	3450	2188	7 Sep	44000	7250	4200	9500	5821
6 Jul	1600	1967	2050	1917	1975	21 Sep	16875	3083	2333	8625	6000
20 Jul	1967	4000	4450	4450	7600	5 Oct	5214	4550	4450	4393	3300
3 Aug	17000	1390	1367	2480	1920	19 Oct	533	533	567	300	200
Geometric mean	3557	2608	2220	3056	2872	Geometric mean	6010	2731	2651	3638	2658

POOR ORIGINAL

5.0 THERMAL PLUME MAPPING

The TMINS plume was monitored to provide temperature data, define the discharge plume, and check the accuracy of the analytical plume model required in the EIS, Appendix B, Section 4.3.1.

5.1 METHODS

The TMINS discharge plume was mapped semimonthly February through November 1976. Additional mappings were conducted during the TMINS refueling shutdown on 21 February. No mappings were conducted in January or December because of ice. Vertical temperature profiles were taken from surface to bottom at 0.5 m intervals with an Endeco Digital Thermometer. The thermister was attached to one end of a 3 m pole (marked off in 0.5 m intervals) to allow consistent vertical measurements. Vertical profiles were taken at 5 m, 20 m, and 40 m from shore.

Two control (ambient river temperature) transects were established upstream of the TMINS Discharge at the Unit 1 Intake and 25 m upstream from the Discharge. Indicator transects were established at the Discharge, and 25 m, 50 m, 100 m, 200 m, 400 m, and 1900 m downstream of the Discharge. Markers were placed on shore at each transect. Station operation level, effluent flow rate, intake and discharge temperature, wind speed and direction, and the number of pumps in use (nuclear service, secondary service, and decay heat) were obtained from TMINS. River flow (ft^3/sec measured at the Harrisburg River Forecast Center) and air temperature were also recorded.

A return of the discharge temperature to within 2.7 C of ambient was used to define the plume.

1565 216

5.2 RESULTS

The TMINS discharge plume was mapped 33 times in 1976 at various river flow and station operation levels (Tables 5.2-1 through 5.2-33). Surface discharge and ambient river temperature differences are summarized in Table 5.2-34.

The delta T at the Discharge ranged from 0.9 C below to 0.6 C above ambient temperature from April through September (Table 5.2-34). The delta T was within the range established in the ETS, Appendix B, "the discharge temperature shall be no greater than 3 F [1.7 C] below inlet temperature or 7 F [3.9 C] above inlet temperature," for normal operation. From February through March, and October through November the allowable ETS discharge temperature is to be "no greater than 3 F [1.7 C] below inlet temperature or 12 F [6.7 C] above inlet temperature." The delta T ranged from 0.4 C below to 4.6 C above ambient temperature and was within the allowable limits.

The Pennsylvania Code, Title 25, Chapter 97.82 states, "The heat content of discharges shall be limited to an amount which could not raise the temperature of the entire stream at the point of discharge 5 F [2.7 C] above ambient temperature or a maximum of 87 F [30.6 C], whichever is less." In 28 of 33 surveys the plume was limited to 5 m offshore and 25 m downstream from the Discharge. The dates the plume exceeded these limits were on 16 February and during the cooldown operation for refueling on 21 February. On 16 February the plume extended 5 m offshore and 50 m downstream from the Discharge. The highest ΔT recorded during the cooldown operations was 4.6 C. At 25 m downstream from the Discharge

the ΔT was 1.7 C. The ETS, Appendix B states that "during reactor cool-down conditions discharge temperature shall not exceed 20 F [11.1 C] above inlet temperature." This temperature was never exceeded.

The analytical plume model was described in the Final Environmental Statement Related to Operation of Three Mile Island Nuclear Station Units 1 and 2 (A.E.C. 1972) for normal cooldown conditions. This was compared with the ten plume maps taken during the cooldown for refueling of Unit 1. In eight of the ten surveys the plume extended no further than 5 m offshore and to 1000 m downstream from the Discharge, where the temperature was within ten dilutions of the ambient temperature. In the surveys at 2200 hr and 2330 hr on 21 February the plume was detectible 20 m offshore, 25 m downstream from the Discharge. The plume was back to 5 m offshore, 50 m downstream from the Discharge. This varied from the plume model. The model used a river flow of 10,000 cfs; the river flow during the cooldown operations ranged from 193,000 to 229,000 cfs. In the model the plume extends 225 ft (68.6 m) into the river and travels 300 ft (91.4 m) downstream before reaching the 10 dilution criteria. During the cooldown operation the plume extended 5 to 20 m offshore and travel 1000 m downstream from the Discharge.

A.E.C 1972. Final environmental statement related to operation of Three Mile Island Nuclear Station. United States Atomic Energy Commission, Washington, D.C. pp. D30-D34.

Pennsylvania Code. No date. Title 25, Rules and regulations. Chapter 97.82. Industrial wastes, heat, pollution, allowable discharges.

POOR ORIGINAL

1565 218

Thermal plume data* for TMIIS on 16 February 1976.

Station Operation Level (m): 100 Time: 1330
 Nuclear Service Pumps: NA Intake Temp. (C): 2.2
 Secondary Service Pumps: NA Effluent Temp. (C): 6.9
 Decay Heat Pumps: NA Air Temp. (C): 13
 Effluent Rate (cfs): 20.1 Wind Speed (mph): 3
 River Flow (cfs): 55,400 Wind Dir.: N

Distance from Three Mile Island Shore				
40 m	20 m	5 m	Depth	
2.0	2.1	2.1	S**	Unit 1 Intake
1.9	2.1	2.1	0.5 m	
1.9	2.1	2.1	1.0	
1.9	2.1	2.1	1.5	
1.9	2.1	2.1	2.0	
1.9	2.1	2.1	2.5	
		2.0	3.0	
2.0	2.1	2.1	S	25 m Upstream of Discharge
2.0	2.1	2.1	0.5	
2.0	2.1	2.1	1.0	
2.0	2.1		1.5	
2.0	2.1		2.0	
2.0	2.1		2.5	
2.0	2.1	4.7	S	Discharge (D)
2.0	2.1	5.1	0.5	
2.0	2.1	5.4	1.0	
2.0	2.1	5.6	1.5	
2.0	2.1	5.9	2.0	
2.0	2.1	5.0	S	25 m Downstream of D
2.0	2.1	4.9	0.5	
2.0	2.1	4.9	1.0	
2.0	2.1		1.5	
2.0	2.1		2.0	
2.0	2.1	2.5	S	50 m Downstream of D
2.0	2.1	2.7	0.5	
2.0	2.1	2.8	1.0	
2.0	2.1		1.5	
2.0	2.1		2.0	
2.0	2.1		2.5	
2.0	2.1	2.7	S	100 m Downstream of D
2.0	2.1	2.7	0.5	
2.0	2.1	3.0	1.0	
2.0	2.1		1.5	
2.0	2.1		2.0	
2.0	2.1		2.5	
2.0	2.1	3.3	S	200 m Downstream of D
2.0	2.1	3.3	0.5	
2.0	2.1	3.3	1.0	
2.0	2.1		1.5	
2.0	2.1		2.0	
2.0	2.1		2.5	
2.0	2.1	2.9	S	400 m Downstream of D
2.0	2.1	3.0	0.5	
2.0	2.1	3.0	1.0	
2.0	2.1		1.5	
2.0	2.1		2.0	
2.0	2.1		2.5	
2.0	2.1		3.0	
2.1	2.1	2.7	S	800 m Downstream of D
2.0	2.1	2.7	0.5	
2.0	2.1	2.7	1.0	
2.0	2.1		1.5	
2.0	2.1		2.0	
2.0	2.1		2.5	

Distance from Three Mile Island Shore				
40 m	20 m	5 m	Depth	
2.1	2.1	2.4	S	1500 m Downstream of D
2.0	2.1	2.5	0.5 m	
2.0	2.1	2.5	1.0	
2.0	2.0		1.5	
2.0	2.0		2.0	
2.0	2.0		2.5	
2.0	2.1	2.4	S	1900 m Downstream of D
2.0	2.1	2.5	0.5	
2.0	2.1	2.5	1.0	
2.0	2.1	2.5	1.5	
2.0	2.1	2.5	2.0	
2.0	2.1	*	2.5	
2.0			3.0	

* Temperature in C.

** S = Surface.

NA = Not Available.

POOR ORIGINAL

1565 219

Table 5.2-2

Thermal plume data for THINS on 20 February 1976

Station Operation Level (T): 0			Time: 1415		
Nuclear Service Pumps: 2			Intake Temp. (C): 1.5		
Secondary Service Pumps: 2			Effluent Temp. (C): 3.5		
Decay Heat Pumps: 1			Air Temp. (C): 8.2		
Effluent Rate (cfs): 37.9			Wind Speed (mph): 6		
River Flow (cfs): 210,000			Wind Dir.: NW		

Distance From Three Mile Island Shore			Distance From Three Mile Island Shore		
20 m	5 m	Depth	20 m	5 m	Depth
2.4	2.4	5	2.4	2.6	5
2.4	2.4	0.5 m	2.4	2.6	0.5 m
2.4	2.4	1.0	2.4	2.6	1.0
2.4	2.4	1.5	2.4	2.6	1.5
2.4	2.4	2.0	2.4		2.0
2.4	2.4	2.5	2.4		2.5
2.4	2.4	3.0	2.4		3.0
2.4	2.4	5	2.5	5	300 m Downstream of D
2.4	2.4	0.5	2.5	0.5	
2.4	2.4	1.0	2.5	1.0	
2.4	2.4	1.5			
2.4		2.0	2.5	5	400 m Downstream of D
2.4		2.5	2.5	0.5	
2.4		3.0	2.5	1.0	
2.4		3.5			
2.4	3.0	5	2.4	2.5	5
2.4	3.1	0.5	2.4	2.5	0.5
2.4	3.4	1.0	2.4	2.5	1.0
2.4	3.6	1.5	2.4	2.5	1.5
2.4	3.6	2.0	2.4	2.5	2.0
2.4		2.5	2.4		2.5
2.4		3.0	2.4		3.0
2.4	2.7	5	2.4	2.5	5
2.4	2.7	0.5	2.4	2.5	0.5
2.4	2.8	1.0	2.4	2.5	1.0
2.4	2.8	1.5	2.4		1.5
2.4		2.0	2.4		2.0
2.4		2.5	2.4		2.5
2.4			2.4		3.0
2.4			2.4		3.5
2.4	2.7	5		2.5	5
2.4	2.7	0.5		2.5	0.5
2.4	2.6	1.0		2.5	1.0
2.4	2.6	1.5		2.5	1.5
2.4		2.0		2.5	2.0
2.4		2.5		2.5	2.5
2.4		3.0		2.5	3.0
	2.7	5	2.4	2.5	5
	2.7	0.5	2.4	2.5	0.5
	2.7	1.0	2.4	2.5	1.0
	2.7	1.5	2.4		1.5
			2.4		2.0
			2.4		2.5
			2.4		3.0
			2.4		3.5
	2.7	5			
	2.7	0.5			
	2.6	1.0			
	2.6	1.5			
2.4	2.6	5			
2.4	2.5	0.5			
2.4	2.5	1.0			
2.4	2.5	1.5			
2.4	2.5	2.0			
2.4		2.5			
2.4		3.0			

POOR ORIGINAL

1565 220

POOR ORIGINAL

Table 5.2-3

Thermal plume data for TMINS on 21 February 1976.

Station Operation Level (%): 0			Time: 0700		
Nuclear Service Pumps: 2			Intake Temp. (C): 1.4		
Secondary Service Pumps: 3			Effluent Temp. (C): 2.8		
Decay Heat Pumps: 0			Air Temp. (C): 3.0		
Effluent Rate (cfs): 55.7			Wind Speed (mph): 8		
River Flow (cfs): 25,000			Wind Dir.: SE		
Distance From Three Mile Island shore					
20 m	5 m	Depth	Distance From Three Mile Island shore		
			20 m	5 m	Depth
2.2	2.2	S	2.2	2.3	S
2.2	2.2	0.5 m	2.2	2.3	0.5 m
2.2	2.3	1.0	2.2	2.3	1.0
2.2	2.3	1.5	2.2	2.4	1.5
2.2	2.3	2.0	2.2	2.4	2.0
2.2	2.3	2.5	2.2		2.5
2.2	2.2	3.0	2.2		3.0
2.2	2.3	3.5	2.2		3.5
2.2	2.2	4.0	2.2	2.4	S
			2.2	2.3	S
2.2	2.2	S	2.2	2.4	0.5
2.2	2.2	0.5	2.2	2.4	1.0
2.2	2.2	1.0	2.2	2.4	1.5
2.2	2.2	1.5	2.2		2.0
2.2	2.3	2.0	2.2		2.5
2.2		2.5	2.2		3.0
2.2		3.0			
2.2		3.5	2.2	2.3	S
			2.2	2.4	0.5
2.2	2.2	S	2.2	2.4	1.0
2.2	2.3	0.5	2.2	2.4	1.5
2.2	2.2	1.0	2.2		2.0
2.2	2.3	1.5	2.2		2.5
2.2	2.8	2.0	2.2		3.0
2.2		2.5	2.2		3.5
2.2		3.0			
2.2		3.5	2.2	2.3	S
			2.2	2.3	0.5
2.2	2.4	S	2.2	2.2	1.0
2.2	2.4	0.5	2.2	2.3	1.5
2.2	2.4	1.0	2.2		2.0
2.2	2.5	1.5	2.2		2.5
2.2		2.0	2.2		3.0
2.2		2.5	2.2		3.5
2.2		3.0	2.2		4.0
2.2		3.5			
			2.2	2.3	S
2.2	2.3	S	2.2	2.3	0.5
2.2	2.3	0.5	2.2	2.3	1.0
2.2	2.3	1.0	2.2	2.3	1.5
2.2	2.4	1.5	2.2		2.0
2.2		2.0	2.2		2.5
2.2		2.5	2.2		3.0
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4				

1565 221

Thermal plume data for INRS on 21 February 1976.

Station Operation Level (C): 0			Time: 0615			
Nuclear Service Pumps: 2			Intake Temp. (C): 1.3			
Secondary Service Pumps: 3			Effluent Temp. (C): 2.8			
Decay Heat Pumps: 0			Air Temp. (C): 3.0			
Effluent Rate (cfs): 44.6			Wind Speed (mph): 8			
River Flow (cfs): 200,000			Wind Dir.: SSE			
Distance From Three Mile Island Shore			Distance From Three Mile Island Shore			
20 m	5 m	Depth	20 m	5 m	Depth	
2.2	2.2	S	2.2	2.4	S	125 m Downstream of D
2.2	2.2	0.5 m	2.2	2.4	0.5 m	
2.2	2.2	1.0	2.2	2.4	1.0	
2.2	2.2	1.5	2.2		1.5	
2.2	2.2	2.0	2.2		2.0	
2.2	2.2	2.5	2.2		2.5	
2.2	2.2	3.0	2.2		3.0	
2.2	2.2	3.5	2.2		3.5	
	2.2	4.0				
2.2	2.5	S	2.2	2.4	S	200 m Downstream of D
2.2	2.2	0.5	2.2	2.4	0.5	
2.2	2.2	1.0	2.2	2.4	1.0	
2.2	2.2	1.5	2.2	2.4	1.5	
2.2		2.0	2.2		2.0	
2.2		2.5	2.2		2.5	
2.2		3.0	2.2		3.0	
2.2		3.5	2.2		3.5	
2.2	2.7	S	2.2	2.3	S	400 m Downstream of D
2.2	2.7	0.5	2.2	2.3	0.5	
2.2	2.7	1.0	2.2	2.3	1.0	
2.2	2.7	1.5	2.2	2.3	1.5	
2.2	2.7	2.0	2.2		2.0	
2.2	2.8	2.5	2.2		2.5	
2.2		3.0	2.2		3.0	
			2.2		3.5	
2.2	2.4	S	2.2	2.3	S	800 m Downstream of D
2.2	2.5	0.5	2.2	2.3	0.5	
2.2	2.5	1.0	2.2	2.3	1.0	
2.2		1.5	2.2	2.3	1.5	
2.2		2.0	2.2		2.0	
2.2		2.5	2.2		2.5	
2.2		3.0	2.2		3.0	
2.2		3.5	2.2		3.5	
2.2	2.4	S	2.2	2.3	S	1000 m Downstream of D
2.2	2.4	0.5	2.2	2.3	0.5	
2.2	2.4	1.0	2.2	2.3	1.0	
2.2		1.5	2.2	2.3	1.5	
2.2		2.0	2.2	2.3	2.0	
2.2		2.5	2.2		2.5	
2.2		3.0	2.2		3.0	
2.2		3.5	2.2		3.5	
2.2	2.3	S				
2.2	2.3	0.5				
2.2	2.3	1.0				
2.2	2.4	1.5				
2.2		2.0				
2.2		2.5				
2.2		3.0				
2.2		3.5				
2.2	2.3	S				
2.2	2.3	0.5				
2.2	2.3	1.0				
2.2	2.3	1.5				
2.2	2.4	2.0				
2.2		2.5				
2.2		3.0				
2.2		3.5				

POOR ORIGINAL

1565 222

POOR ORIGINAL

Table 5.2-5

Thermal plume data for THINS on 21 February 1970.

Station Operation Level (ft): 0			Time: 1000		
Nuclear Service Pumps: 2			Intake Temp. (C): 1.4		
Secondary Service Pumps: 2			Effluent Temp. (C): 2.8		
Decay Heat Pumps: 0			Air Temp. (C): 8.0		
Effluent Rate (cfs): 45.6			Wind speed (mph): 10		
River Flow (cfs): 229,000			Wind Dir.: S		
Distance from Three Mile Island shore			Distance from Three Mile Island shore		
20 m	5 m	Depth	20 m	5 m	Depth
2.2	2.3	S	2.2	2.4	S
2.2	2.3	0.5 m	2.2	2.4	0.5 m
2.2	2.3	1.0	2.2	2.4	1.0
2.2	2.3	1.5	2.2	2.4	1.5
2.2	2.3	2.0	2.2		2.0
2.2	2.3	2.5	2.2		2.5
2.2	2.3	3.0	2.2		3.0
2.2	2.3	3.5	2.2		3.5
2.2	2.3	4.0			
25 m Upstream of Discharge			2.2	2.3	S
2.2	2.2	0.5	2.2	2.3	0.5
2.2	2.2	1.0	2.2	2.3	1.0
2.2	2.2	1.5	2.2	2.3	1.5
2.2	2.2	2.0	2.2		2.0
2.2	2.2	2.5	2.2		2.5
2.2	2.2	3.0	2.2		3.0
2.2	2.2	3.5	2.2		3.5
Discharge (D)			2.2	2.3	S
2.2	2.6	0.5	2.2	2.3	0.5
2.2	2.7	1.0	2.2	2.3	1.0
2.2	2.7	1.5	2.2		1.5
2.2	2.6	2.0	2.2		2.0
2.2	2.6	2.5	2.2		2.5
2.2	2.7	3.0	2.2		3.0
2.2	2.8	3.5	2.2		3.5
25 m Downstream of D					
2.2	2.5	S			
2.2	2.5	0.5			
2.2	2.5	1.0			
2.2		1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
50 m Downstream of D					
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.5	1.0			
2.2		1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
75 m Downstream of D					
2.2	2.4	S			
2.2	2.4	0.5			
2.2	2.4	1.0			
2.2	2.5	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			
100 m Downstream of D					
2.2	2.4	S			
2.2	2.5	0.5			
2.2	2.4	1.0			
2.2	2.4	1.5			
2.2		2.0			
2.2		2.5			
2.2		3.0			
2.2		3.5			

1565 223

Table 5.2-6

Thermal plume data for THIN on 21 February 1976.

Station operation level: 0	Time: 1445
Discharge: Pumps: 2	Intake Temp. (C): 1.6
Effluent: Pumps: 2	Effluent Temp. (C): 3.8
Heat Pump: 6	Air Temp. (C): 15.0
Effluent Rate: 37.9	Wind speed (mph): 8
Water Temp. (C): 12.0	Wind Temp. (C): 15.0

Distance from Three Mile Island Shore			Distance from Three Mile Island Shore		
20 m	5 m	Depth	20 m	5 m	Depth
2.5	2.5	8	2.5	2.7	8
2.5	2.5	0.5 m	2.5	2.7	0.5 m
2.5	2.5	1.0	2.5	2.7	1.0
2.5	2.5	1.5	2.5		1.5
2.5	2.5	2.0	2.5		2.0
2.5	2.5	2.5	2.5		2.5
2.5	2.5	3.0	2.5		3.0
2.5	2.5	3.5			
2.5	2.5	4.0			
2.5	2.5	4.5	2.5	2.7	8
			2.5	2.7	0.5
2.5	2.5	8	2.5	2.7	1.0
2.5	2.5	0.5	2.5	2.7	1.5
2.5	2.5	1.0	2.5		2.0
2.5		1.5	2.5		2.5
2.5		2.0	2.5		3.0
2.5		2.5			
2.5		3.0	2.5	2.7	8
2.5		3.5	2.5	2.7	0.5
2.5		4.0	2.5	2.7	1.0
2.5		4.5	2.5	2.7	1.5
			2.5		2.0
2.5	2.7	8	2.5		2.5
2.5	2.7	0.5	2.5		3.0
2.5	2.7	1.0			
2.5	2.7	1.5	2.5	2.7	8
2.5	2.7	2.0	2.5	2.7	0.5
2.5	2.7	2.5	2.5	2.7	1.0
2.5	2.7	3.0	2.5		1.5
			2.5		2.0
2.5	2.7	8	2.5		2.5
2.5	2.7	0.5			
2.5	2.7	1.0	2.5	2.6	8
2.5	2.7	1.5	2.5	2.6	0.5
2.5	2.7	2.0	2.5	2.6	1.0
2.5	2.7	2.5	2.5		1.5
2.5	2.7	3.0	2.5		2.0
			2.5		2.5
2.5	2.7	8	2.5		3.0
2.5	2.7	0.5			
2.5	2.7	1.0			
2.5		1.5			
2.5		2.0			
2.5		2.5			
2.5		3.0			
2.5	2.7	8			
2.5	2.7	0.5			
2.5	2.7	1.0			
2.5	2.7	1.5			
2.5	2.7	2.0			
2.5	2.7	2.5			
2.5	2.7	3.0			
2.5	2.7	8			
2.5	2.7	0.5			
2.5	2.7	1.0			
2.5	2.7	1.5			
2.5	2.7	2.0			
2.5	2.7	2.5			
2.5	2.7	3.0			

POOR ORIGINAL

1565 224

Table 5.2-7

Thermal plume data for THMS on 21 February 1976.

Station Operation Level (ft): 0	Time: 1635
Nuclear Service Pumps: 1	Intake Temp. (°C): 1.9
Secondary Service Pumps: 2	Effluent Temp. (°C): 4.4
Decay Heat Pumps: 0	Air Temp. (°C): 10.5
Effluent Rate (cfs): 44.6	Wind Speed (mph): 10
River Flow (cfs): 219,000	Wind Dir.: S

Distance from Three Mile Island shore

20 m	5 m	Depth	
2.8	2.7	S	Unit 1 Intake
2.8	2.7	0.5 m	
2.8	2.7	1.0	
2.7	2.7	1.5	
2.7	2.7	2.0	
2.7	2.7	2.5	
2.7	2.7	3.0	
	2.7	3.5	
	2.7	4.0	
2.7	2.7	S	25 m Upstream of Discharge
2.7	2.7	0.5	
2.7	2.7	1.0	
2.7	2.7	1.5	
2.7		2.0	
2.7		2.5	
2.7		3.0	
2.7	3.3	S	Discharge (D)
2.7	3.4	0.5	
2.7	3.4	1.0	
2.7	3.4	1.5	
2.7	3.4	2.0	
2.7	3.4	2.5	
2.7	3.5	3.0	
2.7	3.2	S	25 m Downstream of D
2.7	3.2	0.5	
2.7	3.2	1.0	
2.7		1.5	
2.7		2.0	
2.7		2.5	
2.7		3.0	
2.7	3.2	S	50 m Downstream of D
2.7	3.2	0.5	
2.7	3.2	1.0	
2.7		1.5	
2.7		2.0	
2.7		2.5	
2.7		3.0	
2.7	3.0	S	100 m Downstream of D
2.7	3.0	0.5	
2.7	3.0	1.0	
2.7	3.0	1.5	
2.7		2.0	
2.7		2.5	
2.7		3.0	
2.7	3.0	S	200 m Downstream of D
2.7	3.0	0.5	
2.7	3.0	1.0	
2.7	3.0	1.5	
2.7		2.0	
2.7		2.5	
2.7		3.0	

POOR ORIGINAL

Distance from Three Mile Island shore

20 m	5 m	Depth	
2.8	2.9	S	400 m Downstream of D
2.8	2.9	0.5 m	
2.8	2.9	1.0	
2.8	2.9	1.5	
2.8		2.0	
2.8		2.5	
2.8	2.9	S	500 m Downstream of D
2.8	2.9	0.5	
2.8	2.9	1.0	
2.8	2.9	1.5	
2.8		2.0	
2.8		2.5	
2.8	2.8	S	700 m Downstream of D
2.8	2.8	0.5	
2.8	2.8	1.0	
2.8	2.8	1.5	
2.8		2.0	
2.8		2.5	
2.8	2.8	S	800 m Downstream of D
2.8	2.8	0.5	
2.8	2.8	1.0	
2.8	2.8	1.5	
2.8		2.0	
2.8		2.5	
2.8	2.8	S	1000 m Downstream of D
2.8	2.8	0.5	
2.8	2.8	1.0	
2.8	2.8	1.5	
2.8		2.0	
2.8		2.5	
2.8		3.0	
2.8		3.5	

1565 225

POOR ORIGINAL

Table 5.2-8

Thermal pulse data for THTO on 21 February 1976.

Station operation level (): 0			Time: 1710				
Nuclear Service pumps: 2			Intake Temp. (C): 1.8				
Secondary Service Pumps: 3			Effluent Temp. (C): 3.9				
Decay Heat Pumps: 0			Air Temp. (C): 15.0				
Effluent Rate (cfs): 44.6			Wind Speed (mph): 0				
River Flow (cfs): 11,000			Discharge (D)				
Distance from Three Mile Island shore				Distance from Three Mile Island shore			
20 m	5 m	Depth		20 m	5 m	Depth	
2.6	2.7	S	Unit 1 Intake	2.7	2.8	S	400 m Downstream of D
2.6	2.7	0.5 m		2.7	2.8	0.5 m	
2.6	2.7	1.0		2.7	2.8	1.0	
2.6	2.7	1.5		2.7	2.8	1.5	
2.6	2.7	2.0		2.7	2.8	2.0	
2.6	2.7	2.5		2.7	2.8	2.5	
2.6	2.7	3.0		2.7	2.8	3.0	
2.6	2.7	3.5					
2.7	2.7	S	25 m Upstream of Discharge	2.7	2.8	S	500 m Downstream of D
2.7	2.7	0.5		2.7	2.8	0.5	
2.7	2.7	1.0		2.7	2.8	1.0	
2.7	2.7	1.5		2.7	2.8	1.5	
2.7	2.7	2.0		2.7	2.8	2.0	
2.7	2.7	2.5		2.7	2.8	2.5	
2.7	2.7	3.0		2.7	2.8	3.0	
2.7	3.1	S	Discharge (D)	2.7	2.8	3.5	
2.7	3.1	0.5		2.7	2.8	4.0	
2.7	3.1	1.0		2.7	2.8	S	700 m Downstream of D
2.7	3.2	1.5		2.7	2.8	0.5	
2.7	3.2	2.0		2.7	2.8	1.0	
2.7	3.2	2.5		2.7	2.8	1.5	
2.7	3.2	3.0		2.7	2.8	2.0	
2.7	3.1	S	25 m Downstream of D	2.7	2.8	2.5	
2.7	3.1	0.5		2.7	2.8	3.0	800 m Downstream of D
2.7	3.1	1.0		2.7	2.8	3.5	
2.7	3.1	1.5		2.7	2.8	4.0	
2.7	3.1	2.0		2.7	2.8	S	
2.7	3.1	2.5		2.7	2.8	0.5	
2.7	3.1	3.0		2.7	2.8	1.0	
2.7	3.0	S	50 m Downstream of D	2.7	2.8	1.5	900 m Downstream of D
2.7	3.0	0.5		2.7	2.8	2.0	
2.7	2.9	1.0		2.7	2.8	2.5	
2.7	2.9	1.5		2.7	2.8	3.0	
2.7	2.9	2.0		2.7	2.8	3.5	
2.7	2.9	2.5		2.7	2.8	S	1000 m Downstream of D
2.7	2.9	3.0		2.7	2.8	0.5	
2.7	2.8	S	100 m Downstream of D	2.7	2.8	1.0	
2.7	2.8	0.5		2.7	2.8	1.5	
2.7	2.8	1.0		2.7	2.8	2.0	
2.7	2.8	1.5		2.7	2.8	2.5	
2.7	2.8	2.0		2.7	2.8	3.0	
2.7	2.8	2.5		2.7	2.8	3.5	
2.7	2.8	3.0					
2.7	2.8	3.5					
2.7	2.8	S	200 m Downstream of D				
2.7	2.8	0.5					
2.7	2.8	1.0					
2.7	2.8	1.5					
2.7	2.8	2.0					
2.7	2.8	2.5					
2.7	2.8	3.0					

1565 226

1565 226

Thermal plume data for IMMS on 21 February 1976.

POOR ORIGINAL

1565 227

POOR ORIGINAL

Table 5.2-10

Thermal plume data for N400 on 21 February 1976.

Station Operation Level 100 ft		Time: 1200	
Nuclear Service Pumps 2		Intake Temp. (C): 2.1	
Secondary Service Pumps 2		Effluent Temp. (C): 6.4	
Decay Heat Pumps 2		Air Temp. (C): 5A	
Effluent Rate (cfs): 111.4		Wind Speed (mph): 9	
River Flow (cfs): 1000		Wind Dir: S	

Distance from Three Mile Island shore			Distance from Three Mile Island shore		
20 m	5 m	Depth	20 m	5 m	Depth
2.9	2.4				
2.9	2.8	0.5 m			400 m Downstream of D
2.9	2.8	1.0	2.9	3.5	0.5 m
2.9	2.8	1.5	2.9	3.5	1.0
2.9	2.8	2.0	2.9	3.5	1.5
2.9	2.8	2.5	2.9	3.5	2.0
2.9	2.8	3.0	2.9		2.5
2.9	2.8	3.5	2.9		3.0
	2.8	4.0			
2.9	2.9	S	2.9	3.5	S
2.9	2.9	0.5	2.9	3.5	0.5
2.9	2.9	1.0	2.9	3.5	1.0
2.9	2.9	1.5	2.9		1.5
2.9		2.0	2.9		2.0
2.9		2.5	2.9		2.5
2.9		3.0	2.9		3.0
			2.9		3.5
			2.9		4.0
2.9	5.0		2.9	3.4	S
2.9	5.2	0.5	2.9	3.4	0.5
2.9	5.3	1.0	2.9	3.4	1.0
2.9	5.7	1.5	2.9	3.4	1.5
2.9	5.8	2.0	2.9		2.0
2.9	5.3	2.5	2.9		2.5
2.9	5.2	3.0	2.9		3.0
			2.9		3.5
			2.9		4.0
3.5	4.3	S	2.9	3.3	S
3.4	4.3	0.5	2.9	3.3	0.5
3.3	4.3	1.0	2.9	3.3	1.0
3.3		1.5	2.9		1.5
3.3		2.0	2.9		2.0
3.3		2.5	2.9		2.5
2.9	4.2	S	2.9		3.0
2.9	4.2	0.5	2.9		3.5
2.9	4.2	1.0			
2.9		1.5	3.0	3.2	S
2.9		2.0	3.0	3.2	0.5
2.9		2.5	3.0	3.2	1.0
2.9		3.0	3.0	3.2	1.5
			3.0		2.0
			3.0		2.5
3.1	4.1	S			
3.0	4.1	0.5			
3.0	4.1	1.0			
3.0		1.5			
3.0		2.0			
3.0		2.5			
2.9		3.0			
2.9	3.9	S			
2.9	3.9	0.5			
2.9	3.9	1.0			
2.9		1.5			
2.9		2.0			
2.9		2.5			
2.9		3.0			
2.9		3.5			

1565 228

POOR ORIGINAL

Table 5.2-11

Thermal plume data for TMINS on 21 February 1976.

Station Operation Level (C): 0			Time: 2330
Nuclear Service Pumps: 2			Intake Temp. (C): 2.2
Secondary Service Pumps: 3			Effluent Temp. (C): 6.9
Decay Heat Pumps: 1			Air Temp. (C): NA
Effluent Rate (cfs): 115.9			Wind Speed (mph): 10
River Flow (cfs): 201,000			Wind Dir.: S
Distance from Three Mile Island shore			
20 m	5 m	Depth	
3.0	3.0	S	Unit 1 Intake
3.0	3.0	0.5 m	
3.0	3.0	1.0	
3.0	3.0	1.5	
3.0	3.0	2.0	
3.0	3.0	2.5	
3.0	3.0	3.0	
3.0	3.0	3.5	
	3.0	4.0	
3.0	3.0	S	25 m Upstream of Discharge
3.0	3.0	0.5	
3.0	3.0	1.0	
3.0	3.0	1.5	
3.0		2.0	
3.0		2.5	
3.0	7.6	S	Discharge (D)
3.0	6.0	0.5	
3.0	7.0	1.0	
3.0	7.3	1.5	
3.0	7.3	2.0	
3.0	7.4	2.5	
3.0	7.3	3.0	
3.1	4.7	S	25 m Downstream of D
3.1	4.7	0.5	
3.1	4.7	1.0	
3.4		1.5	
3.6		2.0	
3.7		2.5	
3.3	4.6	S	50 m Downstream of D
3.1	4.7	0.5	
3.1	4.7	1.0	
3.1		1.5	
3.1		2.0	
3.0		2.5	
3.0		3.0	
3.0		3.5	
3.1	4.3	S	100 m Downstream of D
3.1	4.3	0.5	
3.1	4.4	1.0	
3.1	4.4	1.5	
3.0		2.0	
3.0		2.5	
3.0		3.0	
3.0		3.5	
3.4	4.0	S	200 m Downstream of D
3.3	4.0	0.5	
3.3	4.0	1.0	
3.2	4.0	1.5	
3.2		2.0	
3.1		2.5	
3.1		3.0	

Distance from Three Mile Island shore			
20 m	5 m	Depth	
3.1	3.7	S	400 m Downstream of D
3.1	3.7	0.5 m	
3.1	3.7	1.0	
3.1	3.7	1.5	
3.1	3.7	2.0	
3.1		2.5	
3.1		3.0	
3.2	3.6	S	500 m Downstream of D
3.1	3.6	0.5	
3.1	3.6	1.0	
3.2	3.6	1.5	
3.2		2.0	
3.2		2.5	
3.2		3.0	
3.2		3.5	
3.2	3.5	S	700 m Downstream of D
3.1	3.5	0.5	
3.1	3.5	1.0	
3.2		1.5	
3.2		2.0	
3.2		2.5	
3.2		3.0	
3.2	3.4	S	800 m Downstream of D
3.2	3.4	0.5	
3.2	3.4	1.0	
3.2		1.5	
3.3		2.0	
3.3		2.5	
3.2	3.4	S	1000 m Downstream of D
3.2	3.4	0.5	
3.2	3.4	1.0	
3.2	3.4	1.5	
3.2		2.0	
3.2		2.5	

1565 229

POOR ORIGINAL

Thermal plume data for IMFA on 22 February 1976.

Station operation level: 1 at 0			Time: 1100		
Nuclear Service Pumps: 2			Intake Temp. (C): 2.3		
Secondary Service Pumps: 3			Effluent Temp. (C): 6.3		
Leakage Pumps: 2			Air Temp. (C): NA		
Effluent Rate (cfs): 111.4			Wind Speed (mph): 20		
Water Temp (C): 1.4			Wind Dir:		
Distance from Three Mile Island shore			Distance from Three Mile Island shore		
20 m	5 m	Depth	20 m	5 m	Depth
3.1	3.1		3.2	3.7	
3.1	3.1	0.5 m	3.2	3.7	0.5 m
3.1	3.1	1.0	3.2	3.7	1.0
3.1	3.1	1.5	3.2	3.7	1.5
3.1	3.1	2.0	3.2	3.7	2.0
3.1	3.1	2.5	3.2		2.5
3.1	3.1	3.0	3.2		3.0
3.1	3.1	3.5			
	3.1	4.0	3.3	3.7	
			3.3	3.7	0.5
3.1	3.2	5	3.3	3.7	1.0
3.1	3.2	0.5	3.3	3.7	1.5
3.1	3.2	1.0	3.3		2.0
3.1		1.5	3.3		2.5
3.1		2.0	3.3		3.0
3.1		2.5	3.3		3.5
3.1		3.0			
			3.3	3.7	0
3.1	3.7	5	3.3	3.7	0.5
3.1	3.7	0.5	3.3	3.7	1.0
3.1	3.7	1.0	3.3	3.7	1.5
3.1	3.7	1.5	3.3		2.0
3.1	3.7	2.0	3.3		2.5
3.1	3.7	2.5	3.3		3.0
3.1	3.7	3.0	3.3		3.5
3.1	4.4	5	3.3	3.6	5
3.1	4.4	0.5	3.3	3.6	0.5
3.1	4.4	1.0	3.3	3.6	1.0
3.1	4.4	1.5	3.3	3.6	1.5
3.1		2.0	3.3		2.0
3.1		2.5	3.3		2.5
3.1		3.0	3.3		3.0
3.1		3.5	3.3		3.5
3.2	4.3	5	3.3	3.5	5
3.2	4.3	0.5	3.3	3.5	0.5
3.1	4.3	1.0	3.3	3.5	1.0
3.1		1.5	3.3	3.5	1.5
3.1		2.0	3.3		2.0
3.1		2.5	3.3		2.5
3.1		3.0	3.3		3.0
3.1		3.5	3.3		3.5
3.1	4.2	5			
3.1	4.2	0.5			
3.1	4.2	1.0			
3.1		1.5			
3.1		2.0			
3.1		2.5			
3.1		3.0			
3.1		3.5			
3.1		4.0			
3.3	4.1	5			
3.3	4.2	0.5			
3.2	4.2	1.0			
3.2		1.5			
3.2		2.0			
3.2		2.5			
3.2		3.0			
3.2		3.5			

1565 230

POOR ORIGINAL

Table 5.2-13

Thermal plume data for TMINS on 23 February 1976.

Station Operation Level (C): 0				Time: 1-15
Nuclear Service Pumps: 1				Intake Temp. (C): 1.9
Secondary Service Pumps: 1				Effluent Temp. (C): 3.4
Decay Heat Pumps: 1				Air Temp. (C): 4.0
Effluent Rate (cfs): 39.0				Wind Speed (mph): 14
River Flow (cfs): 164,000				Wind Dir.: SW
Distance From Three Mile Island Shore				
40 m	20 m	5 m	Depth	
1.6			S	Unit 1 Intake
1.5	1.6	1.6	0.5 m	
1.6	1.6	1.6	1.0	
1.6	1.6	1.6	1.5	
1.6	1.6	1.6	2.0	
1.6	1.6	1.6	2.5	
1.5	1.6	1.6	3.0	
		1.6	3.5	
		1.6	4.0	
		1.6	4.5	
1.6	1.6	1.6	S	25 m Upstream of Discharge
1.6	1.6	1.6	0.5	
1.6	1.6	1.6	1.0	
1.6	1.6	1.6	1.5	
1.6	1.6		2.0	
1.6	1.6		2.5	
1.6			3.0	
1.6	1.6	1.8	S	Discharge (D)
1.6	1.6	1.7	0.5	
1.6	1.6	1.8	1.0	
1.6	1.6	2.2	1.5	
1.6	1.6	2.3	2.0	
1.6	1.6	2.3	2.5	
1.6			3.0	
1.6	1.6	1.7	S	25 m Downstream of D
1.6	1.6	1.8	0.5	
1.6	1.6	1.8	1.0	
1.6	1.6	1.8	1.5	
1.6	1.6		2.0	
1.6	1.6		2.5	
1.6	1.6		3.0	
1.6	1.6	1.7	S	50 m Downstream of D
1.6	1.6	1.7	0.5	
1.6	1.6	1.7	1.0	
1.6	1.6	1.8	1.5	
1.6	1.6	1.8	2.0	
1.6	1.6		2.5	
1.6	1.6		3.0	
1.6			3.5	
1.6	1.6	1.7	S	100 m Downstream of D
1.6	1.6	1.7	0.5	
1.6	1.6	1.7	1.0	
1.6	1.6	1.7	1.5	
1.6	1.6		2.0	
1.6	1.6		2.5	
1.6	1.6		3.0	

1565 231

Table 5.2-14

Thermal plume data for DMTO on 5 March 1976.

Station Operation Level (m) 0				Time: 11:5
Nuclear Service Pumps: 2				Intake Temp. (C): 14
Secondary Service Pumps: 1				Effluent Temp. (C): 4.5
Recycle Heat Pumps: 1				Air Temp. (C): 20.0
Effluent Rate (cfs): 40.0				Wind Speed (mph): 19
River Flow Rate (cfs): 10				Wind Dir.: 20
Distance from Three Mile Island Core				
50 m	20 m	5 m	Depth	
5.8	5.9	6.0	5	Unit 1 Intake
5.8	5.9	5.9	0.5 m	
5.8	5.9	5.9	1.0	
5.8	5.9	5.9	1.5	
5.8	5.9	5.9	2.0	
5.8	5.9	5.9	2.5	
5.8	5.9	5.9	3.0	
5.8	5.9	6.1	5	25 m Upstream of Discharge
5.8	5.9	6.1	0.5	
5.8	5.9	6.0	1.0	
5.8	5.9	6.1	1.5	
5.8	5.9	6.0	2.0	
5.8	5.9	6.1	2.5	
5.8	5.9	6.1	3.0	
5.8	5.9	6.2	5	Discharge (D)
5.8	5.9	6.1	0.5	
5.8	5.9	6.2	1.0	
5.8	5.9	6.4	1.5	
5.8	5.9	6.5	2.0	
5.8	5.9	6.5	2.5	
5.8	5.9	6.3	5	25 m Downstream of D
5.8	5.9	6.4	0.5	
5.8	5.9	6.4	1.0	
5.8	5.9	6.4	1.5	
5.8	5.9	6.4	2.0	
5.8	5.9	6.4	2.5	
5.8	5.9	6.4	3.0	
5.8	6.0	6.3	5	50 m Downstream of D
5.8	6.0	6.4	0.5	
5.8	5.9	6.4	1.0	
5.8	5.9	6.5	1.5	
5.8	5.9	6.5	2.0	
5.8	5.9	6.5	2.5	
5.8	5.9	6.5	3.0	
5.8	6.0	6.4	5	100 m Downstream of D
5.8	6.0	6.5	0.5	
5.8	5.9	6.5	1.0	
5.8	5.9	6.5	1.5	
5.8	5.9	6.5	2.0	
5.8	5.9	6.5	2.5	
5.8	5.9	6.5	3.0	
5.8	6.0	6.3	5	200 m Downstream of D
5.8	5.9	6.2	0.5	
5.8	5.9	6.3	1.0	
5.8	5.9	6.4	1.5	
5.8	5.9	6.4	2.0	
5.8	5.9	6.4	2.5	
5.8	5.9	6.4	3.0	
5.8	6.0	6.2	5	400 m Downstream of D
5.8	6.0	6.2	0.5	
5.8	6.0	6.2	1.0	
5.8	6.0	6.2	1.5	
5.8	6.0	6.2	2.0	
5.8	6.0	6.2	2.5	
5.8	6.0	6.2	3.0	
6.0	6.0	6.3	5	600 m Downstream of D
5.8	6.0	6.3	0.5	
6.0	6.0	6.3	1.0	
5.8	6.0	6.3	1.5	
5.8	6.0	6.3	2.0	
5.8	6.0	6.3	2.5	
5.8	6.0	6.3	3.0	
6.0	6.0	6.3	5	1000 m Downstream of D
6.0	6.0	6.3	0.5	
6.0	6.0	6.3	1.0	
6.0	6.0	6.3	1.5	
6.0	6.0	6.3	2.0	
6.0	6.0	6.3	2.5	
6.0	6.0	6.3	3.0	

Table 5.2-15

Thermal plume data for DMTO on 19 March 1976.

Station Operation Level (m) 0				Time: 11:5
Nuclear Service Pumps: 2				Intake Temp. (C): 14
Secondary Service Pumps: 2				Effluent Temp. (C): 4.7
Recycle Heat Pumps: 1				Air Temp. (C): 14.5
Effluent Rate (cfs): 40.10				Wind Speed (mph): 19
River Flow Rate (cfs): 10				Wind Dir.: 20
Distance from Three Mile Island Core				
50 m	20 m	5 m	Depth	
3.3	3.3	3.4	5	Unit 1 Intake
3.3	3.3	3.4	0.5 m	
3.3	3.3	3.4	1.0	
3.3	3.3	3.4	1.5	
3.3	3.3	3.4	2.0	
3.3	3.3	3.4	2.5	
3.3	3.4	3.6	5	25 m Upstream of Discharge
3.3	3.4	3.6	0.5	
3.3	3.4	3.6	1.0	
3.3	3.4	3.6	1.5	
3.3	3.4	3.6	2.0	
3.4	3.5	4.7	5	Discharge (D)
3.4	3.5	4.7	0.5	
3.4	3.5	4.7	1.0	
3.4	3.5	4.7	1.5	
3.4	3.5	4.7	2.0	
3.4	4.0	4.2	5	25 m Downstream of D
3.4	4.0	4.2	0.5	
3.4	4.0	4.2	1.0	
3.4	3.8	4.2	1.5	
3.4	3.8	4.2	2.0	
3.4	3.8	4.2	5	50 m Downstream of D
3.4	3.8	4.2	0.5	
3.4	3.8	4.2	1.0	
3.4	3.8	4.2	1.5	
3.4	3.8	4.2	2.0	
3.4	3.8	4.1	5	100 m Downstream of D
3.4	3.8	4.1	0.5	
3.4	3.8	4.1	1.0	
3.4	3.8	4.1	1.5	
3.4	3.8	4.1	2.0	
3.5	3.8	4.1	5	200 m Downstream of D
3.5	3.8	4.1	0.5	
3.4	3.8	4.0	1.0	
3.4	3.8	4.0	1.5	
3.4	3.8	4.0	2.0	
3.4	3.8	3.9	5	400 m Downstream of D
3.4	3.8	3.9	0.5	
3.4	3.8	3.9	1.0	
3.4	3.8	3.9	1.5	
3.4	3.8	3.9	2.0	
3.5	3.8	3.9	5	1000 m Downstream of D
3.5	3.8	3.9	0.5	
3.5	3.8	4.0	1.0	
3.5	3.8	4.0	1.5	
3.5	3.8	4.0	2.0	
3.5	3.7	3.9	2.5	
3.5	3.7	3.9	3.0	

POOR ORIGINAL

1565 232

Table 5.2-16

Thermal plume data for TMINS on 1 April 1976.

Station Operation Level (%): 0				Time: 1350
Nuclear Service Pumps: 2				Intake Temp. (C): 9.22
Secondary Service Pumps: 0				Effluent Temp. (C): 8.44
Decay Heat Pumps: 1				Air Temp. (C): NA
Effluent Rate (cfs): 33.57				Wind Speed (mph): 10
River Flow (cfs): 28,600				Wind Dir: 4
Distance from Three Mile Island shore				
40 m	20 m	5 m	Depth	
9.7	9.7	9.7	S	Unit 1 Intake
9.7	9.7	9.7	0.5 m	
9.7	9.7	9.7	1.0	
9.7	9.7	9.7	1.5	
9.7	9.7		2.0	
9.7	9.7		2.5	
	9.7		3.0	
9.7	9.8	9.8	S	25 m Upstream of Discharge
9.7	9.8	9.8	0.5	
9.7	9.8		1.0	
9.7	9.7		1.5	
9.7			2.0	
9.8	9.7	9.1	S	Discharge (D)
9.8	9.8	9.1	0.5	
9.8	9.8	9.1	1.0	Depth at Discharge
9.8	9.8	9.1	1.5	Pipe = 1.75 m
9.8		9.1	2.0	
9.8	9.7	9.4	S	25 m Downstream of D
9.8	9.7	9.4	0.5	
9.8	9.7		1.0	
9.8	9.7		1.5	
9.8			2.0	
9.8	9.7	9.5	S	50 m Downstream of D
9.8	9.7		0.5	
9.8	9.8		1.0	
9.8	9.8		1.5	
9.8	9.7		2.0	
9.8			2.5	
9.8	9.7	9.6	S	100 m Downstream of D
9.8	9.7	9.5	0.5	
9.8	9.7		1.0	
9.8	9.7		1.5	
9.8	9.7		2.0	
9.8			2.5	
9.8	9.7	9.6	S	200 m Downstream of D
9.8	9.8	9.6	0.5	
9.8	9.7	9.6	1.0	
9.8	9.7		1.5	
9.8	9.7		2.0	
9.7	9.7	9.7	S	400 m Downstream of D
9.7	9.7	9.7	0.5	
9.7	9.7		1.0	
9.7	9.7		1.5	
9.7			2.0	
9.7			2.5	
9.7	9.7	9.8	S	1900 m Downstream of D
9.6	9.7	9.8	0.5	
9.7	9.7	9.8	1.0	
9.7	9.7		1.5	
9.7	9.7		2.0	
9.7	9.7		2.5	

Table 5.2-17

Thermal plume data for TMINS on 15 April 1976.

Station Operation Level (%): 0				Time: 1335
Nuclear Service Pumps: 2				Intake Temp. (C): 11.4
Secondary Service Pumps: 1				Effluent Temp. (C): 11.5
Decay Heat Pumps: 0				Air Temp. (C): 21.0
Effluent Rate (cfs): 24.51				Wind Speed (mph): 10
River Flow (cfs): 28,700				Wind Dir: 8
Distance from Three Mile Island shore				
40 m	20 m	5 m	Depth	
11.8	11.8	11.8	S	Unit 1 Intake
11.8	11.8	11.8	0.5 m	
11.8	11.8	11.8	1.0	
11.8	11.8	11.8	1.5	
11.8	11.8	11.8	2.0	
11.8		11.8	2.5	
		11.8	3.0	
11.9	11.8	11.9	S	25 m Upstream of Discharge
11.9	11.8	11.9	0.5	
11.8	11.8	11.9	1.0	
11.8	11.8		1.5	
11.8	11.8		2.0	
11.9	11.9	11.8	S	Discharge (D)
11.9	11.9	11.8	0.5	
11.9	11.9	11.8	1.0	
11.9	11.9	11.7	1.5	
11.8			2.0	
11.9	11.9	11.7	S	25 m Downstream of D
11.9	11.9	11.7	0.5	
11.9	11.9	11.7	1.0	
11.9	11.9		1.5	
11.9	11.9		2.0	
11.9	11.9	11.8	S	50 m Downstream of D
11.9	11.9	11.8	0.5	
11.9	11.9	11.7	1.0	
11.9	11.9		1.5	
11.9	11.9		2.0	
11.9	11.9	11.9	S	100 m Downstream of D
11.9	11.9	11.9	0.5	
11.9	11.9	11.8	1.0	
11.9	11.9		1.5	
11.9	11.9		2.0	
11.7	11.7	11.8	S	1900 m Downstream of D
11.7	11.7	11.8	0.5	
11.7	11.7	11.8	1.0	
11.7	11.7	11.8	1.5	
11.7	11.7		2.0	
11.7			2.5	
11.7			3.0	

POOR ORIGINAL

1565 233

Table 5.2-18

Thermal plume data for THINS on 19 April 1976.

Station Operation Level: 0				Time: 1440
Nuclear Service Pumps: 2				Intake Temp. (C): 12.3
Secondary Service Pumps: 1				Effluent Temp. (C): 12.2
Decay Heat Pumps: 1				Air Temp. (C): 18.0
Effluent Rate (cfs): 44.56				Wind Speed (mph): 10
River Flow (cfs): 20,000				Wind Dir.: 090
Distance from Three Mile Island Shore				
40 m	20 m	5 m	Depth	
12.6	12.7	12.8	S	Unit 1 Intake
12.6	12.7	12.8	0.5 m	
12.6	12.7	12.8	1.0	
12.6	12.7	12.8	1.5	
12.6	12.7	12.8	2.0	
12.6	12.7	12.8	2.5	
12.6	12.7	12.8	3.0	
12.7	12.7	13.4	S	25 m Upstream of Discharge
12.7	12.7	13.4	0.5	
12.7	12.7	13.4	1.0	
12.7	12.7	13.4	1.5	
12.7	12.7	13.4	2.0	
12.7	12.8	12.2	S	Discharge (D)
12.7	12.8	12.2	0.5	
12.7	12.8	12.2	1.0	
12.7	12.7	12.2	1.5	
12.7	12.7	12.2	2.0	
12.7	12.8	12.4	S	25 m Downstream of D
12.7	12.7	12.4	0.5	
12.7	12.7	12.4	1.0	
12.7	12.7	12.4	1.5	
12.7	12.7	12.4	2.0	
12.7	12.8	12.5	S	50 m Downstream of D
12.7	12.8	12.5	0.5	
12.7	12.8	12.5	1.0	
12.7	12.7	12.5	1.5	
12.7	12.7	12.5	2.0	
12.7	12.8	12.6	S	100 m Downstream of D
12.7	12.8	12.6	0.5	
12.7	12.7	12.6	1.0	
12.7	12.7	12.6	1.5	
12.7	12.7	12.6	2.0	
12.7	12.8	12.7	S	200 m Downstream of D
12.7	12.8	12.7	0.5	
12.7	12.8	12.7	1.0	
12.7	12.8	12.7	1.5	
12.7	12.7	12.7	2.0	
12.7	12.8	12.8	S	400 m Downstream of D
12.7	12.8	12.9	0.5	
12.7	12.8	12.9	1.0	
12.7	12.8	12.9	1.5	
12.7	12.8	12.9	2.0	
12.7	12.8	12.9	2.5	

Table 5.2-19

Thermal plume data for THINS on 13 May 1976.

Station Operation Level: 0				Time: 1300
Nuclear Service Pumps: 2				Intake Temp. (C): 17.1
Secondary Service Pumps: 1				Effluent Temp. (C): 17.1
Decay Heat Pumps: 2				Air Temp. (C): 19.8
Effluent Rate (cfs): 77.97				Wind Speed (mph): 5
River Flow (cfs): 20,000				Wind Dir.: 0
Distance from Three Mile Island Shore				
40 m	20 m	5 m	Depth	
17.0	17.2	17.5	S	Unit 1 Intake
17.0	17.2	17.5	0.5 m	
17.0	17.1	17.5	1.0	
17.0	17.1	17.5	1.5	
17.0	17.1	17.4	2.0	
		17.3	2.5	
		17.4	3.0	
17.4	17.5	17.9	S	25 m Upstream of Discharge
17.4	17.4	17.7	0.5	
17.4	17.4	17.6	1.0	
17.4	17.4	17.6	1.5	
17.3	17.4	17.6	2.0	
17.4	17.5	17.2	S	Discharge (D)
17.3	17.5	17.2	0.5	
17.2	17.5	17.2	1.0	
17.2	17.5	17.2	1.5	
17.2	17.5	17.2	2.0	
17.4	17.4	17.4	S	25 m Downstream of D
17.4	17.4	17.4	0.5	
17.4	17.4	17.4	1.0	
17.4	17.4	17.4	1.5	
17.4	17.4	17.4	2.0	
17.4	17.4	17.4	S	50 m Downstream of D
17.4	17.4	17.4	0.5	
17.4	17.4	17.3	1.0	
17.4	17.4	17.3	1.5	
17.4	17.4	17.3	2.0	
17.4	17.4	17.4	S	100 m Downstream of D
17.4	17.4	17.4	0.5	
17.3	17.4	17.4	1.0	
17.3	17.4	17.4	1.5	
	17.4	17.4	2.0	
17.4	17.4	17.4	S	200 m Downstream of D
17.4	17.4	17.4	0.5	
17.4	17.4	17.4	1.0	
17.4	17.4	17.4	1.5	
17.4	17.4	17.4	2.0	
17.4	17.5	17.6	S	400 m Downstream of D
17.4	17.5	17.6	0.5	
17.4	17.5	17.6	1.0	
17.3	17.4	17.6	1.5	
17.3	17.4	17.6	2.0	
17.2	17.4	17.4	S	1900 m Downstream of D
17.2	17.4	17.4	0.5	
17.1	17.4	17.4	1.0	
17.0	17.3	17.3	1.5	
17.0	17.1	17.3	2.0	

POOR ORIGINAL

Table 5.2-20

Thermal Plume data for EMMS on 27 May 1976.

Station Operation Level (ft): 40				Time: 1310
Nuclear Service Pumps: 2				Intake Temp. (C): 15.7
Secondary Service Pumps: 1				Effluent Temp. (C): 15.0
Decay Heat Pumps: 1				Air Temp. (C): 21.0
Effluent Rate (cfs): 93.57				Wind Speed (mph): 4
River Flow (cfs): 35,000				Wind Dir: 100
Distance From Three Mile Island Shore				
40 m	20 m	5 m	Depth	
15.1	15.3	15.3	S	Unit 1 Intake
15.1	15.3	15.3	0.5 m	
15.1	15.2	15.3	1.0	
15.1	15.2	15.3	1.5	
15.1	15.2	15.2	2.0	
15.2	15.3	15.7	S	25 m Upstream of Discharge
15.2	15.3	15.6	0.5	
15.2	15.3		1.0	
15.2			1.5	
15.2			2.0	
15.4	15.5	15.4	S	Discharge (D)
15.4	15.4	15.4	0.5	
15.3	15.4	15.4	1.0	
15.3	15.4	15.4	1.5	
15.3			2.0	
15.4	15.5	15.5	S	25 m Downstream of D
15.4	15.5	15.5	0.5	
15.4	15.5	15.5	1.0	
15.4	15.5		1.5	
15.4			2.0	
15.3	15.4	15.6	S	50 m Downstream of D
15.3	15.4	15.5	0.5	
15.3	15.4	15.3	1.0	
15.3	15.4		1.5	
15.3	15.4		2.0	
15.3	15.5	15.6	S	100 m Downstream of D
15.3	15.5	15.5	0.5	
15.3	15.5	15.5	1.0	
15.3	15.5	15.5	1.5	
15.3	15.4		2.0	
15.4	15.5	15.8	S	200 m Downstream of D
15.4	15.5	15.7	0.5	
15.4	15.5		1.0	
15.4	15.5		1.5	
15.4	15.5		2.0	
15.3	15.6	15.8	S	400 m Downstream of D
15.3	15.6	15.8	0.5	
15.3	15.5		1.0	
15.3	15.5		1.5	
15.3	15.5		2.0	
15.3			2.5	
15.4	15.7	15.9	S	1900 m Downstream of D
15.4	15.7	15.8	0.5	
15.4	15.6		1.0	
15.3	15.5		1.5	
15.3	15.4		2.0	
15.3	15.3		2.5	

Table 5.2-21

Thermal plume data for EMMS on 2 June 1976.

Station Operation Level (ft): 40				Time: 1320
Nuclear Service Pumps: 1				Intake Temp. (C): 20.56
Secondary Service Pumps: 2				Effluent Temp. (C): 17.76
Decay Heat Pumps: 0				Air Temp. (C): 16.0
Effluent Rate (cfs): 40.10				Wind Speed (mph): calm
River Flow (cfs): 47,700				Wind Dir: 100
Distance From Three Mile Island Shore				
40 m	20 m	5 m	Depth	
19.0	18.9	18.9	S	Unit 1 Intake
19.0	18.9	18.9	0.5 m	
19.0	18.9	18.9	1.0	
19.0	18.9	18.9	1.5	
19.0	18.9	18.9	2.0	
19.0	18.9	19.0	2.5	
19.0		19.0	3.0	
18.9	18.9	18.9	S	25 m Upstream of Discharge
18.9	18.9	18.9	0.5	
18.9	18.9	18.9	1.0	
18.9	18.9		1.5	
18.9	18.9		2.0	
18.9			2.5	
18.9			3.0	
18.9	18.9	18.3	S	Discharge (D)
18.9	18.9	18.3	0.5	
18.9	18.9	18.3	1.0	
18.9	18.9		1.5	
18.9	18.9		2.0	
18.9	18.9	18.4	S	25 m Downstream of D
18.9	18.9	18.4	0.5	
18.9	18.9	18.4	1.0	
18.9	18.9		1.5	
18.9	18.9		2.0	
18.9	18.9	18.5	S	50 m Downstream of D
18.9	18.9	18.5	0.5	
18.9	18.9	18.5	1.0	
18.9	18.9		1.5	
18.9	18.9		2.0	
18.9	18.9	18.6	S	100 m Downstream of D
18.9	18.9	18.6	0.5	
18.9	18.9	18.6	1.0	
18.9	18.9		1.5	
18.9	18.9		2.0	
18.9	18.9	18.6	S	200 m Downstream of D
18.9	18.9	18.6	0.5	
18.9	18.9	18.6	1.0	
18.9	18.9		1.5	
18.9	18.9		2.0	
18.9	18.9	18.7	S	400 m Downstream of D
18.9	18.9	18.7	0.5	
18.9	18.9		1.0	
18.9	18.9		1.5	
18.9	18.9		2.0	
18.9			2.5	
18.9			3.0	
19.0	19.0	18.9	S	1900 m Downstream of D
19.0	19.0	18.9	0.5	
19.0	19.0	18.9	1.0	
19.0	19.0		1.5	
19.0	19.0		2.0	
19.0	18.9		2.5	

POOR ORIGINAL

187

Table 5.2-22

Thermal plume data for TMS on 10 June 1976.

Station Operation Level (ft)	Time	Intake Temp. (°C)	Effluent Temp. (°C)	Air Temp. (°C)	Wind Speed (mph)
Nuclear Service Pumps: 2	1300	16.0	21.67	30.0	0
Secondary Service Pumps: 2					
Leak Heat Pumps: 1					
Effluent Rate (cfs): 40.10					
River flow (cfs): 1000					
Distance from Three Mile Island (ft)					
40 m	25 m	5 m	fe, th		
25.7	25.8	25.6	5	Unit 1 Intake	
25.7	25.8	25.8	0.5 m		
25.7	25.8	25.8	1.0		
25.7	25.7	25.8	1.5		
25.7	25.7	25.8	2.0		
25.8	25.8	26.2	5	25 m Upstream of Discharge	
25.8	25.8	26.2	0.5		
25.8	25.8		1.0		
25.8	25.8		1.5		
25.7			2.0		
25.8	25.7	25.5	5	Discharge (D)	
25.8	25.7	25.4	0.5		
25.8	25.8	25.3	1.0		
25.8	25.8		1.5		
25.8			2.0		
25.8	25.9	25.6	5	25 m Downstream of D	
25.8	25.8	25.5	0.5		
25.8	25.8	25.5	1.0		
25.8	25.8		1.5		
25.8			2.0		
25.8	25.9	25.7	5	50 m Downstream of D	
25.8	25.9		0.5		
25.8	25.9		1.0		
25.8	25.8		1.5		
25.8	25.8		2.0		
25.9	25.9	25.9	5	100 m Downstream of D	
25.9	25.9	25.9	0.5		
25.8	25.9		1.0		
25.8	25.8		1.5		
25.8	25.8		2.0		
25.9	25.9	25.9	5	200 m Downstream of D	
25.9	25.9	25.9	0.5		
25.9	25.9	25.9	1.0		
25.9	25.9		1.5		
25.9	25.9		2.0		
25.9	25.9	26.2	5	400 m Downstream of D	
25.9	25.9	26.1	0.5		
25.9	25.9	26.1	1.0		
25.9	25.9		1.5		
25.9	25.9		2.0		
25.9			2.5		
26.4	26.4	26.1	5	1900 m Downstream of D	
26.0	26.3	26.0	0.5		
25.9	26.1	25.9	1.0		
25.8	25.9	25.8	1.5		
25.8	26.3		2.0		
23.2			2.5		

Table 5.2-23

Thermal plume data for TMS on 14 June 1976.

Station Operation Level (ft)	Time	Intake Temp. (°C)	Effluent Temp. (°C)	Air Temp. (°C)	Wind Speed (mph)
Nuclear Service Pumps: 2	1300	16.0	21.67	30.0	0
Secondary Service Pumps: 2					
Leak Heat Pumps: 1					
Effluent Rate (cfs): 40.10					
River flow (cfs): 1000					
Distance from Three Mile Island (ft)					
40 m	25 m	5 m	fe, th		
25.7	25.8	25.6	5	Unit 1 Intake	
25.7	25.8	25.8	0.5 m		
25.7	25.8	25.8	1.0		
25.7	25.7	25.8	1.5		
25.7	25.7	25.8	2.0		
25.8	25.8	26.2	5	25 m Upstream of Discharge	
25.8	25.8	26.2	0.5		
25.8	25.8		1.0		
25.8	25.8		1.5		
25.7			2.0		
25.8	25.7	25.5	5	Discharge (D)	
25.8	25.7	25.4	0.5		
25.8	25.8	25.3	1.0		
25.8	25.8		1.5		
25.8			2.0		
25.8	25.9	25.6	5	25 m Downstream of D	
25.8	25.8	25.5	0.5		
25.8	25.8	25.5	1.0		
25.8	25.8		1.5		
25.8			2.0		
25.8	25.9	25.7	5	50 m Downstream of D	
25.8	25.9		0.5		
25.8	25.9		1.0		
25.8	25.8		1.5		
25.8	25.8		2.0		
25.9	25.9	25.9	5	100 m Downstream of D	
25.9	25.9	25.9	0.5		
25.8	25.9		1.0		
25.8	25.8		1.5		
25.8	25.8		2.0		
25.9	25.9	25.9	5	200 m Downstream of D	
25.9	25.9	25.9	0.5		
25.9	25.9	25.9	1.0		
25.9	25.9		1.5		
25.9	25.9		2.0		
25.9	25.9	26.2	5	400 m Downstream of D	
25.9	25.9	26.1	0.5		
25.9	25.9	26.1	1.0		
25.9	25.9		1.5		
25.9	25.9		2.0		
25.9			2.5		
26.4	26.4	26.1	5	1900 m Downstream of D	
26.0	26.3	26.0	0.5		
25.9	26.1	25.9	1.0		
25.8	25.9	25.8	1.5		
25.8	26.3		2.0		
23.2			2.5		

1565 236

POOR ORIGINAL

Table 5.2-24

Thermal plume data for TMINS on 15 July 1976.

Station Operation Level (ft): 100			Time: 0930	
Nuclear Service Pumps: 3			Intake Temp. (C): 22.78	
Secondary Service Pumps: 2			Effluent Temp. (C): 23.33	
Decay Heat Pumps: 1			Air Temp. (C): 24.5	
Effluent Rate (cfs): 40.10			Wind Speed (mph): 2-3	
River Flow (cfs): 19,200			Wind Dir.: N	
Distance From Three Mile Island Shore				
40 m	20 m	5 m	Depth	
22.0	21.9	21.8	S	Unit 1 Intake
22.0	21.9	21.8	0.5 m	
22.0	21.9	21.8	1.0	
22.0	21.9	21.9	1.5	
22.0	21.9	21.8	2.0	
21.9	21.9	21.8	S	25 m Upstream of Discharge
21.9	21.9	21.8	0.5	
21.9	21.8	21.8	1.0	
21.9	21.8		1.5	
21.9	21.8		2.0	
21.9			2.5	
21.9			3.0	
21.9	21.7	22.3	S	Discharge (D)
21.9	21.8	22.2	0.5	
21.9	21.7	22.2	1.0	
21.9	21.8		1.5	
21.9			2.0	
21.9	22.0	22.2	S	25 m Downstream of D
21.9	22.0	22.3	0.5	
21.9	22.0		1.0	
21.9	22.0		1.5	
21.9	22.0		2.0	
22.1	22.0	22.2	S	50 m Downstream of D
22.1	22.0	22.2	0.5	
22.1	22.0	22.1	1.0	
22.0	22.0		1.5	
22.0	21.9		2.0	
22.1	22.0	22.2	S	100 m Downstream of D
22.1	22.0	22.2	0.5	
22.1	22.0		1.0	
22.1	22.0		1.5	
22.0	22.0		2.0	
22.1	22.1	22.1	S	200 m Downstream of D
22.1	22.0	22.1	0.5	
22.1	22.0	22.0	1.0	
22.1	22.0	21.0	1.5	
22.1	22.0		2.0	
22.1	22.0	22.0	S	400 m Downstream of D
22.1	22.0	22.0	0.5	
22.1	22.0	22.0	1.0	
22.1	22.0		1.5	
22.1	22.0		2.0	
22.1			2.5	
21.9	21.9	21.8	S	1000 m Downstream of D
21.9	21.9	21.8	0.5	
21.9	21.9	21.8	1.0	
21.9	21.9	21.8	1.5	
21.8	21.8	21.7	2.0	
21.8	21.7		2.5	
21.7			3.0	

Table 5.2-25

Thermal plume data for TMINS on 30 July 1976.

Station Operation Level (ft): 100			Time: 1330	
Nuclear Service Pumps: 3			Intake Temp. (C): 25.0	
Secondary Service Pumps: 2			Effluent Temp. (C): 25.0	
Decay Heat Pumps: 0			Air Temp. (C): 27.0	
Effluent Rate (cfs): 44.66			Wind Speed (mph): 5-7	
River Flow (cfs): 13,200			Wind Dir.: S	
Distance From Three Mile Island Shore				
40 m	20 m	5 m	Depth	
23.0	23.6	23.8	S	Unit 1 Intake
22.9	23.5	23.9	0.5 m	
22.9	23.5	23.9	1.0	
22.9	23.5	23.9	1.5	
22.9		23.8	2.0	
		23.7	2.5	
		23.7	3.0	
23.2	23.8	24.0	S	25 m Upstream of Discharge
23.2	23.8	24.0	0.5	
23.1	23.8	24.0	1.0	
23.1	23.8		1.5	
23.1			2.0	
23.5	23.8	24.4	S	Discharge (D)
23.5	23.9	24.4	0.5	
23.5	23.9		1.0	Depth at Discharge
23.4	23.9		1.5	Pipe = 0.75 m
23.4			2.0	
23.2	23.7	24.2	S	25 m Downstream of D
23.2	23.8	24.2	0.5	
23.2	23.8		1.0	
23.1	23.8		1.5	
23.5	24.1	24.1	S	50 m Downstream of D
23.5	24.1	24.1	0.5	
23.5	24.1		1.0	
23.5	24.1		1.5	
23.3	24.0	24.1	S	100 m Downstream of D
23.3	24.0	24.1	0.5	
23.3	24.0	24.1	1.0	
23.3	23.9		1.5	
23.5	23.9	24.2	S	200 m Downstream of D
23.5	23.9	24.2	0.5	
23.5	23.9	24.2	1.0	
23.4	23.9		1.5	
23.4	23.9	24.2	S	400 m Downstream of D
23.3	23.9	24.2	0.5	
23.3	23.9	24.2	1.0	
23.2	23.9		1.5	
23.2	23.9		2.0	
23.9	24.1	24.1	S	1000 m Downstream of D
23.8	23.9	23.9	0.5	
23.7	23.8	23.8	1.0	
23.7	23.8	23.8	1.5	
23.6	23.7		2.0	

1565 237

Table 5.2-26

Thermal plume data for TMIN on 12 August 1976.

Station Operation Level (°): 100			Time: 1450
Nuclear Service Pumps: 3			Intake Temp. (°C): 25.45
Secondary Service Pumps: 2			Effluent Temp. (°C): 23.3
Decay Heat Pumps: 0			Air Temp. (°C): 26.5
Effluent Rate (cfs): 26.73			Wind Speed (mph): 5
River Flow (cfs): 19,200			Wind Dir.: N
Distance from Three Mile Island Shore			
40 m	20 m	5 m	Depth
24.6	24.7	24.8	S Unit 1 Intake
24.6	24.7	24.8	0.5 m
24.6	24.7	24.8	1.0
24.6	24.7	24.8	1.5
24.6	24.7	24.8	2.0
		24.8	2.5
		24.8	3.0
24.7	24.8	25.1	S 25 m Upstream of Discharge
24.7	24.8	25.1	0.5
24.7	24.8		1.0
24.7	24.8		1.5
24.7	24.8		2.0
24.8	24.9	24.7	S Discharge (D)
24.8	24.9	24.7	0.5
24.8	24.9	24.7	1.0
24.8	24.9	24.7	1.5
24.8	24.9	24.8	S 25 m Downstream of D
24.8	24.9	24.8	0.5
24.8	24.9		1.0
24.8	24.9		1.5
24.8	24.9		2.0
24.8	24.9	24.9	S 50 m Downstream of D
24.8	24.9		0.5
24.8	24.9		1.0
24.8	24.9		1.5
24.8	24.9		2.0
24.8	24.8	24.8	S 100 m Downstream of D
24.8	24.8	24.8	0.5
24.7	24.8	24.8	1.0
24.7	24.8		1.5
24.7	24.8		2.0
24.8	24.9	24.9	S 200 m Downstream of D
24.8	24.9	24.9	0.5
24.8	24.9	24.9	1.0
24.8	24.8		1.5
24.8	24.8		2.0
24.9	24.9	25.1	S 400 m Downstream of D
24.9	24.9	25.1	0.5
24.9	24.9		1.0
24.8	24.9		1.5
24.8	24.8		2.0
24.9	24.9	25.2	S 1900 m Downstream of D
24.9	24.9	25.2	0.5
24.9	24.9	25.1	1.0
24.8	24.8	25.1	1.5
24.8	24.8		2.0
24.6			2.5

Table 5.2-27

Thermal plume data for TMIN on 16 August 1976.

Station Operation Level (°): 100			Time: 1325
Nuclear Service Pumps: 3			Intake Temp. (°C): 24.67
Secondary Service Pumps: 2			Effluent Temp. (°C): 25.83
Decay Heat Pumps: 0			Air Temp. (°C): 24.0
Effluent Rate (cfs): 53.47			Wind Speed (mph): 4
River Flow (cfs): 11,600			Wind Dir.: N
Distance from Three Mile Island Shore			
40 m	20 m	5 m	Depth
26.6	26.4	26.2	S Unit 1 Intake
26.6	26.4	26.2	0.5 m
26.6	26.4	26.2	1.0
26.6	26.4	26.1	1.5
	26.3	26.1	2.0
	26.3	26.1	2.5
26.4	26.2	27.1	S 25 m Upstream of Discharge
26.4	26.2	26.9	0.5
26.4	26.1		1.0
26.4	26.1		1.5
26.4	26.1		2.0
26.3	26.3	26.0	S Discharge (D)
26.3	26.3	26.0	0.5
26.2	26.3	26.0	1.0
26.2	26.3	26.0	1.5
26.2	26.2	26.9	S 25 m Downstream of D
26.2	26.2	26.9	0.5
26.2	26.2		1.0
26.2	26.2		1.5
26.6	26.2	26.3	S 50 m Downstream of D
26.6	26.2	26.3	0.5
26.5	26.2		1.0
26.4	26.2		1.5
26.3	26.3	26.4	S 100 m Downstream of D
26.3	26.2	26.4	0.5
26.3	26.2		1.0
26.3	26.2		1.5
26.4	26.3	26.3	S 200 m Downstream of D
26.4	26.3	26.3	0.5
26.4	26.3	26.3	1.0
26.4	26.3		1.5
26.3	26.3	26.3	S 400 m Downstream of D
26.3	26.3	26.3	0.5
26.3	26.3		1.0
26.3	26.3		1.5
27.0	27.4	28.0	S 1900 m Downstream of D
27.0	27.3	27.7	0.5
26.9	27.1	27.3	1.0
26.9	27.0	27.0	1.5
26.4	26.5		2.0

POOR ORIGINAL

1565 238

Table 5.2-28

Thermal plume data for TMIN on 9 September 1976.

Station Operation Level (C): 100			Time: 1310
Nuclear Service Pumps: 3			Intake Temp. (C): 23.72
Secondary Service Pumps: 2			Effluent Temp. (C): 22.50
Decay Heat Pumps: 1			Air Temp. (C): 27.0
Effluent Rate (cfs): 62.38			Wind Speed (mph): 4
River Flow (cfs): 4300			Wind Dir.: E
Distance From Three Mile Island Shore			
40 m	20 m	5 m	Depth
23.0	22.7	22.7	S Unit 1 Intake
23.0	22.7	22.7	0.5 m
23.0	22.7	22.7	1.0
23.0	22.7	22.7	1.5
22.9	22.6	22.7	2.0
		22.6	2.5
		22.6	3.0
23.0	23.2	23.0	S 25 m Upstream of Discharge
22.9	23.0	23.1	0.5
22.9	22.9		1.0
22.8	22.8		1.5
22.7			2.0
23.1	23.0	22.8	S Discharge (D)
23.0	23.0	22.9	0.5
22.9	23.0	23.0	1.0
22.8			1.5
22.9	23.0	23.0	S 25 m Downstream of D
22.9	23.0	22.9	0.5
22.9	23.0		1.0
22.8	23.0		1.5
23.0	23.0	23.0	S 50 m Downstream of D
23.0	23.0	23.0	0.5
23.0	23.0		1.0
23.0	23.0		1.5
23.0	23.0	23.1	S 100 m Downstream of D
22.9	23.0	23.1	0.5
22.9	23.0		1.0
22.9	23.0		1.5
23.0	23.2	23.4	S 200 m Downstream of D
23.0	23.1	23.3	0.5
23.0	23.1		1.0
22.9	23.0		1.5
23.1	23.2	23.3	S 400 m Downstream of D
23.1	23.1	23.4	0.5
23.1	23.1		1.0
23.1	23.1		1.5
23.0			2.0
23.0			2.5
24.0	24.4	24.9	S 1900 m Downstream of D
23.7	23.3	24.3	0.5
23.4	23.1	23.0	1.0
23.1	23.0		1.5
22.9	22.5		2.0

Table 5.2-29

Thermal plume data for TMIN on 22 September 1976.

Station Operation Level (C): 100			Time: 1315	
Nuclear Service Pumps: 3			Intake Temp. (C): 18.5	
Secondary Service Pumps: 2			Effluent Temp. (C): 18.06	
Decay Heat Pumps: 0			Air Temp. (C): 19.0	
Effluent Rate (cfs): 44.56			Wind Speed (mph): 10	
River Flow (cfs): 14,500			Wind Dir.: E	
Distance from Three Mile Island Shore				
40 m	20 m	5 m	Depth	
18.5	18.3	18.3	S	Unit 1 Intake
18.5	18.3	18.3	0.5 m	
18.5	18.3	18.3	1.0	
18.5	18.3	18.3	1.5	
18.5	18.3	18.3	2.0	
		18.3	2.5	
		18.3	3.0	
			4	
18.6	18.5	18.6	S	25 m Upstream of Discharge
18.6	18.4	18.7	0.5	
18.7	18.4		1.0	
18.7	18.4		1.5	
18.7	18.4		2.0	
18.7			2.5	
18.3	18.4	17.4	S	Discharge (D)
18.3	18.3	17.4	0.5	
18.3	18.3	17.5	1.0	Depth at Discharge
18.3			1.5	Pipe = 1.0 m
18.2	18.0	18.1	S	25 m Downstream of D
18.2	18.0	18.1	0.5	
18.2	18.0		1.0	
18.3	18.0		1.5	
18.3	18.3	17.9	S	50 m Downstream of D
18.4	18.3	18.0	0.5	
18.4	18.3		1.0	
18.4	18.3		1.5	
18.4	18.3	18.2	S	100 m Downstream of D
18.5	18.4	18.2	0.5	
18.5	18.4		1.0	
18.5	18.4		1.5	
18.4	18.3	18.1	S	200 m Downstream of D
18.4	18.3	18.2	0.5	
18.4	18.3	18.2	1.0	
18.3	18.2		1.5	
18.3	18.4	18.4	S	400 m Downstream of D
18.3	18.4	18.4	0.5	
18.3	18.4		1.0	
18.3	18.4		1.5	
18.3	18.3		2.0	
18.4			2.5	
18.2	18.2	18.5	S	1900 m Downstream of D
18.3	18.3	18.5	0.5	
18.3	18.3	18.5	1.0	
18.3	18.3		1.5	
18.4			2.0	

POOR ORIGINAL

1565 239

POOR ORIGINAL

Table 5.2-30

Thermal plume data for TMINS on 7 October 1976.

Station Operation Level (C): 100				Time: 1330
Nuclear Service Pumps: 2				Intake Temp. (C): 16.19
Secondary Service Pumps: 2				Affluent Temp. (C): 16.78
Decay Heat Pumps: 0				Air Temp. (C): 19.0
Effluent Rate (cfs): 44.56				Wind Speed (mph): 2
River Flow (cfs): 23,100				Wind Dir.: N
Distance from Three Mile Island shore				
40 m	20 m	5 m	Depth	
17.2	17.0	16.5	S	Unit 1 Intake
17.2	17.0	16.5	0.5 m	
17.2	17.0	16.5	1.0	
17.2	17.0	16.5	1.5	
17.2	17.0	16.5	2.0	
17.3	17.0	16.5	2.5	
	17.0	16.5	3.0	
17.1	16.7	16.7	S	25 m Upstream of Discharge
17.1	16.7	16.7	0.5	
17.1	16.7		1.0	
17.1	16.6		1.5	
17.1	16.6		2.0	
17.1			2.5	
17.1			3.0	
17.0	16.6	17.3	S	Discharge (D)
16.9	16.6	17.3	0.5	
16.9	16.6	17.3	1.0	Depth at Discharge
16.9	16.6	17.2	1.5	Pipe = 1.5 m
16.9	16.6		2.0	
17.0	16.7	17.0	S	25 m Downstream of D
17.0	16.7	16.9	0.5	
17.0	16.7		1.0	
17.0	16.7		1.5	
17.1			2.0	
17.1	16.7	16.9	S	50 m Downstream of D
17.1	16.7	16.9	0.5	
17.1	16.7		1.0	
17.1	16.7		1.5	
17.1	16.7		2.0	
17.1	16.7	16.8	S	100 m Downstream of D
17.1	16.7	16.8	0.5	
17.1	16.7		1.0	
17.1	16.7		1.5	
17.1	16.7		2.0	
17.2	16.8	16.8	S	200 m Downstream of D
17.2	16.8	16.8	0.5	
17.2	16.8		1.0	
17.2	16.8		1.5	
17.3	16.8		2.0	
17.2	16.9	16.9	S	400 m Downstream of D
17.2	16.9	16.9	0.5	
17.2	16.9		1.0	
17.2	16.9		1.5	
17.2	16.9		2.0	
17.2	16.9		2.5	
17.0	16.8	16.9	S	1900 m Downstream of D
17.0	16.8	16.9	0.5	
17.0	16.8		1.0	
17.0	16.8		1.5	
17.0	16.8		2.0	
17.0	16.8		2.5	

Table 5.2-31

Thermal plume data for TMINS on 21 October 1976.

Station Operation Level (C): 50				Time: 1336
Nuclear Service Pumps: 2				Intake Temp. (C): 10.22
Secondary Service Pumps: 2				Affluent Temp. (C): 10.47
Decay Heat Pumps: 0				Air Temp. (C): 11.0
Effluent Rate (cfs): 66.84				Wind Speed (mph): 18
River Flow (cfs): 43,600				Wind Dir.: W
Distance from Three Mile Island shore				
40 m	20 m	5 m	Depth	
10.3	10.4	10.5	S	Unit 1 Intake
10.3	10.5	10.6	0.5 m	
10.3	10.5	10.6	1.0	
10.3	10.5	10.6	1.5	
10.3	10.5	10.6	2.0	
	10.5	10.5	2.5	
	10.5	10.5	3.0	
10.5	10.6	10.7	S	25 m Upstream of Discharge
10.5	10.6	10.6	0.5	
10.5	10.5		1.0	
10.4	10.5		1.5	
10.4	10.5		2.0	
10.5	10.5	10.2	S	Discharge (D)
10.5	10.6	10.2	0.5	
10.4	10.6	10.2	1.0	Depth at Discharge
10.4	10.6	10.2	1.5	Pipe = 1.5 m
10.4			2.0	
10.5	10.4	10.4	S	25 m Downstream of D
10.5	10.5	10.4	0.5	
10.5	10.5		1.0	
10.5	10.5		1.5	
10.5	10.4		2.0	
10.5	10.5	10.4	S	50 m Downstream of D
10.5	10.5	10.4	0.5	
10.5	10.5	10.4	1.0	
10.5	10.5		1.5	
10.5	10.5		2.0	
10.6	10.5	10.5	S	100 m Downstream of D
10.5	10.5	10.5	0.5	
10.5	10.5		1.0	
10.5	10.5		1.5	
10.5	10.5		2.0	
10.5	10.5	10.5	S	200 m Downstream of D
10.5	10.5	10.5	0.5	
10.5	10.5		1.0	
10.5	10.5		1.5	
10.5	10.5		2.0	
10.5			2.5	
10.5	10.6	10.6	S	400 m Downstream of D
10.5	10.5	10.6	0.5	
10.5	10.5		1.0	
10.5	10.5		1.5	
10.5			2.0	
10.4	10.5	10.5	S	1900 m Downstream of D
10.4	10.5	10.5	0.5	
10.4	10.5	10.5	1.0	
10.4	10.4		1.5	
10.4	10.4		2.0	
10.4			2.5	

1565 240

Table 5.2-32

Thermal plume data for TMI-2 on 4 November 1979.

Station Operation Level (C): 100 Time: 1327			
Nuclear Service Pumps: 2 Intake Temp (C): 5.72			
Secondary Service Pumps: 2 Effluent Temp (C): 6.39			
Decay Heat Pumps: 0 Air Temp (C): 9.0			
Effluent Rate (cfs): 49.01 Wind Speed (mph): 3			
River Flow (cfs): 22,200 Wind Dir: 75			
Distance from Three Mile Island Shore			
40 m	20 m	5 m	Depth
6.4	6.4	6.5	S Unit 1 Intake
6.4	6.4	6.5	0.5 m
6.4	6.4	6.5	1.0
6.4	6.4	6.5	1.5
6.4	6.4	6.5	2.0
6.4	6.4	6.5	2.5
6.4	6.4	6.5	3.0
6.4	6.5	6.5	S 25 m Upstream of Discharge
6.4	6.5	6.5	0.5
6.4	6.5	6.5	1.0
6.4	6.5	6.5	1.5
6.4	6.5	6.5	2.0
6.4	6.5	6.5	2.5
6.4	6.4	7.1	S Discharge (D)
6.4	6.4	7.1	0.5
6.4	6.4	7.1	1.0
6.4	6.4	7.1	1.5
6.4	6.4	7.1	2.0
6.4	6.4	7.1	2.5
6.4	6.5	6.9	S 25 m Downstream of D
6.4	6.5	6.9	0.5
6.4	6.5	6.9	1.0
6.4	6.5	6.9	1.5
6.4	6.5	6.9	2.0
6.4	6.5	6.9	2.5
6.4	6.4	6.8	S 50 m Downstream of D
6.4	6.4	6.8	0.5
6.4	6.4	6.8	1.0
6.4	6.4	6.8	1.5
6.4	6.4	6.8	2.0
6.4	6.4	6.8	2.5
6.4	6.5	6.8	S 100 m Downstream of D
6.4	6.5	6.8	0.5
6.4	6.5	6.8	1.0
6.4	6.5	6.8	1.5
6.4	6.5	6.8	2.0
6.4	6.5	6.8	2.5
6.4	6.5	6.8	S 200 m Downstream of D
6.4	6.5	6.8	0.5
6.4	6.5	6.8	1.0
6.4	6.5	6.8	1.5
6.4	6.5	6.8	2.0
6.4	6.5	6.8	2.5
6.4	6.5	6.7	S 400 m Downstream of D
6.4	6.5	6.7	0.5
6.4	6.5	6.7	1.0
6.4	6.5	6.7	1.5
6.4	6.5	6.7	2.0
6.4	6.5	6.7	2.5
6.4	6.5	6.8	S 600 m Downstream of D
6.4	6.5	6.8	0.5
6.4	6.5	6.8	1.0
6.4	6.5	6.8	1.5
6.4	6.5	6.8	2.0
6.4	6.5	6.8	2.5
6.4	6.5	6.8	S 1900 m Downstream of D
6.4	6.5	6.8	0.5
6.4	6.5	6.8	1.0
6.4	6.5	6.8	1.5
6.4	6.5	6.8	2.0
6.4	6.5	6.8	2.5

Table 5.2-33

Thermal plume data for TMI-2 on 18 November 1979.

Station Operation Level (C): 0 Time: 0850			
Nuclear Service Pumps: 1 Intake Temp (C): 2.53			
Secondary Service Pumps: 1 Effluent Temp (C): 2.44			
Decay Heat Pumps: 1 Air Temp (C): 1.5			
Effluent Rate (cfs): 57.92 Wind Speed (mph): 4			
River Flow (cfs): 22,600 Wind Dir: 75			
Distance from Three Mile Island Shore			
40 m	20 m	5 m	Depth
2.7	3.1	3.5	S Unit 1 Intake
2.6	3.1	3.5	0.5 m
2.6	3.1	3.5	1.0
2.6	3.1	3.5	1.5
2.6	3.2	3.5	2.0
2.6	3.2	3.5	2.5
2.6	3.1	3.5	3.0
2.7	3.4	3.5	S 25 m Upstream of Discharge
2.7	3.4	3.5	0.5
2.7	3.4	3.5	1.0
2.7	3.4	3.5	1.5
2.7	3.4	3.5	2.0
2.7	3.4	3.5	2.5
2.7	3.5	4.3	S Discharge (D)
2.7	3.5	4.3	0.5
2.7	3.4	4.3	1.0
2.7	3.4	4.2	1.5
2.7	3.4	4.2	2.0
2.6	3.7	3.8	S 25 m Downstream of D
2.6	3.7	3.8	0.5
2.6	3.7	3.8	1.0
2.6	3.6	3.8	1.5
2.6	3.5	3.8	2.0
3.1	3.5	3.9	S 50 m Downstream of D
3.1	3.6	3.9	0.5
3.2	3.6	3.8	1.0
3.1	3.5	3.8	1.5
3.1	3.5	3.8	2.0
2.8	3.4	3.7	S 100 m Downstream of D
2.8	3.4	3.7	0.5
2.8	3.4	3.7	1.0
2.8	3.4	3.7	1.5
2.8	3.3	3.7	2.0
2.6	3.1	3.7	S 200 m Downstream of D
2.6	3.1	3.7	0.5
2.6	3.1	3.7	1.0
2.6	3.1	3.7	1.5
2.6	3.1	3.7	2.0
3.1	3.5	3.7	S 400 m Downstream of D
3.0	3.5	3.7	0.5
2.9	3.5	3.7	1.0
2.9	3.5	3.7	1.5
2.9	3.5	3.7	2.0
2.9	3.5	3.7	2.5
2.9	3.5	3.8	S 1900 m Downstream of D
2.9	3.5	3.8	0.5
2.9	3.5	3.8	1.0
2.9	3.5	3.8	1.5
2.9	3.5	3.8	2.0
2.9	3.5	3.8	2.5

1565 241

Table 3.2-3a

Summary of surface water temperatures (t) at selected pier rap stations, discharge and ambient river temperature differences, wind conditions, discharge rate, river flow, and station location level tributary through November 1976.

	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
Adit River - 5 m offshore at intake	2.1	2.6	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7	2.7
Discharge - 5 m offshore	4.7	3.0	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
50 m downstream of intake	2.5	2.7	2.3	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4
50 m downstream of intake	2.3	2.4	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
100 m downstream of intake	2.7	2.7	2.3	2.3	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4	2.4
200 m downstream of intake	2.0	2.6	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Discharge temp. minus	2.6	0.6	0.0	0.5	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Air temp. (t)	13.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0	18.0
Wind direction	8	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
Wind speed (mph)	3	6	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8
Discharge rate (ft ³ /sec x 10 ³)	0.07	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08
River flow (ft ³ /sec x 10 ³)	55.6	200.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0	220.0
Station location level (ft)	3.0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

Table 3.2-3b continued.

	10	15	20	25	30	35	40	45	50	55	60	65	70	75	80	85	90	95	100
Adit River - 5 m offshore at intake	12.8	12.5	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3	13.3
Discharge - 5 m offshore	12.2	12.2	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4
50 m downstream of intake	12.5	12.4	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6
50 m downstream of intake	12.7	12.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4	13.4
100 m downstream of intake	12.6	12.5	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6	13.6
200 m downstream of intake	12.8	12.4	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5	13.5
Discharge temp. minus	-0.6	-0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Air temp. (t)	18.0	19.5	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0	21.0
Wind direction	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20	20
Wind speed (mph)	10	5	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
Discharge rate (ft ³ /sec x 10 ³)	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08
River flow (ft ³ /sec x 10 ³)	12.5	20.6	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0	35.0
Station location level (ft)	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

NA = Not Available.

6.0 EFFECTS OF COOLING TOWER SALT DRIFT ON AGRICULTURAL CROPS AND
NATURAL VEGETATION

The objective of this investigation was to observe changes in the incidence of plant pathogens or insect damages on agricultural crops and natural vegetation that could be attributed to salt drift from the cooling towers at TMINS as required by the ETS, Appendix B, Section 4.1.2.2.

6.1 PLANT PATHOGENS

Plant pathology transects established in 1973 and surveyed in 1974 (Woodward-Envicon, Inc. 1974, 1975) and 1975 (Potter and Associates 1976), were examined monthly from April through October 1976. Location of transects is shown in Figure 6.1-1.

6.1.1 METHODS

Visual examination of agricultural crops and natural vegetation along the transect lines was conducted in the latter half of each month. A list of all plants in flower was kept to determine differences in flowering time or appearance of flowers. All plant diseases and significant damage due to insects were recorded; collections were made when necessary for identification. The "Forest Insect and Disease Management Manual" (Pennsylvania Department of Environmental Resources 1975) was used for identification of plant diseases; nomenclature of vascular plants followed Gleason and Cronquist (1963).

1565 243

POOR ORIGINAL

6.1.2 RESULTS

In 1976, 219 species of flowering plants were observed (Table 6.1-1). This represented three species more than in 1975. However, 41 species observed in 1975 were not seen in 1976, and 44 species were first in flower in 1976 which were not observed in 1975. The major difference occurred in April and May; 29 (66%) of the new species in 1976 were observed in these months. The spring of 1976 was unusually mild and many species were observed a month earlier than in 1975. Most species were observed at the same locations as in 1975. Differences were related to habitat or were random with respect to TMINS.

Four parasitic diseases were observed on agricultural crops (Table 6.1-2); none caused significant damage. Plant parasitic diseases were observed on 21 taxa (Table 6.1-3). Five diseases observed in 1975 were not seen in 1976, and one disease was observed for the first time in 1976. The total number of parasitic diseases was 19 in 1973, 26 in 1974, 19 in 1975, and 14 in 1976. Powdery mildews were more prominent in 1976 than in 1975; this was probably the result of prolonged high humidity in late summer (Pennsylvania Department of Environmental Resources 1975). Many diseases occurred at the same localities as in 1975; where differences occurred they were related to the presence of host plants and no pattern was observed with respect to the location of TMINS. Vegetation with significant insect damage is listed in Table 6.1-4. Some defoliation was caused by fall webworm on several species, and by Japanese beetle on elm. Other damage was less extensive. No pattern was observed relative to the location of TMINS.

1565 244

POOR ORIGINAL

6.1.3 OTHER VEGETATION DISTURBANCES

Other disturbances to vegetation, both natural and caused by man, were less noticeable in 1976 than in 1975. Several narrow corridors cut through the south woodlot on Three Mile Island were first observed in April 1976. Except for the cut trees, the effects had largely disappeared by mid-summer. Much less disturbance was observed on Shelley Island in 1976, and some of the areas cleared in 1975 were allowed to revegetate in 1976.

Gleason, H.A. and A. Cronquist. 1963. Manual of Vascular Plants of Northeastern United States and Adjacent Canada. D. Van Nostrand Company, Princeton, N.J. 810 pp.

Pennsylvania Department of Environmental Resources. 1975. Forest Insect and Disease Management Manual. Bureau of Forestry, Division of Pest Management, Middletown, Pennsylvania. 658 pp.

Potter, W.A. and Associates. 1976. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Annual Report for 1975. Ichthyological Associates, Inc. 395 pp.

Woodward-Envicon, Inc. 1974. Three Mile Island Nuclear Station, Units 1 and 2. Terrestrial Environmental Studies. Pre-operational Survey. 330 pp.

Woodward-Envicon, Inc. 1975. Three Mile Island Nuclear Station, Units 1 and 2. Terrestrial Environmental Studies. Post-operational Survey. 53 pp.

1565 245

Table 6.1-1

Plants observed in flower in salt drift transects¹ in the vicinity of TMS in 1976.

Taxon ²	Common Name	TMI	Month observed ³			
			Shelley Is.	RT 262	East Side	
Gramineae						
<u>Agrostis</u> sp.	bentgrass	5	5	-	5	
<u>Anthoxanthum odoratum</u>	sweet vernal grass	-	4	-	-	
<u>Briza media</u>	chess	-	-	5	5	
<u>Dactylis glomerata</u>	orchard grass	-	5	5,6	5	
<u>Glyceria melicaria</u>	glyceria	5	-	-	-	
<u>Phleum pratense</u>	timothy	-	-	6	5,6	
<u>Poa annua</u>	annual bluegrass	-	4	5	4,5	
<u>Poa</u> sp.	bluegrass	5	5	5	5	
<u>Trifolium flava</u>	purpletop	-	-	9	9	
Cyperaceae						
<u>Carex lurida</u>	sedge	-	-	-	6	
<u>Carex</u> spp.	sedge	4	4	-	4	
Araceae						
<u>Arisaema triphyllum</u>	Jack-in-the-pulpit	-	4	-	-	
Commelinaceae						
<u>Commelina communis</u>	dayflower	7,8	7,8	-	7,8,9	
Liliaceae						
<u>Asparagus officinalis</u>	asparagus	-	-	5	5	
<u>Erythronium americanum</u>	dog-tooth violet	-	4	-	4	
<u>Heimerocallis fulva</u>	day lily	-	6	6	6	
<u>Muscari botryoides</u>	grape hyacinth	-	-	4	4	
<u>Polygonatum biflorum</u>	Soloman's seal	-	5	-	-	
<u>Smilacena racemosa</u>	false Soloman's seal	-	-	-	5	
<u>Yucca filamentosa</u>	Spanish bayonet	-	-	-	6	
Anaryllidaceae						
<u>Narcissus pseudonarcissus</u>	daffodil	-	-	4	-	
Betulaceae						
<u>Betula nigra</u>	river birch	4	4	-	4	
Fagaceae						
<u>Quercus bicolor</u>	swamp white oak	-	-	-	4	
<u>Quercus borealis</u>	red oak	-	-	-	4	
<u>Quercus palustris</u>	pin oak	-	-	4	4	
Ulmaceae						
<u>Celtis occidentalis</u>	hackberry	-	-	4	4	
Urticaceae						
<u>Boehmeria cylindrica</u>	false nettle	6	7	-	-	
<u>Pilea pumila</u>	cleaveed	7,8	7,8	-	-	
<u>Urtica dioica</u>	stinging nettle	6,7	6,7	-	-	

1565 246

Table 6.1-1 continued.

Taxon	Common Name	TMI	Month Observed		
			Shelley Is	RT 262	East Side
Polygonaceae					
<u>Polygonum caespitosum</u>	smartweed	6,7,9,10	7,9,10	9,10	9,10
<u>Polygonum cuspidatum</u>	Mexican bamboo	-	-	-	7,8
<u>Polygonum hydropiperoides</u>	smartweed	8,9,10	8,9,10	8,9	9,10
<u>Polygonum pennsylvanicum</u>	smartweed	9	8	8,9	8,9,10
<u>Polygonum persicaria</u>	smartweed	6,7,8	6,7,8	6	6
<u>Polygonum sagittatum</u>	tear-thumb	-	-	9	9
<u>Polygonum scandens</u>	false buckwheat	6,7	6,7,8	-	7,8
<u>Polygonum virginianum</u>	Virginia knotweed	7,8	7,8	-	7
<u>Rumex acetosella</u>	sheep sorrel	-	-	5	4,5
<u>Rumex altissimus</u>	water dock	-	5	5	5
Phytolaccaceae					
<u>Phytolacca americana</u>	pokeweed	6-9	6,7,8	7,8,9	6-9
Portulacaceae					
<u>Claytonia virginica</u>	spring beauty	4	4	4	4
Caryophyllaceae					
<u>Agrostemma githago</u>	corn cockle	-	-	-	5
<u>Dianthus armeria</u>	deftford pink	-	-	-	6
<u>Lychnis alba</u>	white campion	-	-	-	4,5,6,9,10
<u>Saponaria officinalis</u>	bouncing bet	-	-	7,8,9	6-9
<u>Silene antirrhina</u>	sleepy catchfly	-	-	-	5
<u>Silene cucubalis</u>	bladder campion	-	-	-	5,6,7
<u>Silene nivea</u>	snowy campion	-	6	-	-
<u>Stellaria media</u>	common chickweed	4	4	4	4
Magnoliaceae					
<u>Liriodendron tulipifera</u>	tulip tree	-	5	-	5
Annonaceae					
<u>Asimina triloba</u>	pawpaw	-	4	-	-
Ranunculaceae					
<u>Aquilegia canadensis</u>	columbine	-	-	4	-
<u>Ranunculus abortivus</u>	small-flowered buttercup	-	4	4	4
<u>Ranunculus acris</u>	buttercup	-	-	-	5
<u>Thalictrum polygamum</u>	tall meadow rue	-	6	-	-
Berberidaceae					
<u>Berberis thunbergii</u>	barberry	-	-	-	4
<u>Podophyllum peltatum</u>	may apple	-	4	-	-
Lauraceae					
<u>Lindera benzoin</u>	spicebush	-	4	-	-
<u>Sassafras albidum</u>	sassafras	-	4	-	4
Papaveraceae					
<u>Chelidonium majus</u>	celandine	-	5	4,5,6	4,5,8
<u>Corydalis flavula</u>	corydalis	4	-	-	4
Cruciferae					
<u>Alliaria officinalis</u>	garlic mustard	4,5,6	4,5,6	4	4,5
<u>Arabisopsis thaliana</u>	mouse-ear cress	-	-	4	4
<u>Barbarea vulgaris</u>	winter cress	4	4,5	4,5	4,5
<u>Brassica campestris</u>	field mustard	-	4,5	5	-
<u>Capsella bursa-pastoris</u>	shepard's purse	-	-	-	4
<u>Cardamine bulbosa</u>	bitter cress	-	4	-	-
<u>Cardamine hirsuta</u>	bitter cress	4	4	4	4
<u>Gentiana lacinata</u>	cut-leaf toothwort	-	4	-	-
<u>Erysimum cheiranthoides</u>	wormseed mustard	-	6	-	-
<u>Hesperis matronalis</u>	dame's rocket	5	5,6	-	-
<u>Lepidium campestre</u>	peppergrass	4	-	4	4,5
<u>Thlaspi arvense</u>	penny-cress	-	-	-	4

1565 247

POOR ORIGINAL

Table 6.1-1 continued.

Taxon	Common Name	TMI	Month Observed		
			Shelley Is	RT 262	East side
Crassulaceae					
<u>Sedum ternatum</u>	stonecrop	-	-	5	-
Saxifragaceae					
<u>Saxifraga virginensis</u>	saxifrage	-	-	4	4
Rosaceae					
<u>Geum canadense</u>	avens	6,7	6,7	-	6
<u>Potentilla norvegica</u>	cinquefoil	-	-	6	6,7
<u>Potentilla recta</u>	cinquefoil	-	-	5,6	5
<u>Potentilla simplex</u>	cinquefoil	5	-	4	4
<u>Prunus avium</u>	sweet cherry	-	-	-	4
<u>Prunus malus</u>	apple	-	-	4	4
<u>Rosa multiflora</u>	multiflora rose	-	-	5	5
<u>Rosa palustris</u>	swamp rose	-	-	-	6
<u>Rubus allegheniensis</u>	blackberry	5	5,6	5	5,6
<u>Rubus flageolaris</u>	dewberry	-	-	5	5
<u>Rubus phoenicolasius</u>	wineberry	-	-	-	5
Mimosaceae					
<u>Albizia julibrissin</u>	mimosa-tree	-	-	6	-
Caesalpiniaceae					
<u>Cassia fasciculata</u>	partridge pea	-	-	-	7,8
<u>Cercis canadensis</u>	redbud	-	-	-	4
Leguminosae					
<u>Apocynum androsaemifolium</u>	groundnut	-	8	-	-
<u>Coronilla varia</u>	crown vetch	-	-	6	5-10
<u>Desmodium illinoense</u>	tick-trefoil	8	-	-	8
<u>Lathyrus latifolius</u>	everlasting pea	-	-	-	6
<u>Medicago lupulina</u>	black medick	-	-	6,7	6
<u>Medicago sativa</u>	alfalfa	-	-	-	6,7,9
<u>Melilotus alba</u>	white sweet clover	-	-	6	6,8
<u>Melilotus officinalis</u>	yellow sweet clover	-	-	5,6,7	5,6,7
<u>Robinia pseudoacacia</u>	black locust	-	-	-	5
<u>Strophostyles helvola</u>	wild bean	-	-	-	8
<u>Trifolium agrarium</u>	white clover	-	6	5,6	5,6,9,10
<u>Trifolium arvense</u>	rabbit's foot clover	-	-	-	8
<u>Trifolium hybridum</u>	white clover	-	-	6,7	6,7
<u>Trifolium pratense</u>	red clover	-	-	5,7-10	5-10
<u>Trifolium procumbens</u>	hop clover	-	-	5	5
<u>Vicia cracca</u>	cow vetch	-	-	-	5,6
Oxalidaceae					
<u>Oxalis sp.</u>	wood sorrel	-	6,7	6,7,9	6,8,9
Geraniaceae					
<u>Geranium carolinianum</u>	Carolina crane's bill	-	-	-	5
Euphorbiaceae					
<u>Euphorbia cyparissias</u>	cypress spurge	-	-	-	4
<u>Euphorbia preslii</u>	spurge	-	-	-	7,8
Anacardiaceae					
<u>Rhus radicans</u>	poison ivy	5	5	5	5
<u>Rhus typhina</u>	staghorn sumac	-	-	-	5
Balsaminaceae					
<u>Impatiens biflora</u>	jewelweed	6,7,8	6-9	7,8	6-9
<u>Impatiens pallida</u>	jewelweed	9	6,7,8	7,8	7,8,9

1565 248

POOR ORIGINAL

Table 6.1-1 continued.

Taxon	Common Name	CMT	Month Observed		
			Shelley Is	RT 262	East Side
Vitaceae					
<u>Vitis aestivalis</u>	summer grape	5	5	5	5
Hypericaceae					
<u>Hypericum perforatum</u>	St. John's wort	-	-	7	6,7
Violaceae					
<u>Viola eriocarpa</u>	yellow violet	-	4	-	-
<u>Viola papilionacea</u>	blue violet	4	4	4	4
<u>Viola rafinesquii</u>	wild pansy	4	-	4	4
<u>Viola sororia</u>	blue violet	4	-	-	-
<u>Viola striata</u>	white violet	4	4,5	-	-
Lythraceae					
<u>Lythrum salicaria</u>	purple loosestrife	-	-	-	8
Onagraceae					
<u>Epilobium coloratum</u>	willow herb	-	-	8	8
<u>Oenothera biennis</u>	evening primrose	-	-	7,9	7,8,10
Umbelliferae					
<u>Conium maculatum</u>	poison hemlock	5	-	-	-
<u>Ranuncus acris</u>	Queen Anne's lace	-	-	6-10	6-10
<u>Osmorhiza lanceolata</u>	sweet cicely	-	-	4	4
Cornaceae					
<u>Cornus florida</u>	flowering dogwood	-	4	4	4
Ericaceae					
<u>Malina latifolia</u>	mountain laurel	-	-	5	-
Primulaceae					
<u>Anagallis arvensis</u>	scarlet pimpernel	-	-	-	8
<u>Lythrum salicaria</u>	fringed loosestrife	-	6,7	-	6,7
Apocynaceae					
<u>Apocynum androsaemifolium</u>	dogbane	6	-	-	5,6,7,9
<u>Vinca minor</u>	periwinkle	-	-	-	4
Asclepiadaceae					
<u>Asclepias incarnata</u>	swamp milkweed	-	6	-	7,8
<u>Asclepias syriaca</u>	common milkweed	-	-	6	6,7
Convolvulaceae					
<u>Convolvulus sepium</u>	black bindweed	-	-	-	6
<u>Ipomoea purpurea</u>	wild morning-glory	-	-	-	8,9
Polemoniaceae					
<u>Phlox divaricata</u>	blue phlox	-	-	4	4
Hydrophyllaceae					
<u>Hydrophyllum virginianum</u>	waterleaf	-	-	5	-
Scraginaceae					
<u>Echium vulgare</u>	blueweed	-	-	-	6,8
<u>Hackelia virginiana</u>	beggar's lice	7	-	-	8
<u>Veronica virginica</u>	bluebells	-	4	4	4

POOR ORIGINAL

1565 249

Table 6.1-1 continued.

Taxon	Common Name	TMI	Month Observed		
			Shelley Is	RT 262	East Side
Verbenaceae					
<u>Verbena urticifolia</u>	white vervain	7,8	7	8	8
Labiatae					
<u>Glechoma hederacea</u>	ground ivy	4	4,6	4	4
<u>Lamium amplexicaule</u>	henbit	-	-	4	4
<u>Lamium purpureum</u>	dead nettle	-	-	-	4
<u>Munarda ciliatopodia</u>	bergamot	-	7	-	-
<u>Neretta cataria</u>	catnip	-	-	-	6,7,8
<u>Perilla frutescens</u>	perilla	-	-	-	9
<u>Prunella vulgaris</u>	self-heal	-	-	-	8,9,10
<u>Teucrium canadense</u>	wood-sage	7	-	7	6,7
<u>Trichostema dichotomum</u>	blue-curls	-	-	-	8
Solanaceae					
<u>Physalis longifolia</u>	ground cherry	-	-	7,8	8,9
<u>Solanum carolinense</u>	horse nettle	-	-	-	6-9
<u>Solanum dulcamara</u>	bittersweet	-	-	5,6,7	6
Scrophulariaceae					
<u>Chelone glabra</u>	turtlehead	-	9	-	-
<u>Linaria vulgaris</u>	butter-and-eggs	-	-	5,7-10	6-10
<u>Verbascum blattaria</u>	moth mullein	-	-	-	6,8
<u>Verbascum thapsis</u>	common mullein	-	-	6	6,7,8
<u>Veronica serpyllifolia</u>	speedwell	-	4	5	5
Plantaginaceae					
<u>Plantago lanceolata</u>	English plantain	-	-	6,7	6,7
Rubiaceae					
<u>Galium aparine</u>	cleavers	5	-	-	5
<u>Houstonia sacrales</u>	bluets	-	-	-	4
Caprifoliaceae					
<u>Lonicera japonica</u>	Japanese honeysuckle	5,6	5,6,9	5-10	5-10
<u>Lonicera tatarica</u>	Tartarian honeysuckle	-	4,5	-	-
<u>Sambucus canadensis</u>	common elderberry	-	5,6	6	5,6
<u>Viburnum prunifolium</u>	black haw	4	-	-	4
Valerianaceae					
<u>Valerianella locusta</u>	corn salad	-	-	4	4
Cucurbitaceae					
<u>Sicora angulatus</u>	bur cucumber	8,9	8,9	-	8
Campanulaceae					
<u>Triodamus perfoliata</u>	Venus' looking glass	-	-	-	6
Lobeliaceae					
<u>Lobelia cardinalis</u>	cardinal flower	-	8,9	-	-
<u>Lobelia inflata</u>	Indian tobacco	-	-	-	9
<u>Lobelia siphilitica</u>	great lobelia	-	-	-	8,9,10

POOR ORIGINAL

1565 250

Table 6.1-1 continued.

Taxon	Common Name	TMI	Month observed		
			Shelley Is	RT 262	East Side
Asteraceae (Compositae)					
<u>Achillea millefolium</u>	yarrow	-	-	5,6,7,10	6,7
<u>Ambrosia artemisiifolia</u>	ragweed	-	-	8	8,9
<u>Ambrosia trifida</u>	giant ragweed	-	8	8	8,9
<u>Anthemis cotula</u>	chamomille	-	-	5	5
<u>Arctium minus</u>	burdock	-	-	-	8,9,10
<u>Aster cordifolius</u>	aster	-	-	9,10	9,10
<u>Aster divaricatus</u>	wood aster	-	-	9	9
<u>Aster ericoides</u>	heath aster	-	9,10	9,10	9,10
<u>Aster novae-angliae</u>	New England aster	-	-	-	9,10
<u>Aster prenanthoides</u>	aster	-	-	-	9,10
<u>Aster simplex</u>	aster	-	-	-	9,10
<u>Bidens frondosa</u>	beggar-ticks	-	-	9,10	10
<u>Bidens laevis</u>	bur-marigold	-	9	9	9
<u>Chrysanthemum leucanthemum</u>	ox-eye daisy	-	-	9	9
<u>Cichorium intybus</u>	chicory	6	-	5-8,10	5,6
<u>Cirsium arvense</u>	Canada thistle	-	-	6-10	6-10
<u>Cirsium vulgare</u>	bull thistle	-	-	6,9	6,7,8
<u>Conyza canadensis</u>	horseweed	-	-	6	7,8
<u>Erechtites hieracifolia</u>	fireweed	-	-	7,8	6,8
<u>Erigeron annuus</u>	fleabane	-	-	-	9
<u>Eupatorium fistulosum</u>	Joe-pye weed	5,6	4,5,6	5-8	5-10
<u>Eupatorium perfoliatum</u>	boneset	-	7,8	-	7,8
<u>Eupatorium rugosum</u>	white snakeroot	-	7,8	8	8,9
<u>Galinsoga ciliata</u>	galinsoga	7-10	8,9,10	8,9,10	6-10
<u>Gnaphalium obtusifolium</u>	cudweed	-	8	8,9	7-10
<u>Helenium autumnale</u>	sneezeweed	-	-	9	-
<u>Helianthus decapetalus</u>	sunflower	-	9,10	-	-
<u>Hieracium pilosella</u>	mouse-ear hawkweed	8	7,8,9	-	9
<u>Hieracium pratense</u>	hawkweed	-	-	5	5
<u>Lactuca canadensis</u>	wild lettuce	-	-	5	5
<u>Lactuca scariola</u>	wild lettuce	-	-	8	-
<u>Lapsana communis</u>	nipplewort	-	-	7	7,8,9
<u>Picris hieracioides</u>	bitterweed	-	-	5	-
<u>Rudbeckia hirta</u>	black-eyed susan	-	-	7-10	-
<u>Rudbeckia laciniata</u>	coneflower	-	-	6	6,7
<u>Rudbeckia triloba</u>	coneflower	8	7	7,8	9
<u>Solidago canadensis</u>	goldenrod	-	-	-	8
<u>Solidago flexicaulis</u>	goldenrod	-	-	9	9
<u>Solidago gigantea</u>	goldenrod	-	-	-	9
<u>Solidago graminifolia</u>	grass-leaf goldenrod	7,8	8	8	8,9
<u>Solidago juncea</u>	early goldenrod	-	-	8,9	7,8,9
<u>Solidago nemoralis</u>	little grey goldenrod	7	-	7,8,9	7
<u>Solidago rigida</u>	rough goldenrod	-	-	8	-
<u>Taraxacum officinale</u>	dandelion	8,9,10	-	8,9,10	8,9,10
<u>Tragopogon pratensis</u>	goat's beard	4	4,5	4-10	4-10
<u>Verbesina alternifolia</u>	wingstem	-	-	5	5
<u>Vernonia noveboracensis</u>	ironweed	8,9	7-10	7-10	8,9
		-	-	-	8,9

1. Location of Transects

Three Mile Island, south woodlot (PPT-9A1-9B1) - TMI

Shelley Island, north woodlot and nature trail (PPT-13A1-15A1, PPT-13S1-15A1) - Shelley Is

Route 262 (PPT-11B1-14B1) - RT 262

Combined routes (PPT-12B1-22B1, PPT-2A1-5A1, PPT-5A2-7C1) - East Side (Route 441 and adjacent routes).

2. Arrangement of families and nomenclature follows Gleason and Cronquist (1963); scientific names arranged alphabetically within families.

3. Month indicated by number (4 = April, 5 = May, etc.).

1565 251

POOR ORIGINAL

Table 6.1-2

Parasitic plant diseases observed on agricultural crops in plant pathology transects, April through October 1976.

Host Species	Disease ¹	Location ²
<u>Cucurbita pepo</u> pumpkin	powdery mildew <u>Erysiphe cichoracearum</u>	1B1-8B1
<u>Trifolium pratense</u> red clover	powdery mildew <u>Erysiphe polygoni</u>	1B1-8B1, 14B1-11B1
<u>Triticum aestivum</u> wheat	wheat smut <u>Ustilago ritici</u>	1B1-8B1, 2A1-5A1
<u>Zea mays</u> corn	corn smut <u>Ustilago maydis</u>	5A2-7C1

¹ Nomenclature from U.S. Department of Agriculture, 1960. Index of plant diseases in the United States, Agriculture Handbook No. 165. U.S. Government Printing Office, Washington, D.C. 531 pp.

² Locations keyed as follows (prefix PPT omitted in Table)
 PPT-14B1 to PPT-11B1 - Plot 24, Cly Road (Rt. 262) to Midway Road, Goldsboro
 PPT-9A1 to PPT-9B1 - TMI, South woodlot
 PPT-15A1 to PPT-13A1 - Shelley Island Nature Trail
 PPT-15A1 to PPT-13A1 - Shelley Island, north woodlot baseline
 PPT-1B1 to PPT-8B1 - Rt. 441 from first farm south of RR at Royalton to south access to TMI
 PPT-2A1 to PPT-5A1 - Geyers Church Road from Rt. 441 to Gingrich Road to Zion Road to Pecks Road
 PPT-5A2 to PPT-7C1 - Pecks Road at Rt. 441 to Falmouth Turnpike (RD 4).

1565 252

Table 6.1-3

Parasitic plant diseases observed on natural vegetation in plant pathology transects, April through October 1976.

Host Species	Disease ¹	Location ²
<u>Acer negundo</u>	leaf spot <u>Phyllosticta minima</u>	181-881, 15A1-11A1, 15A1-1381
<u>Acer saccharinum</u>	leaf spot <u>Phyllosticta minima</u>	181-881, 15A1-11A1, 15A1-1381
<u>Ambrosia artemisiifolia</u>	powdery mildew <u>Erysiphe cichoracearum</u>	2A1-5A1, 5A2-701
<u>Ambrosia trifida</u>	powdery mildew <u>Erysiphe cichoracearum</u>	1481-1181
<u>Aster simplex</u>	leaf rust <u>Coleosporium solidaginis</u>	1481-1181
<u>Carya ovalis</u>	anthracnose <u>Gnomonia caryae</u>	2A1-5A1, 1481-1181
<u>Fraxinus</u> sp.	anthracnose <u>Gloeosporium aridum</u>	181-881, 5A2-701, 1481-1181, 15A1-1381
<u>Lonicera japonica</u>	powdery mildew <u>Erysiphe cichoracearum</u>	181-881 *
<u>Platanus occidentalis</u>	anthracnose <u>Gnomonia platani</u>	181-881
<u>Platanus occidentalis</u>	powdery mildew <u>Microsphaera alni</u>	1481-1181
<u>Pyrus malus</u>	apple scab <u>Venturia inaequalis</u>	1481-1181
<u>Quercus bicolor</u>	anthracnose <u>Gnomonia venetia</u>	2A1-5A1
<u>Rubus allegheniensis</u>	orange rust <u>Puccinia nitens</u>	2A1-5A1, 5A1-701, 1481-1181, 15A1-13A1
<u>Rudbeckia laciniata</u>	powdery mildew <u>Erysiphe cichoracearum</u>	1481-1181
<u>Scrophularia marilandica</u>	powdery mildew <u>Erysiphe cichoracearum</u>	9A1-981
<u>Solidago canadensis</u>	leaf rust <u>Coleosporium solidaginis</u>	181-881, 1481-1181
<u>Solidago canadensis</u>	powdery mildew <u>Erysiphe cichoracearum</u>	181-881
<u>Solidago graminifolia</u>	tar spot <u>Placosphaera haydeni</u>	1481-1181
<u>Solidago rugosa</u>	leaf rust <u>Coleosporium solidaginis</u>	9A1-981, 1481-1181
<u>Solidago rugosa</u>	powdery mildew <u>Erysiphe cichoracearum</u>	181-881, 9A1-981, 1481-1181
<u>Syringa vulgaris</u>	powdery mildew <u>Microsphaera alni</u>	181-881, 5A2-701
<u>Trifolium pratense</u>	powdery mildew <u>Erysiphe polygoni</u>	181-881, 2A1-5A1, 1481-1181
<u>Verbesina alternifolia</u>	powdery mildew <u>Erysiphe cichoracearum</u>	181-881, 2A1-5A1, 5A2-701, 9A1-981, 1481-1181, 15A1-1381
<u>Vernonia noveboracensis</u>	powdery mildew <u>Erysiphe cichoracearum</u>	2A1-5A1

¹ Nomenclature from USDA (1960)

² Locations listed in Table 6.1-2.

1565 253

Table 6.1-4


Vegetation exhibiting significant insect damage in plant pathology transects, April through October 1976.

Host Species	Insect Agent	Location*
<u>Acer saccharinum</u>	leaf gall <u>Eriophyes</u> sp.	13A1-15A1
<u>Celtis occidentalis</u>	leaf gall <u>Pachypsilla</u> sp.	1B1-8B1, 2A1-5A1, 5A2-7C1, 14B1-11B1, 15A1-13A1, 15A1-13S1
<u>Juglans nigra</u>	fall webworm <u>Hyphantria cunea</u>	14B1-11B1
<u>Prunus serotina</u>	gall mite <u>Eriophyes</u> sp.	9A1-9B1
<u>Prunus serotina</u>	tent caterpillar <u>Malacosoma americanum</u>	1B1-8B1, 2A1-5A1
<u>Prunus serotina</u>	fall webworm <u>Hyphantria cunea</u>	1B1-8B1, 2A1-5A1, 14B1-11B1
<u>Pyrus malus</u>	fall webworm <u>Hyphantria cunea</u>	14B1-11B1
<u>Robinia pseudoacacia</u>	locust leaf miner <u>Odontota dorsalis</u>	1B1-8B1, 9A1-9B1
<u>Sassafras albidum</u>	Japaness beetle <u>Popillia japonica</u>	2A1-5A1
<u>Ulmus americana</u>	Japaness beetle <u>Popillia japonica</u>	1B1-8B1, 5A2-7C1, 14B1-11B1
<u>Vitis</u> sp.	leaf gall unidentified	1B1-8B1

* Locations listed in Table 6.1-2.

POOR ORIGINAL

1565 254



POOR ORIGINAL

1565 255

6.2 QUANTITATIVE VEGETATION STUDIES

6.2.1 METHODS

Woodlots and fields surveyed in 1973, 1974, and 1975 were surveyed during late August through mid-October 1976. The bottomland hardwood forest on Shelley Island and the south woodlot on Three Mile Island were surveyed for trees, saplings, shrubs, vines, and groundcover (Figure 6.1-1). The north field, south field, and field near the south ponds on Three Mile Island, and the field near the 500 kv substation just east of Route 441 were sampled for groundcover. Permanent plots and transect lines used in previous years (1973-1975) were used again. Methods for sampling followed those of previous years (Woodward-Envicon, Inc. 1974, 1975; Potter and Associates 1976; Nardacci et al. 1976).

Data from 1976 were compared to those from 1973, the pre-operational sampling, (Woodward-Envicon, Inc. 1974), 1974 (Woodward-Envicon, Inc. 1975) and 1975 (Potter and Associates 1976). Plot by plot tests for significance of changes were made using Student's t-test for paired observations. The number of stems was used for trees, saplings, shrubs, and woody vines, and the percent coverage for all groundcover plots. Statistical formulas used are given in Potter and Associates (1976). Data from 1976 were statistically compared with those from 1975 and 1973. The comparisons with 1974 data were not tested statistically since 1974 and 1973 data were similar in general, and significance of changes from 1974 to 1976 can be inferred from 1976 to 1975 and 1976 to 1973 tests. Significance tests were made on the same taxa as in 1975, except where a taxon was absent or in a frequency too low to permit tests (present in only one or two plots) in 1976.

POOR ORIGINAL

1565 256

Common names are used throughout the text and in the tables where space permits. To avoid confusion, scientific names are used in tables and given at the first use of each taxon in the text. Scientific nomenclature follows Gleason and Cronquist (1963).

6.2.2 RESULTS

In the bottomland hardwood forest on Shelley Island changes in the overstory (trees and saplings) were small (Table 6.2-1). Elm (Ulmus americana) decreased between 1973 and 1976, and silver maple (Acer saccharinum) saplings increased significantly between 1973 and 1976. There were no significant changes in shrubs and vines between 1976 and 1975 or 1973 (Table 6.2-2). The most important groundcover taxa were jewelweed (Impatiens spp.), Virginia knotweed (Polygonum virginianum), Japanese honeysuckle (Lonicera japonica), violet (Viola spp.), poison ivy (Rhus radicans), and false nettle (Boehmeria cylindrica); 57 taxa were represented (Table 6.2-3). Non-vegetative cover decreased significantly between 1975 and 1976, but increased significantly from 1973 to 1976 (Table 6.2-4). Some of the areas cleared in 1975 were again covered with vegetation in 1976, causing a decrease in the amount of non-vegetative cover. Jewelweed increased significantly in 1976, and was one of the important invaders in areas previously cleared. Other significant increases observed included poison ivy (1975-1976), Virginia creeper (Parthenocissus quinquefolia) (1975-1976), Japanese honeysuckle (1973-1976), false nettle (1973-1976), and avens (Geum canadense) (1973-1976). Significant decreases occurred between 1975 and 1976 in avens, and between 1973 and 1976 in Virginia knotweed, violet, poison ivy, Virginia creeper, and clearweed

POOR ORIGINAL

1565 257

(Pilea pumila) (Table 6.2-4). Seedlings of nine trees were found (Table 6.2-5). Silver maple decreased but was still the most abundant; ash (Fraxinus sp.) increased and was the second most abundant, with river birch (Betula nigra) third in abundance.

In the south forest on Three Mile Island the only significant change was a decrease in black locust (Robinia pseudoacacia) saplings (Table 6.2-6). Non-significant decreases also occurred for most other dominant trees and saplings. There were no significant changes in the shrub or vine taxa in this forest (Table 6.2-7). The groundcover is divided into "vine" and "non-vine" portions of the forest. The most important plants in the "vine" portion were clearweed, Japanese honeysuckle, violet, Virginia knotweed, and false nettle; non-vegetative cover had the highest importance value in the forest groundcover (Table 6.2-8). Thirty-seven taxa were recorded (40 in 1975). Non-vegetative cover increased significantly from 1975 to 1976 and 1973 to 1976 (Table 6.2-9). Violet and avens also increased from 1975 to 1976; Virginia knotweed, false buckwheat (Polygonum scandens), jewelweed, pokeweed (Phytolacca americana), Virginia creeper, and poison ivy decreased from 1975 to 1976. Some of these changes continue trends from 1973 to 1975, others such as Virginia creeper and jewelweed are reversals of trends. Seedlings of five tree species were found; tree-of-heaven (Ailanthus altissima) was the most abundant (Table 6.2-10).

In the "non-vine" portion of the south forest dominants were clearweed, grasses, goldenrod (Solidago spp.), false nettle, and avens (Table 6.2-11). Non-vegetative cover had the highest importance value in the groundcover.

Forty taxa were recorded (46 in 1975). Significant increases from 1975 to 1976 were found in non-vegetative cover, and clearweed, and significant decreases from 1975 to 1976 in violet, blackberry (Rubus spp.), false buckwheat, and cinquefoil (Potentilla simplex) (Table 6.2-12). All of these except false buckwheat showed the same change from 1973 to 1976 (non-significant for violet). Other significant changes from 1973 to 1976 (Table 6.2-12) were the same as changes reported for 1973 to 1975. Seedlings of five tree species were found (Table 6.2-13).

In the south field on Three Mile Island, 66 taxa were found in 1976 (Table 6.2-14). Heath aster (Aster ericoides), grasses, goldenrod, blackberry, and white sweet clover (Melilotus alba) were the dominant taxa. Non-vegetative cover, goldenrod, and Canada thistle (Cirsium arvense) decreased significantly, and blackberry increased from 1975 to 1976 (Table 6.2-15). Significant decreases from 1973 to 1976 were found in Canada thistle, nodding foxtail (Setaria faberii), horse nettle (Solanum carolinense), ragweed (Ambrosia artemisiifolia), and beggar-ticks (Bidens frondosa), and significant increases in heath aster, blackberry, and non-vegetative cover from 1973 to 1976. Seedlings of three taxa were found (Table 6.2-16).

In the field near the south ponds on Three Mile Island, 38 taxa were recorded (Table 6.2-17). Heath aster, grasses, evening primrose (Oenothera biennis), goldenrod, and ragweed were the dominants. Significant increases from 1975 to 1976 were found in heath aster, goldenrod, ragweed, and smartweed (Polygonum spp.), and significant decreases from 1975 to 1976 in sheep sorrel (Rumex acetosella), Canada thistle, horse nettle, and horsetweed (Conyza

1565 259

canadensis) (Table 6.2-18). Significant increases from 1973 to 1976 were found for heath aster, evening primrose, and non-vegetative cover, and decreases for goldenrod, Canada thistle, smartweed, nodding foxtail, horse nettle, and horseweed. No beggar-ticks or cocklebur (Xanthium strumarium) were found in 1976 (Table 6.2-18). Seedlings of silver maple and ash were found; these represent the first seedlings encountered in this field (Table 6.2-19).

In the north field on Three Mile Island, 80 taxa were recorded in 1976 (Table 6.2-20). Heath aster, grasses, goldenrod, blackberry, and Japanese honeysuckle were the dominants. Significant increases from 1975 to 1976 were found for goldenrod, and decreases for blackberry, tick trefoil (Desmodium sp.), nodding foxtail, and St. John's wort (Hypericum punctatum) (Table 6.2-21). These changes were also found from 1973 to 1976, plus increases in heath aster and non-vegetative cover, and decreases in horse nettle and ragweed. Seedlings of four taxa were found; three of these represented new taxa, and eight species of seedlings found in previous years were not recorded in 1976 (Table 6.2-22).

In the field near the 500 kv substation, east of Route 441, 39 taxa were recorded in 1976 (Table 6.2-23). Heath aster, goldenrod, crown vetch (Coronilla varia), and Canada thistle were the dominants. Heath aster and goldenrod were dominant in the upper part of the field, with crown vetch forming dense cover on much of the lower part of the field. Canada thistle was found mostly on the upper part of the field. The only significant changes between 1975 and 1976 were an increase in goldenrod and a decrease in

1565 260

POOR ORIGINAL

common plantain (Plantago major) (Table 6.2-24). Goldenrod and crown vetch increased significantly from 1974 to 1976; heath aster, common plantain, dandelion (Taraxacum officinale), and red clover (Trifolium pratense) decreased from 1974 to 1976 (Table 6.2-24). Seedlings of white pine (Pinus strobus), and red maple (Acer rubrum) were found (Table 6.2-25).

6.2.3 DISCUSSION

As in 1975, there were few changes in the overstory and understory of the forests in 1976. The increase in jewelweed and decrease in non-vegetative cover in the Shelley Island forest were related to less disturbance of the forest in 1976. No plots were omitted because of new disturbance in this forest in 1976.

In the Three Mile Island forest some changes, including increase in non-vegetative cover and violet, and decrease in goldenrod, blackberry, and cinquefoil are probably successional in nature. This trend was also observed in 1975 (Potter and Associates 1976). Other changes show no pattern with respect to succession and may represent seasonal responses to growing conditions in the forests. Local disturbances by man and animals may also cause temporary changes in the patterns of vegetation. No pattern was apparent when the Shelley Island forest and the Three Mile Island forest were compared that could be related to the operation of TMINS.

Most of the changes in the four fields were successional in nature: increases in perennials such as heath aster, goldenrod, and blackberry and decreases in annuals including horseweed, nodding foxtail, beggar-ticks,

1565 261

and ragweed, and short-lived perennials including horse nettle and Canada thistle. This pattern is evident in the south field (Table 6.2-15) and the field near the south ponds (Table 6.2-18). In the north field on Three Mile Island an increase in moisture is evidently occurring at the edge of the field near the TMINS dike. Cat-tail (Typha latifolia) was recorded for the first time in 1976; a small stand has developed along the base of the dike. There was also more open water in 1976 than in 1975.

No pattern of change was observed in any of the fields, or in comparison between the fields with respect to the location of TMINS, that could be related to the operation of the TMINS cooling towers.

Gleason, H.A. and A. Cronquist. 1963. Manual of Vascular Plants of Northeastern United States and Adjacent Canada. D. Van Nostrand Company, Princeton, N.J. 810 pp.

Nardacci, G.A., W.A. Potter, J.H. Epler, III, R.F. Eppley, Jr., R.E. Evans, H.A. Hagerty, J.H. Kennedy, B.F. Lathrop, R.W. Malick, Jr., J.D. Montgomery, J.L. Polk, P.C. Ritson, and L. Wike. 1976. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Supplemental Report for 1975. Ichthyological Associates, Inc. 249 pp.

Potter, W.A. and Associates. 1976. An ecological study of the Susquehanna River in the vicinity of the Three Mile Island Nuclear Station. Annual Report for 1975. Ichthyological Associates, Inc. 395 pp.

Woodward-Envicon, Inc. 1974. Three Mile Island Nuclear Station, Units 1 and 2. Terrestrial Environmental Studies. Pre-operational Survey. 330 pp.

Woodward-Envicon, Inc. 1975. Three Mile Island Nuclear Station, Units 1 and 2. Terrestrial Environmental Studies. Post-operational Survey. 53 pp.

1565 262

Table 6.2-1

Comparison of trees and saplings on 24 selected plots in the bottomland hardwood forest on Shalley Island, 1973-1976.

Species	Common Name	Number of Plots				Number of Stems				Basal Area				1976 vs 1975		1976 vs 1973	
		1973	1974	1975	1976	1973	1974	1975	1976	1973	1974	1975	1976	t	Significance ¹	t	Significance
TREES																	
<i>Acer saccharinum</i>	silver maple	12	10	11	10	21	20	18	15	25.106	23.114	28.257	26.895	-1.606	NS	-1.904	NS
<i>Fraxinus</i> sp.	ash	8	7	7	6	15	14	10	9	3.677	3.488	2.809	2.689	-0.446	NS	-1.596	NS
<i>Ulmus americana</i>	elm	7	7	5	3	10	10	8	5	5.400	5.729	5.921	5.290	-1.702	NS	-2.358	*
<i>Betula nigra</i>	river birch	4	4	3	3	13	13	8	6	5.851	5.790	3.197	2.577	-	-	-	-
<i>Robinia pseudacacia</i>	black locust	3	3	3	3	5	4	5	4	3.091	2.955	4.913	4.471	-	-	-	-
<i>Platanus occidentalis</i>	sycamore	3	3	3	2	3	3	3	2	1.925	1.925	2.187	0.841	-	-	-	-
<i>Carya cordiformis</i>	bitternut hickory	3	3	2	1	4	4	2	1	6.936	6.940	2.377	2.181	-	-	-	-
<i>Sassafras albidum</i>	sassafras	2	2	1	1	2	2	3	1	0.403	0.485	0.632	0.349	-	-	-	-
<i>Liriodendron tulipifera</i>	tulip tree	1	1	1	1	1	1	1	1	0.921	1.068	1.068	1.068	-	-	-	-
<i>Catalpa speciosa</i>	catalpa	1	1	1	1	1	1	1	1	0.442	0.442	0.394	0.394	-	-	-	-
SAPLINGS																	
<i>Acer saccharinum</i>	silver maple	10	12	13	14	36	44	656	404	-	-	-	-	-0.582	NS	2.297	*
<i>Acer negundo</i>	box elder	9	10	10	12	69	77	49	47	-	-	-	-	0.628	NS	-0.458	NS
<i>Fraxinus</i> sp.	ash	7	8	7	10	40	44	27	22	-	-	-	-	-0.146	NS	-0.596	NS
<i>Ulmus americana</i>	elm	7	7	7	8	25	23	18	21	-	-	-	-	0.674	NS	-0.065	NS
<i>Carya cordiformis</i>	bitternut hickory	3	3	4	5	14	17	22	15	-	-	-	-	-	-	-	-
<i>Prunus serotina</i>	black cherry	2	4	5	4	2	7	18	23	-	-	-	-	-	-	-	-
<i>Morus rubra</i>	red mulberry	1	3	4	4	1	6	6	7	-	-	-	-	-	-	-	-
<i>Asimina triloba</i>	paw-paw	2	3	3	4	33	47	54	28	-	-	-	-	-	-	-	-
<i>Sassafras albidum</i>	sassafras	0	2	2	2	0	29	36	18	-	-	-	-	-	-	-	-
<i>Liriodendron tulipifera</i>	tulip tree	0	1	2	1	0	1	6	5	-	-	-	-	-	-	-	-
<i>Ailanthus altissima</i>	tree-of-heaven	0	0	2	1	0	0	2	1	-	-	-	-	-	-	-	-
<i>Celtis occidentalis</i>	hackberry	2	1	2	1	2	1	2	1	-	-	-	-	-	-	-	-
<i>Platanus occidentalis</i>	sycamore	1	1	1	1	1	1	1	1	-	-	-	-	-	-	-	-
<i>Quercus borealis</i>	red oak	1	1	1	1	1	1	1	1	-	-	-	-	-	-	-	-
<i>Robinia pseudacacia</i>	black locust	1	1	1	1	1	1	1	1	-	-	-	-	-	-	-	-
<i>Betula nigra</i>	river birch	0	0	2	0	0	0	2	0	-	-	-	-	-	-	-	-
<i>Carya</i> sp.	hickory	0	0	1	0	0	0	1	0	-	-	-	-	-	-	-	-
<i>Cornus florida</i>	flowering dogwood	0	0	1	0	0	0	1	0	-	-	-	-	-	-	-	-

¹ Significance tests made on taxa occurring in five or more plots; ** = significant at 0.01 level, * = significant at 0.05 level, NS = not significant.

Table 6.2-2

Comparison of shrubs and vines on 24 selected plots in the bottomland hardwood forest on Shalley Island, 1933-1976.

Species	Number of plots				Number of stems				1976 vs 1975		1976 vs 1973	
	1933	1976	1973	1976	1973	1976	1973	1976	t	Significance	t	Significance
<i>Quercus</i>												
<i>Liquidambar styraciflua</i>	71	22	22	21	252	250	233	-0.738	NS	-0.759	NS	
<i>Alnus incana</i>	11	16	17	15	148	186	144	0.578	NS	0.926	NS	
<i>Alnus serrulata</i>	4	5	3	4	37	19	8	-	-	-	-	
<i>Loniceria latifolia</i>	1	1	2	3	2	2	5	-	-	-	-	
<i>Rhus typhina</i>	0	0	2	1	0	0	1	-	-	-	-	
<i>Staghorn sumac</i>												
<i>Vitis</i>												
<i>Vitis</i> sp.	9	11	13	12	68	61	46	-0.787	NS	-0.325	NS	
<i>Parthenocissus quinquefolia</i>	12	12	8	12	43	50	21	1.933	NS	-0.620	NS	
<i>Rhus radicans</i>	12	11	9	11	43	48	48	0.298	NS	-0.615	NS	
<i>Celastrus scandens</i>	1	1	4	4	3	1	4	-	-	-	-	
<i>Bittersweet</i>	1	1	4	4	3	1	7	-	-	-	-	

1 Significance tests made on taxa occurring in five or more plots; * = significant at 0.01 level, * = significant at 0.05 level, NS = not significant.

POOR ORIGINAL

1565 264

Table 6.2-3

Ground cover occurring in the bottomland hardwood forest on Shelley Island, 1976.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
Non-vegetative cover	-	10.25	34.67	44.92
<i>Impatiens</i> spp.	jewelweed	9.54	18.77	28.31
<i>Polypogon virginianum</i>	Virginia knotweed	8.24	7.44	15.68
<i>Lonicera japonica</i>	Japanese honeysuckle	6.70	7.95	14.65
<i>Viola</i> spp.	violet	8.60	2.81	11.41
<i>Rhus radicans</i>	poison ivy	6.28	4.88	11.16
<i>Boehmeria cylindrica</i>	false nettle	6.05	4.38	10.43
<i>Ceanothus americanus</i>	avena	5.33	1.14	6.47
<i>Parthenocissus quinquefolia</i>	Virginia creeper	5.33	0.96	6.29
<i>Polypogon scandens</i>	false buckwheat	3.38	2.73	6.11
grass (unidentified)	grass	4.03	1.27	5.30
<i>Pilea pumila</i>	clearweed	2.67	1.05	3.72
<i>Celastrus scandens</i>	bittersweet	1.54	1.97	3.51
<i>Oenothera sensibilis</i>	sensitive fern	2.37	1.08	3.45
<i>Cornelina communis</i>	dayflower	1.78	1.31	3.09
<i>Helianthus decapetalus</i>	sunflower	1.54	1.42	2.96
<i>Rubus</i> sp.	blackberry	1.54	0.88	2.42
<i>Verbesina alternifolia</i>	wingstem	1.42	0.52	1.94
<i>Urtica dioica</i>	nettle	1.13	0.67	1.80
<i>Eupatorium rugosum</i>	white snakeroot	1.13	0.50	1.63
<i>Phytolacca americana</i>	pokeweed	1.19	0.40	1.59
<i>Sicyos angulatus</i>	bur cucumber	1.19	0.24	1.43
<i>Polypogon</i> sp.	smartweed	0.95	0.44	1.39
<i>Martmannia struthiopteris</i>	ostrich fern	0.47	0.75	1.22
<i>Vitis</i> sp.	grape	0.83	0.25	1.18
<i>Glechoma hederacea</i>	ground ivy	0.83	0.29	1.12
<i>Oxalis</i> sp.	wood sorrel	0.77	0.06	0.83
<i>Rudbeckia laciniata</i>	coneflower	0.53	0.17	0.70
<i>Lysimachia ciliata</i>	loosestrife	0.59	0.09	0.68
<i>Hesperis matronalis</i>	dame's rocket	0.47	0.05	0.52
<i>Lysimachia nummularia</i>	moneywort	0.30	0.04	0.34
<i>Rumex crispus</i>	water dock	0.30	0.04	0.34
<i>Polypogon cuspidatum</i>	Mexican bamboo	0.12	0.17	0.29
<i>Solidago</i> sp.	goldenrod	0.18	0.10	0.28
<i>Menispermum canadense</i>	moonseed	0.18	0.06	0.24
<i>Teucrium canadense</i>	wood sage	0.18	0.06	0.24
<i>Athyrium filix-femina</i>	lady fern	0.18	0.04	0.22
<i>Carex</i> sp.	sedge	0.18	0.04	0.22
<i>Eupatorium maculatum</i>	Joe-Pye weed	0.18	0.03	0.21
<i>Acalypha virginica</i>	three-seeded mercury	0.18	0.02	0.20
<i>Aster diversiflorus</i>	white wood aster	0.12	0.06	0.18
<i>Potentilla simplex</i>	cinquefoil	0.12	0.03	0.15
<i>Thalictrum polygamum</i>	tall meadow rue	0.12	0.02	0.14
<i>Solanum carolinense</i>	horse nettle	0.12	0.01	0.13
<i>Allium vineale</i>	wild garlic	0.12	0.00*	0.12
<i>Scrophularia marilandica</i>	figwort	0.06	0.03	0.09
<i>Arisaema triphyllum</i>	jack-in-the-pulpit	0.06	0.02	0.08
<i>Monarda clinopodia</i>	bergamot	0.06	0.02	0.08
<i>Rubus strigosus</i>	red raspberry	0.06	0.02	0.08
<i>Gelium aparine</i>	bedstraw	0.06	0.01	0.07
<i>Hypericum punctatum</i>	St. John's wort	0.06	0.01	0.07
<i>Oenothera biennis</i>	evening primrose	0.06	0.01	0.07
<i>Plantago major</i>	common plantain	0.06	0.01	0.07
<i>Rumex acetosella</i>	sheep sorrel	0.06	0.01	0.07
<i>Portulaca oleraceus</i>	grape fern	0.06	0.00	0.06
<i>Chelidonium majus</i>	celandine	0.06	0.00	0.06
<i>Cuscuta</i> sp.	dodder	0.06	0.00	0.06
<i>Stellaria media</i>	chickweed	0.06	0.00	0.06

* Indicates less than 0.005 percent relative coverage.

Table 6.2-4

Comparison of dominant ground cover¹ occurring in the bottomland hardwood forest on Shelley Island, 1973-1976.

Species	Relative Frequency				Relative Coverage				Importance Value				1976 vs 1975		1976 vs 1973	
	1973	1974	1975	1976	1973	1974	1975	1976	1973	1974	1975	1976	t	Significance ²	t	Significance
Non-vegetative cover	9.7	9.8	11.5	10.2	17.0	14.3	49.3	34.7	26.7	24.1	60.8	44.9	-5.716	**	12.612	**
<u>Impatiens</u> spp.	9.0	8.5	7.7	9.5	11.4	13.6	7.2	18.8	20.4	22.1	14.9	28.3	10.633	**	6.023	**
<u>Polygonum virginianum</u>	8.9	9.7	9.1	8.2	18.0	16.3	6.6	7.4	26.9	26.0	15.7	15.7	1.698	NS	-6.850	**
<u>Lonigera japonica</u>	5.4	7.6	7.3	6.7	6.0	9.7	7.6	8.0	11.4	17.3	14.9	14.7	1.478	NS	4.069	**
<u>Viola</u> spp.	9.3	4.3	8.5	8.6	10.5	4.1	3.3	2.8	19.8	8.4	11.8	11.4	0.152	NS	-7.369	**
<u>Rhus radicans</u>	7.8	7.7	4.9	6.3	8.7	6.6	1.7	4.9	16.5	14.3	6.6	11.2	2.202	*	-7.550	**
<u>Boehmeria cylindrica</u>	5.1	6.9	7.7	6.0	4.0	5.4	4.3	4.4	9.1	12.3	12.0	10.4	0.139	NS	2.156	*
<u>Cean. canadense</u>	0.6	4.6	5.5	5.3	0.2	1.3	2.3	1.1	0.8	5.9	7.8	6.5	-2.541	*	8.202	**
<u>Parthenocissus quinquefolia</u>	5.6	3.7	2.9	5.3	2.1	1.7	1.1	1.0	7.7	5.4	4.0	6.3	4.275	**	-3.282	**
<u>Pilea pumila</u>	5.4	5.1	2.7	2.7	4.4	4.4	0.7	1.0	9.8	9.5	3.4	3.7	1.265	NS	-5.645	**

¹ Dominant ground cover is defined as any taxon with percent frequency equal to or greater than .30.² ** = significant at 0.01 level, * significant at 0.05 level, NS = not significant.

Table 6.2-5

Tree seedlings in bottomland hardwood forest on Shelley Island, 1973, 1975, and 1976.

Species	Common Name	Seedlings/Acre		
		1973	1975	1976
<u>Acer saccharinum</u>	silver maple	9663.5	672.8	290.5
<u>Fraxinus</u> sp.	ash	67.3	83.4	150.8
<u>Betula nigra</u>	river birch	0.0	150.1	134.1
<u>Carya cordiformis</u>	bitternut hickory	38.5	66.7	72.6
<u>Prunus serotina</u>	black cherry	81.7	33.4	33.5
<u>Quercus palustris</u>	pin oak	0.0	5.6	22.4
<u>Acer negundo</u>	box elder	96.2	44.5	5.6
<u>Alnus incana</u>	tree-of-heaven	19.2	5.6	5.6
<u>Liriodendron tulipifera</u>	tulip tree	0.0	0.0	5.6

1565 266

POOR ORIGINAL

Table 6.2-6

Comparison of trees and saplings on 17 selected plots in the south forest on Three Mile Island, 1973-1976.

Species	Common Name	Number of Plots					Number of Stems					Basal Area		1976 vs 1975		1976 vs 1973	
		1973	1974	1975	1976	1977	1973	1974	1975	1976	1977	1973	1975	1976	Significance 1	Significance	
TREES																	
<i>Rubia fraxinosa</i>	black locust	14	14	13	13	43	47	33	36	15.062	15.830	10.159	9.185	0.424	NS	-1.187	NS
<i>Prunus serotina</i>	black cherry	7	7	9	9	19	19	16	16	5.828	6.079	5.489	5.677	-0.104	NS	-0.074	NS
<i>Juglans nigra</i>	black walnut	4	4	3	2	4	4	4	2	0.964	1.024	1.168	0.443	-	-	-	-
<i>Quercus palustris</i>	pin oak	1	1	1	1	2	2	2	2	0.578	0.578	0.796	0.927	-	-	-	-
<i>Alnus incana</i>	tree-of-heaven	2	3	3	1	7	9	7	1	1.034	1.159	1.065	0.230	-	-	-	-
<i>Acer negundo</i>	box elder	1	1	1	0	1	1	1	0	0.267	0.267	0.307	0.009	-	-	-	-
SAPLINGS																	
<i>Alnus incana</i>	tree-of-heaven	10	11	9	10	78	69	35	32	-	-	-	-	-0.263	NS	-1.723	NS
<i>Prunus serotina</i>	black cherry	11	14	12	9	54	57	42	33	-	-	-	-	-2.554	NS	-1.895	NS
<i>Rubia fraxinosa</i>	black locust	10	10	8	7	22	20	19	12	-	-	-	-	-1.520	NS	-2.358	NS
<i>Juglans nigra</i>	black walnut	4	5	4	4	5	6	4	5	-	-	-	-	0.372	NS	-0.060	NS
<i>Fraxinus</i> sp.	ash	1	4	3	3	1	24	19	19	-	-	-	-	-	-	-	-
<i>Acer saccharinum</i>	silver maple	0	0	1	1	0	0	1	2	-	-	-	-	-	-	-	-
<i>Celtis occidentalis</i>	hackberry	0	0	0	1	0	0	0	0	-	-	-	-	-	-	-	-
<i>Acer negundo</i>	box elder	1	1	3	1	1	1	3	1	-	-	-	-	-	-	-	-
<i>Sassafras albidum</i>	sassafras	2	3	1	1	3	8	1	1	-	-	-	-	-	-	-	-
<i>Acer rubrum</i>	red maple	0	0	1	0	0	0	0	2	-	-	-	-	-	-	-	-

1 Significance tests made on taxa occurring in five or more plots; ** = significant at 0.01 level, * = significant at 0.05 level, NS = not significant.

Table 6.2-7

Comparison of shrubs and vines on 17 selected plots in the south forest on Three Mile Island, 1973-1976.

Species	Common Name	Number of Plots					Number of Stems					1976 vs 1975		1976 vs 1973	
		1973-1976					1973-1976					t	Significance	t	Significance
		1973	1974	1975	1976	1977	1973	1974	1975	1976	1977				
SHRUBS															
<i>Lonicera hirculus</i>	spicebush	6	7	7	7	15	19	31	36	0.021	NS	-	1.332	NS	-
<i>Rhus typhina</i>	staghorn sumac	0	0	0	1	0	0	0	1	-	-	-	-	-	-
VINES															
<i>Vitis</i> sp.	grape	10	12	10	11	82	82	65	64	0.470	NS	-	-0.410	NS	-
<i>Parthenocissus quinquefolia</i>	Virginia creeper	10	10	11	8	33	40	17	21	-0.376	NS	-	-1.567	NS	-
<i>Rhus radicans</i>	poison ivy	5	7	4	5	23	23	11	18	1.247	NS	-	-0.142	NS	-
<i>Celastrus scandens</i>	bittersweet	0	0	0	2	0	0	0	6	-	-	-	-	-	-

1 Significance tests made on taxa occurring in five or more plots; ** = significant at 0.01 level, * = significant at 0.05 level, NS = not significant.

Table 6.2-8

Ground cover occurring in the "vine" portion of the black locust forest on Three Mile Island, 1976.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
Non-vegetative cover	-	9.32	45.13	54.45
<u>Pilea pumila</u>	clearweed	7.96	11.61	16.57
<u>Lonicera japonica</u>	Japanese honeysuckle	8.14	5.87	14.01
<u>Viola</u> spp.	violet	8.23	3.41	11.64
<u>Polypodium virginicum</u>	Virginia knotweed	6.67	3.21	9.88
<u>Boehmeria cylindrica</u>	false nettle	5.48	4.20	9.68
Grass (unidentified)	grass	5.48	4.04	9.52
<u>Ceanothus canadensis</u>	avens	7.50	1.80	9.30
<u>Polypodium scandens</u>	false buckwheat	4.39	3.67	8.06
<u>Impatiens</u> spp.	jewelweed	5.30	1.84	7.14
<u>Eupatorium rugosum</u>	white snakeroot	3.29	2.20	5.49
<u>Solidago</u> spp.	goldenrod	3.11	2.28	5.39
<u>Phytolacca americana</u>	pokeweed	2.29	2.81	5.10
<u>Verbesina alternifolia</u>	wingstem	2.29	2.74	5.03
<u>Polygonum</u> spp.	smartweed	2.83	1.67	4.50
<u>Teucrium canadense</u>	wood sage	2.47	0.74	3.21
<u>Parthenocissus quinquefolia</u>	Virginia creeper	2.83	0.27	3.10
<u>Rubus</u> sp.	blackberry	2.10	0.50	2.60
<u>Rhus radicans</u>	poison ivy	2.10	0.12	2.22
<u>Celastrus scandens</u>	bittersweet	1.19	0.35	1.54
<u>Rumex crispus</u>	water dock	1.19	0.12	1.31
<u>Urtica dioica</u>	stinging nettle	0.55	0.39	0.94
<u>Camellia sasanqua</u>	dayflower	0.73	1.18	0.91
<u>Vitis</u> sp.	grape	0.55	0.15	0.70
<u>Cucurbita angulatus</u>	bur cucumber	0.64	0.05	0.63
<u>Glechoma hederacea</u>	ground ivy	0.46	0.10	0.56
<u>Carex</u> sp.	sedge	0.46	0.06	0.52
<u>Allium vineale</u>	wild garlic	0.46	0.03	0.49
<u>Hieracium virginianum</u>	beggar's lice	0.27	0.15	0.42
<u>Potentilla simplex</u>	cinquefoil	0.37	0.03	0.40
<u>Galium</u> sp.	wood sorrel	0.37	0.02	0.39
<u>Asarum canadense</u>	three-seeded mercury	0.27	0.04	0.31
<u>Lonicera villosa</u>	wild yam	0.18	0.09	0.27
<u>Leucanthemum laciniatum</u>	coneflower	0.18	0.01	0.19
<u>Asarum canadense</u>	figwort	0.09	0.06	0.15
<u>Aster</u> sp.	aster	0.09	0.02	0.11
<u>Lythrum virginicum</u>	bugleweed	0.09	0.02	0.11
<u>Leguminosae helveticus</u>	wild bean	0.09	0.02	0.11

1565 268

Table 6.2-9

Comparison of the dominant ground cover¹ occurring in the "vine" portion of the south woodlot on Three Mile Island, 1973-1976.

Species	Relative Frequency				Relative Coverage				Importance Value				1976 vs 1975		1976 vs 1973	
	1973	1974	1975	1976	1973	1974	1975	1976	1973	1974	1975	1976	t	Significance ²	t	Significance
Non-vegetative cover	1.2	2.9	9.1	9.3	0.8	2.1	37.7	45.1	2.0	5.0	46.8	56.4	5.115	**	25.800	**
<i>Pilea pumila</i>	9.7	9.9	7.1	8.0	33.5	28.1	12.7	11.6	43.2	28.0	19.8	19.6	-0.909	NS	-6.645	**
<i>Lonicera japonica</i>	6.8	6.8	7.6	8.1	8.4	10.1	5.3	5.9	15.2	16.9	12.9	14.0	1.134	NS	0.135	NS
<i>Viola</i> spp.	5.5	4.8	7.9	8.2	5.2	4.4	2.9	3.4	10.7	9.2	10.6	11.6	2.730	**	0.733	NS
<i>Polygonum virginianum</i>	6.7	5.9	6.3	6.7	5.9	5.1	4.2	3.2	12.6	11.0	10.5	9.9	-2.117	*	-2.404	*
<i>Echmeria cylindrica</i>	5.2	6.8	6.4	5.5	5.6	6.1	4.3	4.2	10.8	12.9	10.7	9.7	-1.686	NS	-0.011	NS
<i>Geum canadense</i>	4.3	6.2	5.8	7.5	1.7	2.6	1.9	1.8	6.0	8.8	7.7	9.3	2.536	*	3.419	**
<i>Polygonum scandens</i>	4.7	5.9	5.0	4.4	6.9	9.7	5.1	3.7	11.6	15.6	10.1	8.1	-2.870	**	-1.975	NS
<i>Impatiens</i> spp.	5.2	5.9	7.0	5.3	2.4	6.2	5.3	1.8	7.6	12.1	12.3	7.1	-6.709	**	-0.546	NS
<i>Solidago</i> spp.	5.0	4.9	4.1	3.1	5.1	3.6	2.2	2.3	10.1	8.5	6.3	5.4	-1.773	NS	-4.310	**
<i>Phytolacca americana</i>	3.2	3.8	3.0	2.3	3.1	4.9	4.8	2.8	6.3	8.7	7.8	5.1	-2.812	**	-0.156	NS
<i>Polygonum</i> spp.	3.2	1.9	2.7	2.8	2.2	1.3	2.1	1.7	5.4	3.2	4.8	4.5	-1.447	NS	-0.269	NS
<i>Ilex</i> spp.	4.6	3.1	2.1	2.5	4.1	1.8	0.6	0.7	8.7	4.9	2.7	3.2	0.818	NS	-5.216	**
<i>Parthenocissus quinquefolia</i>	3.4	4.0	1.6	2.8	0.7	1.6	0.2	0.3	4.1	5.6	1.6	3.1	-2.008	*	3.020	**
<i>Rubus</i> sp.	5.5	4.6	2.6	3.1	2.9	2.2	0.6	0.5	8.4	6.8	3.2	2.6	-1.356	NS	-6.190	**
<i>Rhus radicans</i>	4.7	3.8	3.3	2.1	1.5	0.9	0.5	0.1	6.2	4.7	3.8	2.2	-2.481	*	-5.932	**

¹ Dominant ground cover is defined in Table 6.2-4.

² ** = significant at 0.01 level, * = significant at 0.05 level, NS = not significant.

Table 6.2-10

Tree seedlings in the "vine" portion of the black locust forest on Three Mile Island, 1973-1976.

Species	Common Name	Seedlings/Acre		
		1973	1975	1976
<i>Ailanthus altissima</i>	tree-of-heaven	84.7	9.1	58.8
<i>Prunus serotina</i>	black cherry	50.8	27.3	19.6
<i>Acer negundo</i>	box elder	59.3	9.1	9.8
<i>Salix americana</i>	hackberry	16.9	18.2	9.8
<i>Asclepias tuberosa</i>	asclepias	84.7	9.1	9.8
<i>Acer saccharinum</i>	silver maple	0.0	9.1	0.0
<i>Betula nigra</i>	river birch	0.0	9.1	0.0

1565 269

POOR ORIGINAL

Table 6.2-11

Ground cover occurring in the "non-vine" portion of the black locust forest on Three Mile Island, 1976.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
Non-vegetative cover	-	8.04	24.97	33.01
<u>Pilea pumila</u>	cleaveveed	7.32	15.50	22.82
Grass (unidentified)	grass	7.44	10.63	18.07
<u>Solidago</u> spp.	goldenrod	6.48	7.72	14.20
<u>Boehmeria cylindrica</u>	false nettle	5.64	7.68	13.32
<u>Ceanothus canadensis</u>	avena	7.68	4.06	11.74
<u>Viola</u> spp.	violet	6.36	3.00	9.36
<u>Polypogon virginianum</u>	Virginia knotweed	6.24	3.10	9.34
<u>Impatiens</u> spp.	jewelweed	5.64	3.60	9.24
<u>Lonicera japonica</u>	Japanese honeysuckle	4.32	3.46	7.78
<u>Leucophaea canadensis</u>	wood sage	4.68	2.66	7.34
<u>Phytolacca americana</u>	pokeweed	3.12	3.66	6.78
<u>Rubus</u> spp.	blackberry	3.60	1.33	4.93
<u>Polypogon scandens</u>	false buckwheat	3.12	1.81	4.93
<u>Polypogon</u> spp.	smartweed	2.28	1.47	3.75
<u>Verbesina alternifolia</u>	wingstem	1.20	1.81	3.01
<u>Eupatorium rugosum</u>	white snakeroot	1.56	1.09	2.65
<u>Rhus radicans</u>	poison ivy	2.04	0.25	2.29
<u>Carex</u> sp.	sedge	1.80	0.48	2.28
<u>Coreopsis cornuta</u>	dayflower	1.56	0.31	1.87
<u>Solanum carolinense</u>	horse nettle	1.20	0.18	1.36
<u>Parthenocissus quinquefolia</u>	Virginia creeper	1.20	0.08	1.28
<u>Hackelia virginiana</u>	beggar's lice	0.96	0.22	1.18
<u>Potentilla simplex</u>	cinquefoil	1.08	0.07	1.15
<u>Rumex crispus</u>	water dock	1.09	0.07	1.16
<u>Sisyrinchium angustifolium</u>	bur cucumber	0.72	0.19	0.91
<u>Urtica dioica</u>	stinging nettle	0.48	0.28	0.76
<u>Allium vineale</u>	wild garlic	0.60	0.03	0.63
<u>Acalypha virginica</u>	three-seeded mercury	0.48	0.04	0.52
<u>Oxalis</u> sp.	wood sorrel	0.48	0.02	0.50
<u>Verbena urticifolia</u>	blue vervain	0.24	0.09	0.33
<u>Aster ericoides</u>	heath aster	0.24	0.04	0.28
<u>Celastrus scandens</u>	bittersweet	0.12	0.04	0.16
<u>Fragaria virginiana</u>	strawberry	0.12	0.03	0.15
<u>Lycopus virginicus</u>	bugleweed	0.12	0.03	0.15
<u>Vitis</u> sp.	grape	0.12	0.03	0.15
<u>Ambrosia artemisiifolia</u>	ragweed	0.12	0.02	0.14
<u>Rosa</u> sp.	wild rose	0.12	0.01	0.13
<u>Taraxacum officinale</u>	dandelion	0.12	0.01	0.13
<u>Trifolium repens</u>	white clover	0.12	0.01	0.13
<u>Botrychium dissectum</u>	grape fern	0.12	0.00*	0.12

* Indicates less than 0.001 percent relative coverage.

POOR ORIGINAL

1565 270

Table 6.2-12

Comparison of the dominant ground cover¹ occurring in the "non-vine" portion of the south woodlot on Three Mile Island, 1973-1976.

Species	Relative Frequency					Relative Coverage					Importance Value					1976 vs 1975		1976 vs 1973	
	1973	1974	1975	1976		1973	1974	1975	1976		1973	1974	1975	1976		t	Significance	t	Significance
Non-vegetative cover	1.1	1.7	7.7	8.0		1.3	1.9	20.7	25.0		2.4	3.6	28.4	33.0		2.857	**	17.725	**
<i>Pilea pumila</i>	5.2	7.0	7.3	7.3		5.3	11.4	11.4	15.5		10.5	18.4	18.7	22.8		2.633	*	6.663	**
<i>Solidago</i> spp.	8.1	6.0	6.6	6.5		12.3	8.5	8.1	7.7		20.4	14.3	14.7	14.2		-0.121	NS	-0.932	NS
<i>Psidium coccineum</i>	2.1	5.8	6.0	5.6		0.8	4.7	9.1	7.7		2.9	10.5	15.1	13.3		-1.071	NS	6.977	**
<i>Ceanothus americanus</i>	6.3	7.4	6.7	7.7		2.7	6.1	3.4	4.1		9.0	13.5	10.1	11.7		1.904	NS	4.619	**
<i>Viola</i> spp.	7.2	5.1	6.4	6.4		7.0	6.2	6.1	3.0		14.2	11.3	10.5	9.4		-2.090	*	-1.657	NS
<i>Polypodium virginianum</i>	4.1	6.7	5.7	6.2		3.9	4.5	2.9	3.1		8.0	9.2	8.6	9.3		1.055	NS	1.622	NS
<i>Junonia</i> spp.	4.1	4.0	6.1	5.6		1.6	7.0	4.7	3.6		5.7	7.0	10.8	9.2		-1.307	NS	3.494	**
<i>Leuciseta japonica</i>	4.2	3.7	4.4	4.3		6.9	4.6	3.7	3.5		11.1	8.3	8.1	7.8		-0.079	NS	-0.480	NS
<i>Leuciseta canadensis</i>	2.9	5.4	3.3	4.7		18.1	5.5	1.9	2.7		27.0	10.9	5.2	7.3		1.971	NS	-2.681	**
<i>Phlox</i> spp.	2.6	2.2	2.9	3.1		1.9	2.4	5.0	3.7		4.5	4.6	7.9	6.8		-0.650	NS	2.488	*
<i>Rubus</i> spp.	6.3	5.1	3.7	3.6		7.1	3.8	2.6	1.1		13.4	10.9	6.3	4.9		-2.865	**	-6.368	**
<i>Polypodium polypodioides</i>	3.4	3.7	4.9	3.1		0.6	6.5	4.0	1.8		4.0	12.2	8.9	4.9		-3.435	**	2.148	*
<i>Rhus typhina</i>	4.1	3.1	2.2	2.0		1.3	0.4	0.2	0.3		5.4	3.5	2.4	2.3		-0.114	NS	-2.307	*
<i>Parthenocissus quinquefolia</i>	4.4	4.0	1.3	1.2		1.2	1.0	0.1	0.1		5.6	5.0	1.4	1.3		-0.628	NS	-4.315	**
<i>Potentilla simplex</i>	4.2	4.7	2.7	1.1		0.6	0.9	0.3	0.1		4.8	5.6	3.0	1.2		-3.576	*	-3.968	**

¹ Dominant ground cover is defined in Table 6.2-4.

* = significant at 0.01 level, ** = significant at 0.05 level, NS = Not significant.

222

POOR ORIGINAL

1565 271

Table 6.2-13

Tree seedlings in the "non-vine" portion of the black locust forest on Three Mile Island, 1973, 1975, and 1976.

Species	Crown Name		Seedlings/Acre	
	1973	1975	1976	
<i>Prunus serotina</i>	black cherry	14.1	29.0	29.9
<i>Sassafras albidum</i>	sassafras	0.0	0.0	29.9
<i>Robinia pseudoacacia</i>	black locust	0.0	29.0	14.9
<i>Ailanthus altissima</i>	tree-of-heaven	197.2	14.5	14.9
<i>Betula nigra</i>	river birch	0.0	14.5	14.9

Ground cover occurring in the south field on Three Mile Island, 1976.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
<u>Aster spicatus</u>	heath aster	13.15	36.06	49.31
Grass (unidentified)	grass	13.17	35.48	48.65
<u>Solidago</u> spp.	goldenrod	13.78	14.51	27.79
Non-vegetative cover	-	7.43	2.95	10.78
<u>Rubus</u> spp.	blackberry	5.55	1.59	7.14
<u>Melilotus alba</u>	white sweet clover	3.50	1.96	4.46
<u>Cirsium arvense</u>	Canada thistle	3.42	0.33	4.25
<u>Setaria faberii</u>	nodding foxtail	1.85	1.48	3.33
<u>Hypericum punctatum</u>	St. John's wort	3.14	0.15	3.31
<u>Lonicera japonica</u>	Japanese honeysuckle	1.43	1.37	3.00
<u>Taraxacum officinale</u>	dandelion	2.83	0.11	2.94
<u>Oenothera biennis</u>	evening primrose	2.39	0.19	2.58
<u>Avena canadense</u>	oats	1.76	0.19	2.15
<u>Rumex crispus</u>	water dock	1.45	0.18	2.03
<u>Verbesina alternifolia</u>	wingstem	1.41	0.40	1.89
<u>Aralia virginica</u>	three-seeded mercury	1.74	0.04	1.78
<u>Potentilla simplex</u>	cinquefoil	1.52	0.21	1.73
<u>Conium maculatum</u>	poison hemlock	1.10	0.21	1.41
<u>Thymolaea americana</u>	pokeweed	0.45	0.60	1.20
<u>Solanum carolinense</u>	horse nettle	1.09	0.05	1.14
<u>Oxalis</u> sp.	wood sorrel	1.19	0.03	1.11
<u>Rhus radicans</u>	poison ivy	0.98	0.04	1.01
<u>Elymus</u> spp.	smartweed	0.87	0.03	0.95
<u>Achroia artemisiifolia</u>	ragweed	0.76	0.15	0.91
<u>Scrophularia cylindrica</u>	false nettle	0.76	0.14	0.91
<u>Eupatorium perfoliatum</u>	boneset	0.87	0.03	0.88
<u>Lycopus virginicus</u>	bugleweed	0.87	0.03	0.88
<u>Elymus canadensis</u>	false buckwheat	0.76	0.14	0.88
<u>Viola</u> sp.	violet	0.76	0.12	0.88
<u>Plantago lanceolata</u>	English plantain	0.44	0.36	0.80
<u>Allium vineale</u>	wild garlic	0.45	0.08	0.73
<u>Teucrium canadense</u>	wood sage	0.45	0.05	0.70
<u>Impatiens</u> sp.	jewelweed	0.45	0.03	0.68
<u>Juncus</u> sp.	rush	0.54	0.13	0.67
<u>Vitis</u> sp.	grape	0.54	0.02	0.56
<u>Urtica media</u>	chickweed	0.54	0.01	0.55
<u>Lychnis viscaria</u>	Queen Anne's lace	0.44	0.01	0.45
<u>Phlox inflata</u>	Indian tobacco	0.44	0.01	0.45
<u>Asarum canadense</u>	dogbane	0.33	0.04	0.37
<u>Rumex acetosella</u>	sheep sorrel	0.33	0.02	0.35
<u>Trifolium pratense</u>	willow herb	0.33	0.01	0.34
<u>Achillea millefolium</u>	beggar-ticks	0.22	0.05	0.27
<u>Verbena officinalis</u>	white vervain	0.22	0.03	0.25
Mustard (unidentified)	mustard	0.22	0.03	0.25
<u>Lonicera canadensis</u>	horseradish	0.22	0.01	0.23
<u>Brasica virginiana</u>	strawberry	0.22	0.01	0.23
<u>Potentilla recta</u>	cinquefoil	0.22	0.01	0.23
<u>Trifolium</u> sp.	white clover	0.22	0.01	0.23
<u>Galium</u> sp.	bedstraw	0.11	0.02	0.13
<u>Asclepias tuberosa</u>	beggar's lice	0.11	0.02	0.13
<u>Urtica dioica</u>	sensitive fern	0.11	0.02	0.13
<u>Chenopodium album</u>	lamb's quarters	0.11	0.01	0.12
<u>Lysichiton ciliata</u>	loosestrife	0.11	0.01	0.12
<u>Salvia verticillata</u>	carpetweed	0.11	0.01	0.12
<u>Physalis longifolia</u>	ground cherry	0.11	0.01	0.12
<u>Pilea pumila</u>	cleaver	0.11	0.01	0.12
<u>Rubus strigosus</u>	red raspberry	0.11	0.01	0.12
<u>Anagallis arvensis</u>	scarlet pimpernel	0.11	0.00*	0.11
<u>Desmodium illinoense</u>	tick trefoil	0.11	0.00	0.11
<u>Eupatorium rugosum</u>	white snakeroot	0.11	0.00	0.11
<u>Plantago major</u>	plantain	0.11	0.00	0.11
<u>Polypodium virginicum</u>	milkwort	0.11	0.00	0.11
<u>Scirpus atrovirens</u>	bulrush	0.11	0.00	0.11
<u>Scrophularia marilandica</u>	figwort	0.11	0.00	0.11
<u>Trifolium pratense</u>	red clover	0.11	0.00	0.11
<u>Urtica dioica</u>	stinging nettle	0.11	0.00	0.11
<u>Xanthoxylum</u>	cocklebur	0.11	0.00	0.11

* Indicates less than 0.005 percent relative coverage.

Table 6.2-15

Comparison of dominant ground cover¹ occurring in the south field on Three Mile Island, 1973-1976.

Species	Relative frequency					Relative coverage					Importance Value					1976 vs 1975		1976 vs 1973	
	1973	1974	1975	1976		1973	1974	1975	1976		1973	1974	1975	1976		F	Significance ²	F	Significance ²
<i>Aster multiflorus</i>	1.7	15.9	16.6	13.3		29.9	45.6	45.3	36.1		43.6	61.5	61.9	49.3		-1.625	NS	1.992	*
<i>Solidago</i> spp.	14.1	14.3	16.0	13.3		16.1	20.2	23.7	14.5		30.2	34.5	39.7	27.8		-3.364	**	-0.940	NS
Non-vegetative cover	1.3	1.7	12.6	7.8		0.3	0.5	4.4	3.0		1.6	2.2	17.0	10.8		-2.943	*	7.914	**
<i>Rubus</i> spp.	3.2	5.4	5.0	5.6		1.0	3.1	1.0	1.6		4.2	8.5	6.0	7.1		2.606	*	2.262	*
<i>Cirsium arvense</i>	5.3	6.4	4.9	3.9		2.6	2.9	1.5	0.3		7.9	9.3	6.4	4.2		-2.654	*	-4.117	**
<i>Sesaria faberii</i>	13.1	7.6	1.6	1.8		25.7	5.9	0.8	1.5		38.8	13.5	2.6	3.3		0.962	NS	-13.866	**
<i>Solanum carolinense</i>	10.0	8.0	0.7	1.1		3.8	1.3	0.1	0.1		13.8	9.3	0.8	1.1		0.570	NS	-10.838	**
<i>Ambrosia artemisiifolia</i>	9.4	0.5	0.7	0.8		7.8	2.5	0.1	0.2		17.2	6.0	0.8	0.9		1.595	NS	-9.189	**
<i>Rudens frondosa</i>	5.8	3.0	0.5	0.2		5.6	3.2	0.1	0.1		11.4	6.2	0.6	0.3		1.296	NS	-6.423	**
<i>Corylus canadensis</i> ³	4.2	4.8	-	0.2		1.0	7.9	-	0.0		5.2	12.7	-	0.2		-	-	-	-

¹ Dominant ground cover is defined in Table 6.2-4.² ** = significant at 0.01 level, * = significant at 0.05 level, NS = not significant.³ *Corylus canadensis* was not reported in 1975, and frequency was too low to test in 1976.

Table 6.2-16

Tree seedlings in the south field on Three Mile Island, 1973-1976.

Species	Common Name	Seedlings/Acre	
		1973	1976
<i>Betula nigræ</i>	river birch	296.3	296.9
<i>Acer saccharinum</i>	silver maple	7.4	23.4
<i>Platanus occidentalis</i>	sycamore	0.0	7.8
<i>Robinia pseudoacacia</i>	black locust	0.0	7.8
<i>Viburnum prunifolium</i>	black haw	0.0	7.8
<i>Fragaria</i> sp.	ash	37.0	0.0
<i>Prunus cerasifera</i>	black cherry	7.4	0.0

POOR ORIGINAL

1565 273

Table 6.2-17

Ground cover occurring in the field near the south ponds on Three Mile Island, 1976.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
<u>Aster ericoides</u>	heath aster	9.76	53.81	63.57
Grass (unidentified)	grass	9.53	11.32	20.85
<u>Oenothera biennis</u>	evening primrose	8.87	8.04	16.91
Non-vegetative cover	-	7.98	7.00	14.98
<u>Solidago</u> spp.	goldenrod	8.65	5.61	14.26
<u>Artemisia artemisiifolia</u>	ragweed	6.87	3.33	10.20
<u>Rumex acetosella</u>	sheep sorrel	7.10	1.82	6.92
<u>Cirsium arvense</u>	Canada thistle	7.32	1.50	8.82
<u>Poligonum</u> spp.	smartweed	3.55	1.21	4.76
<u>Taraxacum officinale</u>	dandelion	3.77	0.26	4.03
<u>Setaria faberii</u>	nodding foxtail	1.57	2.37	3.92
<u>Medicago alba</u>	white sweet clover	2.86	0.70	3.58
<u>Rubus</u> spp.	blackberry	2.44	0.61	3.05
<u>Oxalis</u> sp.	wood sorrel	2.44	0.14	2.58
<u>Solanum carolinense</u>	horse nettle	1.77	0.10	1.87
<u>Potentilla simplex</u>	cinquefoil	0.89	0.80	1.69
<u>Lysimachia ciliata</u>	fringed loosestrife	1.33	0.14	1.47
<u>Ceanothus canadensis</u>	avens	1.33	0.11	1.44
<u>Hypericum punctatum</u>	St. John's wort	1.33	0.11	1.44
<u>Daucus carota</u>	Queen Anne's lace	1.11	0.27	1.38
<u>Trifolium</u> sp.	white clover	1.11	0.10	1.21
<u>Rumex crispus</u>	water dock	1.11	0.09	1.20
<u>Stellaria media</u>	chickweed	1.11	0.07	1.18
<u>Fragaria virginiana</u>	strawberry	1.11	0.04	1.15
<u>Carex</u> sp.	sedge	0.67	0.10	0.77
<u>Viola</u> sp.	violet	0.67	0.02	0.69
<u>Desmodium</u> sp.	tick-trefoil	0.44	0.07	0.51
<u>Asclepias incarnata</u>	swamp milkweed	0.44	0.04	0.48
<u>Conium maculatum</u>	poison hemlock	0.44	0.04	0.48
<u>Eupatorium perfoliatum</u>	boneset	0.44	0.03	0.47
<u>Lonicera japonica</u>	Japanese honeysuckle	0.22	0.05	0.27
<u>Hordeum jubatum</u>	horseweed	0.22	0.04	0.26
<u>Juncus</u> sp.	rush	0.22	0.02	0.24
<u>Verbena alternifolia</u>	wingstem	0.22	0.02	0.24
<u>Cassia fasciculata</u>	partridge pea	0.22	0.01	0.23
<u>Polypodium virginicum</u>	false buckwheat	0.22	0.01	0.23
<u>Physalis longifolia</u>	ground cherry	0.22	0.01	0.23
<u>Teucrium canadense</u>	wood sage	0.22	0.01	0.23
Mustard (unidentified)	mustard	0.22	0.01	0.23

POOR ORIGINAL

1565 274

Comparison of dominant ground cover¹ occurring in the field near the south ponds on Three Mile Island, 1973-1976.

Species	Relative Frequency			Relative Coverage			Importance Value			1976 vs 1975		1976 vs 1973		
	1973	1974	1975	1973	1974	1975	1973	1974	1975	F	Significance	F	Significance	
<i>Aster triflorus</i>	0.9	9.9	9.3	9.8	0.1	16.3	37.6	53.8	1.0	26.2	46.9	63.6	6.464	**
<i>Onoclea biennis</i>	0.9	2.7	8.6	8.9	0.4	0.4	11.9	8.0	1.3	3.1	20.7	16.9	-1.229	NS
Non-vegetative cover	0.2	0.2	7.9	8.0	0.1	0.2	8.7	7.0	0.3	0.5	16.6	15.0	0.510	NS
<i>Solidago</i> spp.	0.9	4.5	6.4	8.7	0.4	2.1	4.4	5.6	1.3	6.6	10.8	14.3	1.990	*
<i>Achillea millefolium</i>	10.3	13.7	1.6	6.9	12.4	11.7	0.7	3.3	22.7	25.4	2.3	10.2	6.234	**
<i>Rumex acetosella</i>	-	0.9	7.5	7.1	-	0.2	0.9	1.8	-	1.1	15.4	8.9	-3.510	**
<i>Cirsium arvense</i>	6.6	10.5	8.4	7.3	5.7	7.2	7.3	1.5	12.3	17.7	15.7	8.8	-4.074	**
<i>Polygonum</i> spp.	8.7	0.6	1.1	3.6	3.2	0.3	0.1	1.2	11.9	0.9	1.2	4.8	4.172	**
<i>Saturnia fabae</i>	11.6	2.4	5.0	1.6	23.9	0.4	1.4	2.4	35.5	2.8	6.4	3.9	-1.833	NS
<i>Solanum carolinense</i>	11.9	14.3	3.8	1.8	12.9	9.8	0.5	0.1	2.6	24.1	4.3	1.9	-1.943	*
<i>Conyza canadensis</i>	2.3	14.9	7.2	0.2	0.3	44.8	8.9	0.0	2.6	59.7	16.1	0.3	-7.818	**
<i>Ribes</i> sp.-4	11.9	4.8	-	-	28.9	3.2	-	-	40.8	8.0	-	-	-	-
<i>Anthrum strumarium</i> ⁵	5.3	0.9	0.5	-	1.7	0.1	0.1	-	7.0	1.0	0.6	-	-	-

Dominant ground cover is defined in Table 6.2-4.

1 Dominant ground cover is defined in Table 0.2-4.
2 dx = significant at 0.1 level. * = significant at 0.05 level, NS = not significant.

2 for = significant at 0.1 level, * = significant at 0.05 level, ** = significant at 0.01 level, *** = significant at 0.001 level.

3 Ranex acetosella not reported in 1973.
4 Ranex sp. not reported in 1975 or 1976.

4 Bildens sp. not reported in 1975 or 1976.

Tree seedlings in the field near the south ponds on Three Mile Island, 1976.

Species	Common Name	Seedlings/Acre		
		1973	1975	1976
<i>Acer saccharinum</i>	silver maple	0.0	0.0	68.2
<i>Fraxinus</i> sp.	ash	0.0	0.0	22.7

Table 6.2-20

Ground cover occurring in the north field on Three Mile Island, 1976.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
<u>Aster ericoides</u>	heath aster	8.76	38.36	47.12
Grass (unidentified)	grass	10.24	22.14	32.38
<u>Solidago</u> spp.	goldenrod	9.61	8.35	17.96
Non-vegetative cover	-	7.50	8.13	15.63
<u>Rubus</u> spp.	blackberry	5.70	6.52	12.02
<u>Lonicera japonica</u>	Japanese honeysuckle	1.80	2.98	4.78
<u>Acalypha virginica</u>	three-seeded mercury	3.70	0.31	4.01
<u>Polygonum scandens</u>	false buckwheat	2.64	0.88	3.52
<u>Desmodium</u> sp.	tick trefoil	2.53	0.94	3.47
<u>Oenothera biennis</u>	evening primrose	2.96	0.42	3.33
<u>Vitis</u> sp.	grape	1.80	1.01	2.81
<u>Eupatorium perfoliatum</u>	boneset	2.32	0.41	2.73
<u>Solanum carolinense</u>	horse nettle	2.43	0.21	2.64
<u>Impatiens</u> sp.	jewelweed	1.80	0.70	2.50
<u>Eupatorium rugosum</u>	white snakeroot	1.69	0.73	2.42
<u>Setaria faberii</u>	nodding foxtail	1.69	0.65	2.34
<u>Rumex al.</u>	water dock	2.11	0.17	2.28
<u>Juncus</u> spp.	rush	1.48	0.65	2.13
<u>Verbesina alternifolia</u>	wingstem	1.69	0.44	2.13
<u>Melilotus alba</u>	white sweet clover	0.84	1.22	2.06
<u>Artemisia artemisiifolia</u>	ragweed	1.37	0.37	1.74
<u>Cassia fasciculata</u>	partridge pea	1.27	0.34	1.61
<u>Bidens</u> spp.	bur marigold	1.16	0.36	1.52
<u>Fragaria virginiana</u>	strawberry	1.27	0.19	1.46
<u>Oxalis</u> sp.	wood sorrel	1.27	0.10	1.37
<u>Viola</u> sp.	violet	1.06	0.15	1.21
<u>Pilea pumila</u>	clearweed	0.95	0.20	1.15
<u>Lysimachia vulgaris</u>	loosestrife	0.84	0.17	1.01
<u>Apocynum medium</u>	dogbane	0.84	0.06	0.90
<u>Boehreria cylindrica</u>	false nettle	0.63	0.26	0.89
<u>Coreopsis grandiflora</u>	dayflower	0.74	0.13	0.87
<u>Polygonum</u> spp.	smartweed	0.74	0.09	0.83
<u>Lycopus americanus</u>	water horehound	0.74	0.08	0.82
<u>Hypericum punctatum</u>	St. John's wort	0.74	0.06	0.80
<u>Cirsium arvense</u>	Canada thistle	0.74	0.04	0.78
<u>Teucrium canadense</u>	wood sage	0.74	0.04	0.78
<u>Urtica dioica</u>	stinging nettle	0.42	0.30	0.72
<u>Gerardia tenuifolia</u>	gerardia	0.53	0.10	0.63
<u>Rhus radicans</u>	poison ivy	0.53	0.09	0.62
<u>Euphorbia maculata</u>	spurge	0.53	0.08	0.61
<u>Mimulus ringens</u>	monkey flower	0.42	0.16	0.58
<u>Parthenocissus quinquefolia</u>	Virginia creeper	0.53	0.05	0.58
<u>Lypha latifolia</u>	cat-tail	0.42	0.15	0.57
Mustard (unidentified)	mustard	0.53	0.04	0.57
<u>Rumex acetosella</u>	sheep sorrel	0.42	0.10	0.52
<u>Conium maculatum</u>	poison hemlock	0.42	0.05	0.47
<u>Daucus carota</u>	Queen Anne's lace	0.42	0.05	0.47
<u>Lythrum salicaria</u>	purple loosestrife	0.32	0.15	0.47
<u>Ceanothus canadensis</u>	avens	0.42	0.03	0.45
<u>Potentilla recta</u>	cinquefoil	0.42	0.03	0.45

POOR ORIGINAL

1565 276

Table 6.2-23 continued.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
<u>Veronica verticillata</u>	white vervain	0.32	0.11	0.43
<u>Aster multiflorus</u>	New England aster	0.32	0.10	0.42
<u>Eriogonum fasciculatum</u>	spike-rush	0.21	0.15	0.36
<u>Anemone patens</u>	scarlet pimpernel	0.32	0.03	0.35
<u>Hieracium sp.</u>	hawkweed	0.32	0.03	0.35
<u>Scrophularia helvola</u>	wild bean	0.32	0.02	0.34
<u>Pentstemon scabellus</u>	ditch stonecrop	0.21	0.05	0.26
<u>Clematis virginiana</u>	virgin's bower	0.21	0.04	0.25
<u>Allium vineale</u>	wild garlic	0.21	0.02	0.23
<u>Aster sp.</u>	aster	0.21	0.02	0.23
<u>Rosa sp.</u>	wild rose	0.11	0.12	0.23
<u>Asclepias tuberosa</u>	swamp milkweed	0.21	0.01	0.22
<u>Crotalaria parvifolia</u>	rattlebox	0.21	0.01	0.22
<u>Hypericum virginicum</u>	marsh St. John's wort	0.21	0.01	0.22
<u>Taraxacum officinale</u>	dandelion	0.21	0.00	0.21
<u>Pyrola asarifolia</u>	mountain mint	0.11	0.06	0.17
<u>Carex lasiocarpa</u>	sedge	0.11	0.05	0.16
<u>Glechoma hederacea</u>	ground ivy	0.11	0.05	0.16
<u>Cyperus sp.</u>	galingale	0.11	0.03	0.14
<u>Helianthus scaberrimus</u>	sunflower	0.11	0.03	0.14
rosette (unidentified)	unidentified rosette	0.11	0.02	0.13
<u>Celastrus scandens</u>	bittersweet	0.11	0.01	0.12
<u>Eupatorium coelestinum</u>	mistflower	0.11	0.01	0.12
<u>Eupatorium serotinum</u>	throughwort	0.11	0.01	0.12
<u>Oxalis sensilis</u>	sensitive fern	0.11	0.01	0.12
<u>Prunella americana</u>	ground cherry	0.11	0.01	0.12
<u>Trifolium sp.</u>	white clover	0.11	0.01	0.12
<u>Conium maculatum</u>	horseweed	0.11	0.00*	0.11
<u>Plantago rugelii</u>	plantain	0.11	0.00	0.11
<u>Potentilla simplex</u>	cinquefoil	0.11	0.00	0.11
<u>Stellaria media</u>	chickweed	0.11	0.00	0.11

* Indicates less than 0.005 percent relative coverage.

POOR ORIGINAL

1565 277

Table 6.2-21

Comparison of the dominant ground cover¹ occurring in the north field on Three Mile Island, 1973-1976.

Species	Relative Frequency				Relative Coverage				Importance Value				1976 vs 1975		1976 vs 1973	
	1973	1974	1975	1976	1973	1974	1975	1976	1973	1974	1975	1976	F	Significance ²	F	Significance
<i>Aster ericoides</i>	12.4	13.1	8.4	8.8	14.2	38.9	37.1	38.4	26.6	52.0	45.5	47.1	-0.047	NS	2.012	*
<i>Galidago</i> spp.	7.5	7.2	8.5	9.6	3.3	5.0	5.8	8.4	11.3	12.2	14.3	18.0	3.973	**	5.077	**
Non-vegetative cover	0.6	0.2	5.4	7.5	0.2	0.0	5.4	8.1	0.8	0.2	10.8	15.6	1.972	NS	9.124	**
<i>Rubus</i> spp.	6.6	8.6	5.7	5.7	7.1	16.8	11.3	6.3	13.7	27.4	17.0	17.0	-4.196	**	-0.972	NS
<i>Desmodium</i> sp.	5.2	5.9	3.6	2.5	3.3	2.4	2.6	0.9	8.5	8.3	6.2	3.5	-7.515	*	-3.609	**
<i>Galium latifolium</i>	5.3	4.5	2.4	2.4	1.7	0.6	0.2	0.2	6.5	5.1	2.6	2.6	-0.251	NS	-5.072	**
<i>Setaria faberii</i>	13.4	9.7	4.0	1.7	20.8	15.4	0.9	0.6	42.2	25.1	4.9	2.3	-2.071	*	-5.090	**
<i>Achroia artemisiifolia</i>	13.9	6.6	2.3	1.4	24.7	4.3	0.3	0.4	38.6	10.9	2.6	1.7	-2.495	NS	-22.616	**
<i>Hypericum punctatum</i>	1.7	5.4	4.8	0.7	0.2	1.6	1.4	0.1	1.9	7.0	6.2	0.8	-6.185	**	-1.722	NS
<i>Coryza canadensis</i> ³	9.2	2.9	-	0.1	7.4	1.4	-	0.0	16.6	4.3	-	0.1	-	-	-	-

¹ Dominant ground cover is defined in Table 6.2-4.² * = significant at 0.01 level, ** = significant at 0.05 level, NS = not significant.³ *Coryza canadensis* was not reported in 1975, and frequency was too low to test in 1976.

POOR ORIGINAL

1565-278

Table 6.2-22

Tree seedlings in the north field on Three Mile Island, 1973-1976.

Species	Common Name	Seedlings/Acre	
		1973	1976
<i>Prunus rubra</i>	red mulberry	0.0	9.9
<i>Prunus serotina</i>	black cherry	9.9	9.9
<i>Rubinia pseudacacia</i>	black locust	0.0	9.9
<i>Ulmus americana</i>	elm	0.0	9.9
<i>Platanus occidentalis</i>	sycamore	108.3	69.3
<i>Acer rubrum</i>	red maple	0.0	39.6
<i>Acer saccharinum</i>	silver maple	19.8	29.7
<i>Betula nigra</i>	river birch	0.0	9.9
<i>Salix</i> sp.	willow	0.0	9.9
<i>Alnus incana</i>	tree-of-heaven	9.9	0.0
<i>Fraxinus</i> sp.	ash	9.9	0.0
<i>Liriodendron tulipifera</i>	tulip tree	9.9	0.0

Table 6.2-23

Ground cover occurring in the field near the 500 kv substation, 1976.

Species	Common Name	Relative Frequency (percent)	Relative Coverage (percent)	Importance Value
<u>Aster ericoides</u>	heath aster	9.69	37.81	47.70
<u>Solidago spp.</u>	goldenrod	9.66	27.02	36.68
<u>Coronilla varia</u>	crown vetch	3.15	11.00	14.15
<u>Cirsium arvense</u>	Canada thistle	7.42	3.20	10.62
<u>Ranuncus cernuus</u>	Queen Anne's lace	6.97	1.14	8.11
<u>Lonicera japonica</u>	Japanese honeysuckle	2.47	5.54	8.01
Grass	grass	5.39	2.10	7.49
<u>Phleum pratense</u>	timothy	3.82	3.27	7.09
<u>Plantago major</u>	common plantain	5.39	0.35	5.74
Non-vegetative cover	-	4.27	1.05	5.32
<u>Taraxacum officinale</u>	dandelion	4.91	0.36	5.27
<u>Trifolium pratense</u>	red clover	4.27	0.90	5.17
<u>Adonis vernalis</u>	dogbane	4.04	0.93	4.97
<u>Clematis heteropetala</u>	ground ivy	2.02	1.66	3.68
<u>Rhus radicans</u>	poison ivy	2.92	0.38	3.30
<u>Rumex sp.</u>	dock	2.25	0.23	2.48
<u>Plantago lanceolata</u>	English plantain	2.25	0.20	2.45
<u>Gem canadense</u>	avens	1.80	0.12	1.92
<u>Polygonum pennsylvanicum</u>	smartweed	1.35	0.51	1.86
<u>Solanum carolinense</u>	horse nettle	1.57	0.21	1.78
<u>Rhus spp.</u>	blackberry	1.12	0.57	1.69
<u>Polygonum scandens</u>	false buckwheat	1.57	0.09	1.66
<u>Prunella vulgaris</u>	self-heal	1.35	1.11	2.46
<u>Vicia sp.</u>	grape	1.12	0.17	1.29
<u>Physalis longifolia</u>	ground cherry	1.12	0.11	1.23
<u>Lactuca pteris</u>	orchard grass	0.67	0.51	1.18
Mustard (unidentified)	mustard	1.12	0.05	1.17
<u>Cirsium vulgare</u>	bull thistle	0.90	0.05	0.95
<u>Lyceum virginicum</u>	bugleweed	0.67	0.07	0.74
<u>Oenothera biennis</u>	evening primrose	0.67	0.06	0.73
<u>Achrosia trifida</u>	giant ragweed	0.67	0.05	0.72
<u>Achrosia artemisiifolia</u>	ragweed	0.67	0.03	0.70
<u>Achillea millefolium</u>	yarrow	0.45	0.03	0.48
<u>Allium vineale</u>	wild garlic	0.45	0.02	0.47
<u>Oxalis sp.</u>	wood sorrel	0.45	0.02	0.47
<u>Agrimonia eupatoria</u>	agrimony	0.22	0.03	0.25
<u>Hieracium sp.</u>	hawkweed	0.22	0.01	0.23
<u>Parthenocissus quinquefolia</u>	Virginia creeper	0.22	0.01	0.23
<u>Viola sp.</u>	violet	0.22	0.01	0.23
<u>Rumex acetosella</u>	sheep sorrel	0.22	0.00*	0.22

* Indicates less than 0.005 percent relative coverage.

1565 279

POOR ORIGINAL

Table 6.2-24

Comparison of the dominant ground cover¹ occurring in the field near the 500 kv substation on Route 441², 1974-1976.

Species	Relative Frequency		Relative Coverage		Importance Value		1976 vs 1975		1976 vs 1974	
	1974	1975	1974	1975	1974	1975	r	Significance	r	Significance
<i>Aster ageratoides</i>	12.4	9.2	9.9	44.9	68.3	37.8	57.3	57.5	47.7	0.235
<i>Colidaga</i> spp.	7.7	8.7	9.7	2.5	10.4	27.0	10.2	19.1	36.7	2.102
<i>Coronilla varia</i>	0.5	4.1	3.2	0.2	16.1	11.0	0.7	29.2	15.2	-0.621
<i>Lirioden arvensis</i>	6.1	6.2	7.6	2.6	2.9	1.2	8.7	9.1	10.6	1.719
<i>Prunus americana</i>	2.8	6.0	7.0	1.5	1.6	1.1	6.3	7.6	8.1	0.206
<i>Plantago patens</i>	2.2	6.7	3.8	1.6	4.3	3.3	3.8	11.0	7.1	-1.618
<i>Loranthum officinale</i>	10.7	6.0	5.4	10.0	1.2	0.4	20.7	7.2	5.7	-2.292
<i>Epilobium prostratum</i>	7.6	4.8	4.9	2.1	1.0	0.4	9.5	5.8	5.1	-1.592
<i>Alnus incana</i>	12.7	3.2	4.3	21.3	0.4	0.9	16.0	1.6	5.2	1.521
<i>Conoclinium ovatum</i>	1.9	3.7	4.0	0.5	1.2	0.9	2.4	4.9	5.0	0.273
	1.0	4.4	-	0.4	0.4	-	1.4	6.8	-	1.719

¹ Dominant ground cover is defined in Table 6.2-4.² This field was not sampled in 1973.³ r = significant at 0.01 level, s = significant at 0.05 level, NS = not significant.⁴ Conoclinium ovatum not reported in 1976.

Table 6.2-25

Tree seedlings in the field near the 500 kv substation, 1974-1976.

Species	Common Name	Seedlings/Acre	
		1974	1975
<i>Pinus strobus</i>	white pine	217.4	260.9
<i>Acer rubrum</i>	red maple	0.0	0.0
<i>Acer negundo</i>	box elder	2.2	0.0
<i>Acer saccharum</i>	sugar maple	2.2	0.0

1565 280